

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2004/026930

International filing date: 20 August 2004 (20.08.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 10/720,448
Filing date: 24 November 2003 (24.11.2003)

Date of receipt at the International Bureau: 14 May 2007 (14.05.2007)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1608340

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

May 09, 2007

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 10/720,448

FILING DATE: November 24, 2003

RELATED PCT APPLICATION NUMBER: PCT/US04/26930

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS US10/720,448



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

UTILITY PATENT APPLICATION TRANSMITTAL

Submit an original and a duplicate for fee processing
(Only for new nonprovisional applications under 37 CFR 1.53(b))

ADDRESS TO:

Mail Stop Patent Application
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Attorney Docket No. 03-465-B (400.138)
First Named Inventor James McSwiggen
Express Mail No. EV 334703995 US

APPLICATION ELEMENTS

1. ☒ Transmittal Form with Fee
2. ☒ Specification (including tables, claims and abstract) [Total Pages 242]
3. ☒ Drawings [Total Sheets 95]
4. ☐ Oath or Declaration [Total Pages]
 - a. ☐ Newly executed
 - b. ☐ Copy from prior application

[Note Boxes 5 and 18 below]

 - i. ☐ Deletion of Inventor(s) Signed statement attached deleting inventor(s) named in the prior application
5. ☐ Incorporation by Reference: The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
6. ☐ Computer Code Listing (See 1.96)
 - a. ☐ Microfiche Appendix
 - b. ☐ CD-Rom (in duplicate, with separate transmittal)
7. ☐ Nucleotide and/or Amino Acid Sequence Submission
 - a. ☐ Computer Readable Copy
 - b. ☐ Paper Copy
 - c. ☐ Statement verifying above copies
8. ☒ Small Entity Status
 - ☐ is claimed
 - ☒ Statement filed in prior application; status still proper and desired
 - ☐ is no longer claimed.

ACCOMPANYING APPLICATION PARTS

9. ☐ Assignment Papers
10. ☐ Power of Attorney
11. ☐ English Translation Document (if applicable)
12. ☐ Information Disclosure Statement (IDS)
 - ☐ PTO-1449 Form
 - ☐ Copies of IDS Citations
13. ☐ Preliminary Amendment
14. ☒ Return Receipt Postcard (Should be specifically itemized)
15. ☐ Certified Copy of Priority Document(s)
16. ☐ A Request for non-publication pursuant to 35 U.S.C. § 122(b)(2)(B)(i)
17. ☒ Other: Patent Application Date Sheet (12 pages)

18. ☒ This is a CONTINUING APPLICATION. Please note the following:

a. ☒ This is a ☐ Continuation ☐ Divisional ☒ Continuation-in-part of prior U.S. Patent Application Serial Nos. 10/693,095 filed October 23, 2003 which is a continuation-in-part of 10/444,853, filed May 23, 2003, which is a continuation-in-part of 10/652,791, filed August 29, 2003, which is a continuation of 10/422,704, filed April 24, 2003, which is a continuation of U.S. Patent Application No. 10/417,012, filed April 16, 2003. This application is also a continuation-in-part of International Patent Application No. PCT/US03/05346, filed February 20, 2003, and a continuation-in-part of International Patent Application No. PCT/US03/05028, filed February 20, 2003, both of which claim the benefit of U.S. Provisional Application No. 60/358,580 filed February 20, 2002, U.S. Provisional Application No. 60/363,124 filed March 11, 2002, U.S. Provisional Application No. 60/386,782 filed June 6, 2002, U.S. Provisional Application No. 60/406,784 filed August 29, 2002, U.S. Provisional Application No. 60/408,378 filed September 5, 2002, U.S. Provisional Application No. 60/409,293 filed September 9, 2002, and U.S. Provisional Application No. 60/440,129 filed January 15, 2003. U.S. Serial No. 10/693,095 is also a continuation-in-part of US Patent Application No. 10/427,160, filed April 30, 2003 and International Patent Application No. PCT/US02/15876 filed May 17, 2002.

- b. ☐ Cancel in this application original claims of the prior application before calculating the filing fee. (At least one claim must remain.)

<p>c. <input type="checkbox"/> Amend the specification by inserting before the first line the sentence: This is a <input type="checkbox"/> continuation <input type="checkbox"/> divisional <input type="checkbox"/> continuation-in-part of of application Serial No.</p> <p>d. <input checked="" type="checkbox"/> The prior application is assigned of record to Sirna Therapeutics, Inc.</p>
--

UTILITY PATENT APPLICATION TRANSMITTAL	Attorney Docket No. 03-465-A (400.136)
---	--

APPLICATION FEES				
BASIC FEE				\$ 770.00
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total Claims	17 -20 =		x \$18.00	\$
Independent Claims	1 - 3 =		x \$84.00	\$.00
<input type="checkbox"/> Multiple Dependent Claims(s) if applicable			+\$280.00	\$
Total of above calculations =				\$
Reduction by 50% for filing by small entity =				\$(385.00)
<input type="checkbox"/> Assignment fee if applicable			+ \$40.00	\$
TOTAL =				\$ 385.00

19. ☐ Please charge my Deposit Account No. 13-2490 in the amount of \$385.00
20. ☒ A check in the amount of \$385.00 is enclosed.
21. The Commissioner is hereby authorized to credit overpayments or charge any additional fees of the following types to Deposit Account No. 13-2490:
- a. ☒ Fees required under 37 CFR 1.16.
 - b. ☒ Fees required under 37 CFR 1.17.
 - c. ☒ Fees required under 37 CFR 1.18.
22. The Commissioner is hereby generally authorized under 37 CFR 1.136(a)(3) to treat any future reply in this or any related application filed pursuant to 37 CFR 1.53 requiring an extension of time as incorporating a request therefor, and the Commissioner is hereby specifically authorized to charge Deposit Account No. 13-2490 for any fee that may be due in connection with such a request for an extension of time.

23. CERTIFICATE OF MAILING

I hereby certify that, under 37 CFR § 1.10, I directed that the correspondence identified above be deposited with the United States Postal Service as "Express Mail Post Office to Addressee," addressed to Mail Stop Patent Application, Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450, on the date indicated below.


24. USPTO CUSTOMER NUMBER

20306

PATENT TRADEMARK OFFICE

McDonnell Boehnen Hulbert & Berghoff

25. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED

Name	Michael S. Greenfield
Reg. No.	37,142
Signature	
Date	November 24, 2003

**RNA INTERFERENCE MEDIATED INHIBITION OF GENE EXPRESSION
USING CHEMICALLY MODIFIED SHORT INTERFERING NUCLEIC ACID
(siNA)**

This application is a continuation-in-part of U.S. Patent Application No. 10/693,095, filed October 23, 2003, which is a continuation-in-part of U.S. Patent Application No. 10/444,853, filed May 23, 2003 which is a continuation-in-part of 10/652,791, filed August 29, 2003, which is a continuation of 10/422,704, filed April 24, 2003, which is a continuation of U.S. Patent Application No. 10/417,012, filed April 16, 2003. U.S. application 10/693,095 is also a continuation-in-part of International Patent Application No. PCT/US03/05346, filed February 20, 2003, and a continuation-in-part of International Patent Application No. PCT/US03/05028, filed February 20, 2003, both of which claim the benefit of U.S. Provisional Application No. 60/358,580 filed February 20, 2002, U.S. Provisional Application No. 60/363,124 filed March 11, 2002, U.S. Provisional Application No. 60/386,782 filed June 6, 2002, U.S. Provisional Application No. 60/406,784 filed August 29, 2002, U.S. Provisional Application No. 60/408,378 filed September 5, 2002, U.S. Provisional Application No. 60/409,293 filed September 9, 2002, and U.S. Provisional Application No. 60/440,129 filed January 15, 2003. U.S. application 10/693,095 is also a continuation-in-part of US Patent Application No. 10/427,160, filed April 30, 2003 and International Patent Application No. PCT/US02/15876 filed May 17, 2002. The instant application claims the benefit of all the listed applications, which are hereby incorporated by reference herein in their entireties, including the drawings.

Field Of The Invention

The present invention concerns methods and reagents useful in modulating gene expression in a variety of applications, including use in therapeutic, diagnostic, target validation, and genomic discovery applications. Specifically, the invention relates to synthetic small nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules capable of mediating RNA interference (RNAi).

Background Of The Invention

The following is a discussion of relevant art pertaining to RNAi. The discussion is provided only for understanding of the invention that follows. The summary is not an admission that any of the work described below is prior art to the claimed invention.

5 Applicant demonstrates herein that chemically modified short interfering nucleic acids possess the same capacity to mediate RNAi as do siRNA molecules and are expected to possess improved stability and activity in vivo; therefore, this discussion is not meant to be limiting only to siRNA and can be applied to siNA as a whole.

10 RNA interference refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Zamore *et al.*, 2000, *Cell*, 101, 25-33; Fire *et al.*, 1998, *Nature*, 391, 806; Hamilton *et al.*, 1999, *Science*, 286, 950-951). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an
15 evolutionarily-conserved cellular defense mechanism used to prevent the expression of foreign genes and is commonly shared by diverse flora and phyla (Fire *et al.*, 1999, *Trends Genet.*, 15, 358). Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or from the random integration of transposon elements into a host genome via a
20 cellular response that specifically destroys homologous single-stranded RNA or viral genomic RNA. The presence of dsRNA in cells triggers the RNAi response through a mechanism that has yet to be fully characterized. This mechanism appears to be different from the interferon response that results from dsRNA-mediated activation of protein kinase PKR and 2',5'-oligoadenylate synthetase resulting in non-specific cleavage of
25 mRNA by ribonuclease L.

The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Hamilton *et al.*, *supra*; Zamore *et al.*, 2000, *Cell*, 101, 25-33; Bernstein *et al.*, 2001, *Nature*, 409, 363). Short
30 interfering RNAs derived from dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes (Hamilton *et al.*, *supra*;

Elbashir *et al.*, 2001, *Genes Dev.*, 15, 188). Dicer has also been implicated in the excision of 21- and 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner *et al.*, 2001, *Science*, 293, 834). The RNAi response also features an endonuclease complex,
 5 commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having sequence complementary to the antisense strand of the siRNA duplex. Cleavage of the target RNA takes place in the middle of the region complementary to the antisense strand of the siRNA duplex (Elbashir *et al.*, 2001, *Genes Dev.*, 15, 188).

10 RNAi has been studied in a variety of systems. Fire *et al.*, 1998, *Nature*, 391, 806, were the first to observe RNAi in *C. elegans*. Bahramian and Zarbl, 1999, *Molecular and Cellular Biology*, 19, 274-283 and Wianny and Goetz, 1999, *Nature Cell Biol.*, 2, 70, describe RNAi mediated by dsRNA in mammalian systems. Hammond *et al.*, 2000, *Nature*, 404, 293, describe RNAi in *Drosophila* cells transfected with dsRNA. Elbashir
 15 *et al.*, 2001, *Nature*, 411, 494, describe RNAi induced by introduction of duplexes of synthetic 21-nucleotide RNAs in cultured mammalian cells including human embryonic kidney and HeLa cells. Recent work in *Drosophila* embryonic lysates (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877) has revealed certain requirements for siRNA length, structure, chemical composition, and sequence that are essential to mediate efficient RNAi activity.
 20 These studies have shown that 21-nucleotide siRNA duplexes are most active when containing 3'-terminal dinucleotide overhangs. Furthermore, complete substitution of one or both siRNA strands with 2'-deoxy (2'-H) or 2'-O-methyl nucleotides abolishes RNAi activity, whereas substitution of the 3'-terminal siRNA overhang nucleotides with 2'-deoxy nucleotides (2'-H) was shown to be tolerated. Single mismatch sequences in the
 25 center of the siRNA duplex were also shown to abolish RNAi activity. In addition, these studies also indicate that the position of the cleavage site in the target RNA is defined by the 5'-end of the siRNA guide sequence rather than the 3'-end of the guide sequence (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). Other studies have indicated that a 5'-phosphate on the target-complementary strand of a siRNA duplex is required for siRNA
 30 activity and that ATP is utilized to maintain the 5'-phosphate moiety on the siRNA (Nykanen *et al.*, 2001, *Cell*, 107, 309).

Studies have shown that replacing the 3'-terminal nucleotide overhanging segments of a 21-mer siRNA duplex having two-nucleotide 3'-overhangs with deoxyribonucleotides does not have an adverse effect on RNAi activity. Replacing up to four nucleotides on each end of the siRNA with deoxyribonucleotides has been reported to be well tolerated, whereas complete substitution with deoxyribonucleotides results in no RNAi activity (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). In addition, Elbashir *et al.*, *supra*, also report that substitution of siRNA with 2'-O-methyl nucleotides completely abolishes RNAi activity. Li *et al.*, International PCT Publication No. WO 00/44914, and Beach *et al.*, International PCT Publication No. WO 01/68836 preliminarily suggest that siRNA may include modifications to either the phosphate-sugar backbone or the nucleoside to include at least one of a nitrogen or sulfur heteroatom, however, neither application postulates to what extent such modifications would be tolerated in siRNA molecules, nor provides any further guidance or examples of such modified siRNA. Kreutzer *et al.*, Canadian Patent Application No. 2,359,180, also describe certain chemical modifications for use in dsRNA constructs in order to counteract activation of double-stranded RNA-dependent protein kinase PKR, specifically 2'-amino or 2'-O-methyl nucleotides, and nucleotides containing a 2'-O or 4'-C methylene bridge. However, Kreutzer *et al.* similarly fails to provide examples or guidance as to what extent these modifications would be tolerated in siRNA molecules.

Parrish *et al.*, 2000, *Molecular Cell*, 6, 1077-1087, tested certain chemical modifications targeting the unc-22 gene in *C. elegans* using long (>25 nt) siRNA transcripts. The authors describe the introduction of thiophosphate residues into these siRNA transcripts by incorporating thiophosphate nucleotide analogs with T7 and T3 RNA polymerase and observed that RNAs with two phosphorothioate modified bases also had substantial decreases in effectiveness as RNAi. Further, Parrish *et al.* reported that phosphorothioate modification of more than two residues greatly destabilized the RNAs *in vitro* such that interference activities could not be assayed. *Id.* at 1081. The authors also tested certain modifications at the 2'-position of the nucleotide sugar in the long siRNA transcripts and found that substituting deoxynucleotides for ribonucleotides produced a substantial decrease in interference activity, especially in the case of Uridine to Thymidine and/or Cytidine to deoxy-Cytidine substitutions. *Id.* In addition, the authors tested certain base modifications, including substituting, in sense and antisense

strands of the siRNA, 4-thiouracil, 5-bromouracil, 5-iodouracil, and 3-(aminoallyl)uracil for uracil, and inosine for guanosine. Whereas 4-thiouracil and 5-bromouracil substitution appeared to be tolerated, Parrish reported that inosine produced a substantial decrease in interference activity when incorporated in either strand. Parrish also reported
 5 that incorporation of 5-iodouracil and 3-(aminoallyl)uracil in the antisense strand resulted in a substantial decrease in RNAi activity as well.

The use of longer dsRNA has been described. For example, Beach *et al.*, International PCT Publication No. WO 01/68836, describes specific methods for attenuating gene expression using endogenously-derived dsRNA. Tuschl *et al.*,
 10 International PCT Publication No. WO 01/75164, describe a *Drosophila in vitro* RNAi system and the use of specific siRNA molecules for certain functional genomic and certain therapeutic applications; although Tuschl, 2001, *Chem. Biochem.*, 2, 239-245, doubts that RNAi can be used to cure genetic diseases or viral infection due to the danger of activating interferon response. Li *et al.*, International PCT Publication No. WO
 15 00/44914, describe the use of specific dsRNAs for attenuating the expression of certain target genes. Zernicka-Goetz *et al.*, International PCT Publication No. WO 01/36646, describe certain methods for inhibiting the expression of particular genes in mammalian cells using certain dsRNA molecules. Fire *et al.*, International PCT Publication No. WO
 20 99/32619, describe particular methods for introducing certain dsRNA molecules into cells for use in inhibiting gene expression. Plaetinck *et al.*, International PCT Publication No. WO 00/01846, describe certain methods for identifying specific genes responsible for conferring a particular phenotype in a cell using specific dsRNA molecules. Mello *et al.*, International PCT Publication No. WO 01/29058, describe the identification of specific
 25 genes involved in dsRNA-mediated RNAi. Deschamps Depaillette *et al.*, International PCT Publication No. WO 99/07409, describe specific compositions consisting of particular dsRNA molecules combined with certain anti-viral agents. Waterhouse *et al.*, International PCT Publication No. 99/53050, describe certain methods for decreasing the phenotypic expression of a nucleic acid in plant cells using certain dsRNAs. Driscoll *et al.*, International PCT Publication No. WO 01/49844, describe specific DNA constructs
 30 for use in facilitating gene silencing in targeted organisms.

Others have reported on various RNAi and gene-silencing systems. For example, Parrish *et al.*, 2000, *Molecular Cell*, 6, 1077-1087, describe specific chemically-modified

siRNA constructs targeting the *unc-22* gene of *C. elegans*. Grossniklaus, International PCT Publication No. WO 01/38551, describes certain methods for regulating polycomb gene expression in plants using certain dsRNAs. Churikov *et al.*, International PCT Publication No. WO 01/42443, describe certain methods for modifying genetic characteristics of an organism using certain dsRNAs. Cogoni *et al.*, International PCT Publication No. WO 01/53475, describe certain methods for isolating a *Neurospora* silencing gene and uses thereof. Reed *et al.*, International PCT Publication No. WO 01/68836, describe certain methods for gene silencing in plants. Honer *et al.*, International PCT Publication No. WO 01/70944, describe certain methods of drug screening using transgenic nematodes as Parkinson's Disease models using certain dsRNAs. Deak *et al.*, International PCT Publication No. WO 01/72774, describe certain *Drosophila*-derived gene products that may be related to RNAi in *Drosophila*. Arndt *et al.*, International PCT Publication No. WO 01/92513 describe certain methods for mediating gene suppression by using factors that enhance RNAi. Tuschl *et al.*, International PCT Publication No. WO 02/44321, describe certain synthetic siRNA constructs. Pachuk *et al.*, International PCT Publication No. WO 00/63364, and Satishchandran *et al.*, International PCT Publication No. WO 01/04313, describe certain methods and compositions for inhibiting the function of certain polynucleotide sequences using certain dsRNAs. Echeverri *et al.*, International PCT Publication No. WO 02/38805, describe certain *C. elegans* genes identified via RNAi. Kreutzer *et al.*, International PCT Publications Nos. WO 02/055692, WO 02/055693, and EP 1144623 B1 describes certain methods for inhibiting gene expression using RNAi. Graham *et al.*, International PCT Publications Nos. WO 99/49029 and WO 01/70949, and AU 4037501 describe certain vector expressed siRNA molecules. Fire *et al.*, US 6,506,559, describe certain methods for inhibiting gene expression in vitro using certain long dsRNA (greater than 25 nucleotide) constructs that mediate RNAi.

SUMMARY OF THE INVENTION

This invention relates to compounds, compositions, and methods useful for modulating RNA function and/or gene expression in a cell. Specifically, the instant invention features synthetic small nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-

RNA (miRNA), and short hairpin RNA (shRNA) molecules capable of modulating gene expression in cells by RNA interference (RNAi). The siNA molecules of the invention can be chemically modified. The use of chemically modified siNA can improve various properties of native siRNA molecules through increased resistance to nuclease degradation *in vivo* and/or improved cellular uptake. The chemically modified siNA molecules of the instant invention provide useful reagents and methods for a variety of therapeutic, diagnostic, agricultural, target validation, genomic discovery, genetic engineering and pharmacogenomic applications.

In a non-limiting example, the introduction of chemically modified nucleotides into nucleic acid molecules provides a powerful tool in overcoming potential limitations of *in vivo* stability and bioavailability inherent to native RNA molecules that are delivered exogenously. For example, the use of chemically modified nucleic acid molecules can enable a lower dose of a particular nucleic acid molecule for a given therapeutic effect since chemically modified nucleic acid molecules tend to have a longer half-life in serum. Furthermore, certain chemical modifications can improve the bioavailability of nucleic acid molecules by targeting particular cells or tissues and/or improving cellular uptake of the nucleic acid molecule. Therefore, even if the activity of a chemically modified nucleic acid molecule is reduced as compared to a native nucleic acid molecule, for example when compared to an all RNA nucleic acid molecule, the overall activity of the modified nucleic acid molecule can be greater than the native molecule due to improved stability and/or delivery of the molecule. Unlike native unmodified siRNA, chemically modified siNA can also minimize the possibility of activating interferon activity in humans.

In one embodiment, the nucleic acid molecules of the invention that act as mediators of the RNA interference gene silencing response are chemically modified double stranded nucleic acid molecules. As in their native double stranded RNA counterparts, these siNA molecules typically consist of duplexes containing about 19 base pairs between oligonucleotides comprising about 19 to about 25 nucleotides. The most active siRNA molecules are thought to have such duplexes with overhanging ends of 1-3 nucleotides, for example 21 nucleotide duplexes with 19 base pairs and 2 nucleotide 3'-overhangs. These overhanging segments are readily hydrolyzed by endonucleases *in vivo*. Studies have shown that replacing the 3'-overhanging segments of a 21-mer siRNA

duplex having 2 nucleotide 3' overhangs with deoxyribonucleotides does not have an adverse effect on RNAi activity. Replacing up to 4 nucleotides on each end of the siRNA with deoxyribonucleotides has been reported to be well tolerated whereas complete substitution with deoxyribonucleotides results in no RNAi activity (Elbashir et al., 2001, EMBO J., 20, 6877). In addition, Elbashir *et al., supra*, also report that substitution of siRNA with 2'-O-methyl nucleotides completely abolishes RNAi activity. Li *et al.*, International PCT Publication No. WO 00/44914, and Beach *et al.*, International PCT Publication No. WO 01/68836 both suggest that siRNA may include modifications to either the phosphate-sugar back bone or the nucleoside to include at least one of a nitrogen or sulfur heteroatom, however neither application teaches to what extent these modifications are tolerated in siRNA molecules nor provide any examples of such modified siRNA. Kreutzer and Limmer, Canadian Patent Application No. 2,359,180, also describe certain chemical modifications for use in dsRNA constructs in order to counteract activation of double stranded-RNA-dependent protein kinase PKR, specifically 2'-amino or 2'-O-methyl nucleotides, and nucleotides containing a 2'-O or 4'-C methylene bridge. However, Kreutzer and Limmer similarly fail to show to what extent these modifications are tolerated in siRNA molecules nor provide any examples of such modified siRNA.

In one embodiment, the invention features chemically modified siNA constructs having specificity for target nucleic acid molecules in a cell. Non-limiting examples of such chemical modifications include without limitation phosphorothioate internucleotide linkages, 2'-O-methyl ribonucleotides, 2'-deoxy-2'-fluoro ribonucleotides, 2'-deoxy ribonucleotides, "universal base" nucleotides, 5-C-methyl nucleotides, and inverted deoxyabasic residue incorporation. These chemical modifications, when used in various siNA constructs, are shown to preserve RNAi activity in cells while at the same time, dramatically increasing the serum stability of these compounds. Furthermore, contrary to the data published by Parrish *et al., supra*, applicant demonstrates that multiple (greater than one) phosphorothioate substitutions are well-tolerated and confer substantial increases in serum stability for modified siNA constructs.

In one embodiment, the chemically-modified siNA molecules of the invention comprise a duplex having two strands, one or both of which can be chemically-modified, wherein each strand is about 19 to about 29 (e.g., about 19, 20, 21, 22, 23, 24, 25, 26, 27,

28, or 29) (e.g., about 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, or 29) nucleotides. In one embodiment, the chemically-modified siNA molecules of the invention comprise a duplex having two strands, one or both of which can be chemically-modified, wherein each strand is about 19 to about 23 (e.g., about 19, 20, 21, 22, or 23) nucleotides. In one
5 embodiment, a siNA molecule of the invention comprises modified nucleotides while maintaining the ability to mediate RNAi. The modified nucleotides can be used to improve *in vitro* or *in vivo* characteristics such as stability, activity, and/or bioavailability. For example, a siNA molecule of the invention can comprise modified nucleotides as a percentage of the total number of nucleotides present in the siNA molecule. As such, a
10 siNA molecule of the invention can generally comprise modified nucleotides from about 5 to about 100% of the nucleotide positions (e.g., 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% of the nucleotide positions). The actual percentage of modified nucleotides present in a given siNA molecule depends on the total number of nucleotides present in the siNA. If the
15 siNA molecule is single stranded, the percent modification can be based upon the total number of nucleotides present in the single stranded siNA molecules. Likewise, if the siNA molecule is double stranded, the percent modification can be based upon the total number of nucleotides present in the sense strand, antisense strand, or both the sense and antisense strands. In addition, the actual percentage of modified nucleotides present in a
20 given siNA molecule can also depend on the total number of purine and pyrimidine nucleotides present in the siNA, for example, wherein all pyrimidine nucleotides and/or all purine nucleotides present in the siNA molecule are modified.

The antisense region of a siNA molecule of the invention can comprise a phosphorothioate internucleotide linkage at the 3'-end of said antisense region. The
25 antisense region can comprise about one to about five phosphorothioate internucleotide linkages at the 5'-end of said antisense region. The 3'-terminal nucleotide overhangs of a siNA molecule of the invention can comprise ribonucleotides or deoxyribonucleotides that are chemically-modified at a nucleic acid sugar, base, or backbone. The 3'-terminal nucleotide overhangs can comprise one or more universal base ribonucleotides. The 3'-
30 terminal nucleotide overhangs can comprise one or more acyclic nucleotides.

In one embodiment, a siNA molecule of the invention comprises blunt ends, i.e., the ends do not include any overhanging nucleotides. For example, a siNA molecule of the

invention comprising modifications described herein (e.g., comprising nucleotides having Formulae I-VII or siNA constructs comprising Stab1-Stab18 or any combination thereof) and/or any length described herein can comprise blunt ends or ends with no overhanging nucleotides.

5 In one embodiment, any siNA molecule of the invention can comprise one or more blunt ends, i.e. where a blunt end does not have any overhanging nucleotides. In a non-limiting example, a blunt ended siNA molecule has a number of base pairs equal to the number of nucleotides present in each strand of the siNA molecule. In another example, a siNA molecule comprises one blunt end, for example wherein the 5'-end of the
10 antisense strand and the 3'-end of the sense strand do not have any overhanging nucleotides. In another example, a siNA molecule comprises one blunt end, for example wherein the 3'-end of the antisense strand and the 5'-end of the sense strand do not have any overhanging nucleotides. In another example, a siNA molecule comprises two blunt ends, for example wherein the 3'-end of the antisense strand and the 5'-end of the sense strand as well as the 5'-end of the antisense strand and 3'-end of the sense strand do not
15 have any overhanging nucleotides. A blunt ended siNA molecule can comprise, for example, from about 18 to about 30 nucleotides (e.g., about 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides). Other nucleotides present in a blunt ended siNA molecule can comprise mismatches, bulges, loops, or wobble base pairs, for example, to
20 modulate the activity of the siNA molecule to mediate RNA interference.

By "blunt ends" is meant symmetric termini or termini of a double stranded siNA molecule having no overhainging nucleotides. The two strands of a double stranded siNA molecule align with each other without over-hanging nucleotides at the termini. For example, a blunt ended siNA construct comprises terminal nucleotides that are
25 complimentary between the sense and antisense regions of the siNA molecule.

In one embodiment, the invention features the use of a double-stranded short interfering nucleic acid (siNA) molecule to down-regulate expression of a target gene, wherein the siNA molecule comprises one or more chemical modifications and each strand of the double-stranded siNA is about 19 to about 23 nucleotides (e.g., about 19, 20,
30 21, 22, or 23 nucleotides) long.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a target gene, wherein the siNA molecule comprises no ribonucleotides and each strand of the double-stranded siNA comprises about 19 to about 23 nucleotides.

5 In one embodiment, one of the strands of a double-stranded siNA molecule of the invention comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of a target gene, and wherein the second strand of a double-stranded siNA molecule comprises a nucleotide sequence substantially similar to the nucleotide sequence or a portion thereof of the target gene.

10 In one embodiment, a siNA molecule of the invention comprises about 19 to about 23 nucleotides, and each strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand.

In one embodiment, a siNA molecule of the invention comprises an antisense region comprising a nucleotide sequence that is complementary to a nucleotide sequence
15 or a portion thereof of a target gene, and the siNA further comprises a sense region, wherein the sense region comprises a nucleotide sequence substantially similar to the nucleotide sequence or a portion thereof of the target gene. The antisense region and the sense region each comprise about 19 to about 23 nucleotides, and the antisense region comprises at least about 19 nucleotides that are complementary to nucleotides of the
20 sense region.

In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region, wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of RNA encoded by a target
25 gene and the sense region comprises a nucleotide sequence that is complementary to the antisense region.

In one embodiment, a siNA molecule of the invention is assembled from two separate oligonucleotide fragments wherein one fragment comprises the sense region and the second fragment comprises the antisense region of the siNA molecule. In another embodiment, the sense region is connected to the antisense region via a linker molecule,
30 which can be a polynucleotide linker or a non-nucleotide linker.

In one embodiment, a siNA molecule of the invention comprises a sense region and antisense region, wherein pyrimidine nucleotides in the sense region comprises 2'-O-methyl pyrimidine nucleotides and purine nucleotides in the sense region comprise 2'-deoxy purine nucleotides. In one embodiment, a siNA molecule of the invention
5 comprises a sense region and antisense region, wherein pyrimidine nucleotides present in the sense region comprise 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein purine nucleotides present in the sense region comprise 2'-deoxy purine nucleotides.

In one embodiment, a siNA molecule of the invention comprises a sense region and antisense region, wherein the pyrimidine nucleotides when present in said antisense
10 region are 2'-deoxy-2'-fluoro pyrimidine nucleotides and the purine nucleotides when present in said antisense region are 2'-O-methyl purine nucleotides.

In one embodiment, a siNA molecule of the invention comprises a sense region and antisense region, wherein the pyrimidine nucleotides when present in said antisense
15 region are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein the purine nucleotides when present in said antisense region comprise 2'-deoxy- purine nucleotides.

In one embodiment, a siNA molecule of the invention comprises a sense region and antisense region, wherein the sense region includes a terminal cap moiety at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the sense region. In another embodiment, the terminal cap moiety is an inverted deoxy abasic moiety.

20 In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a gene. Because many genes can share some degree of sequence homology with each other, siNA molecules can be designed to target a class of genes (and associated receptor or ligand genes) or alternately specific genes by selecting sequences that are either shared amongst different gene targets or alternatively
25 that are unique for a specific gene target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of a RNA sequence having homology between several genes so as to target several genes or gene families (e.g., different gene isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a sequence that is
30 unique to a specific RNA sequence of a specific gene due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

In one embodiment, nucleic acid molecules of the invention that act as mediators of the RNA interference gene silencing response are double-stranded nucleic acid molecules. In another embodiment, the siNA molecules of the invention consist of duplexes containing about 19 base pairs between oligonucleotides comprising about 19 to
5 about 25 (*e.g.*, about 19, 20, 21, 22, 23, 24 or 25) nucleotides. In yet another embodiment, siNA molecules of the invention comprise duplexes with overhanging ends of about 1 to about 3 (*e.g.*, about 1, 2, or 3) nucleotides, for example, about 21-nucleotide duplexes with about 19 base pairs and 3'-terminal mononucleotide, dinucleotide, or trinucleotide overhangs.

10 In one embodiment, the invention features one or more chemically-modified siNA constructs having specificity for nucleic acid molecules that express or encode a protein sequence, such as RNA or DNA encoding a protein sequence. Non-limiting examples of such chemical modifications include without limitation phosphorothioate internucleotide linkages, 2'-deoxyribonucleotides, 2'-O-methyl ribonucleotides, 2'-deoxy-2'-fluoro
15 ribonucleotides, "universal base" nucleotides, "acyclic" nucleotides, 5-C-methyl nucleotides, and terminal glyceryl and/or inverted deoxy abasic residue incorporation. These chemical modifications, when used in various siNA constructs, are shown to preserve RNAi activity in cells while at the same time, dramatically increasing the serum stability of these compounds.

20 In one embodiment, a siNA molecule of the invention does not contain any ribonucleotides. In another embodiment, a siNA molecule of the invention comprises one or more ribonucleotides.

In one embodiment, the invention features the use of compounds or compositions that inhibit the activity of double stranded RNA binding proteins (dsRBPs, see for
25 example Silhavy *et al.*, 2003, *Journal of General Virology*, 84, 975-980). Non-limiting examples of compounds and compositions that can be used to inhibit the activity of dsRBPs include but are not limited to small molecules and nucleic acid aptamers that bind to or interact with the dsRBPs and consequently reduce dsRBP activity and/or siNA molecules that target nucleic acid sequences encoding dsRBPs. The use of such
30 compounds and compositions is expected to improve the activity of siNA molecules in biological systems in which dsRBPs can abrogate or suppress the efficacy of siNA

mediated RNA interference, such as where dsRBPs are expressed during viral infection of a cell to escape RNAi surveillance. Therefore, the use of agents that inhibit dsRBP activity is preferred in those instances where RNA interference activity can be improved via the abrogation or suppression of dsRBP activity. Such anti-dsRBP agents can be administered alone or can be co-administered with siNA molecules of the invention, or
5 can be used to pretreat cells or a subject before siNA administration. In another embodiment, anti-dsRBP agents are used to treat viral infection, such as HCV, HBV, or HIV infection with or without siNA molecules of the invention.

In one embodiment, the invention features a double-stranded short interfering
10 nucleic acid (siNA) molecule that down-regulates expression of a gene, wherein one of the strands of the double-stranded siNA molecule comprises a nucleotide sequence that is complementary to a nucleotide sequence of the gene or RNA encoded by the gene or a portion thereof, and wherein the second strand of the double-stranded siNA molecule comprises a nucleotide sequence substantially similar to the nucleotide sequence of the
15 gene or RNA encoded by the gene or a portion thereof.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a gene, wherein each strand of the siNA molecule comprises about 19 to about 23 nucleotides, and wherein each strand comprises at least about 19 nucleotides that are complementary to the
20 nucleotides of the other strand.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a gene, wherein the siNA molecule comprises an antisense region comprising a nucleotide sequence that is complementary to a nucleotide sequence of the gene or RNA encoded by the gene or a
25 portion thereof, and wherein the siNA further comprises a sense region, wherein the sense region comprises a nucleotide sequence substantially similar to the nucleotide sequence of the gene or RNA encoded by the gene or a portion thereof.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits the expression of a target gene by mediating
30 RNA interference (RNAi) process, wherein the siNA molecule comprises no

ribonucleotides and wherein each strand of the double-stranded siNA molecule comprises about 21 nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits the replication of a virus (e.g, as mammalian virus, plant virus, hepatitis C virus, human immunodeficiency virus, hepatitis B virus, herpes simplex virus, cytomegalovirus, human papilloma virus, respiratory syncytial virus, or influenza virus), wherein the siNA molecule does not require the presence of a ribonucleotide within the siNA molecule for the inhibition of replication of the virus and each strand of the double-stranded siNA molecule comprises about 21 nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a gene, wherein the siNA molecule comprises a sense region and an antisense region and wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of RNA encoded by the gene and the sense region comprises a nucleotide sequence that is complementary to the antisense region or a portion thereof, and wherein the purine nucleotides present in the antisense region comprise 2'-deoxy-purine nucleotides. In another embodiment, the purine nucleotides present in the antisense region comprise 2'-O-methyl purine nucleotides. In either of the above embodiments, the antisense region comprises a phosphorothioate internucleotide linkage at the 3' end of the antisense region. In an alternative embodiment, the antisense region comprises a glyceryl modification at the 3' end of the antisense region. In another embodiment of any of the above described siNA molecules, any nucleotides present in a non-complementary region of the antisense strand (e.g. overhang region) are 2'-deoxy nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a gene, wherein the siNA molecule is assembled from two separate oligonucleotide fragments each comprising 21 nucleotides, wherein one fragment comprises the sense region and the second fragment comprises the antisense region of the siNA molecule, and wherein about 19 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule and wherein at least two 3' terminal nucleotides

of each fragment of the siNA molecule are not base-paired to the nucleotides of the other fragment of the siNA molecule. In one embodiment, each of the two 3' terminal nucleotides of each fragment of the siNA molecule is a 2'-deoxy-pyrimidine nucleotide, such as 2'-deoxy-thymidine. In another embodiment, all 21 nucleotides of each fragment
5 of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule. In another embodiment, about 19 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by the gene. In another embodiment, 21 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA
10 encoded by the gene. In any of the above embodiments, the 5'-end of the fragment comprising said antisense region can optionally include a phosphate group.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits the expression of a RNA sequence (e.g., wherein said target RNA sequence is encoded by a gene or a gene involved in a pathway
15 of gene expression), wherein the siNA molecule does not contain any ribonucleotides and wherein each strand of the double-stranded siNA molecule is about 21 nucleotides long.

In one embodiment, the invention features a medicament comprising a siNA molecule of the invention.

In one embodiment, the invention features an active ingredient comprising a siNA
20 molecule of the invention.

In one embodiment, the invention features the use of a double-stranded short interfering nucleic acid (siNA) molecule to down-regulate expression of a target gene, wherein the siNA molecule comprises one or more chemical modifications and each strand of the double-stranded siNA is about 21 nucleotides long.

25 The invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of a RNA encoded by the gene or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is
30 complementary to a nucleotide sequence of the antisense strand and wherein a majority of

the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification. In one embodiment, the nucleotide sequence of the antisense strand of the double-stranded siNA molecule is complementary to the nucleotide sequence of a RNA which encodes a protein or a portion thereof. In one embodiment, each strand of

5 the siNA molecule comprises about 19 to about 29 (e.g., about 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, or 29) nucleotides, and each strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand. In one embodiment, the siNA molecule is assembled from two oligonucleotide fragments, wherein one fragment comprises the nucleotide sequence of the antisense strand of the siNA molecule and a

10 second fragment comprises nucleotide sequence of the sense region of the siNA molecule. In another embodiment, the sense strand is connected to the antisense strand via a linker molecule, such as a polynucleotide linker or a non-nucleotide linker. In one embodiment, the pyrimidine nucleotides present in the sense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and the purine nucleotides present in the sense region are 2'-deoxy

15 purine nucleotides. In another embodiment, the pyrimidine nucleotides present in the sense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and the purine nucleotides present in the sense region are 2'-O-methyl purine nucleotides. In one embodiment, the sense strand comprises a 3'-end and a 5'-end, wherein a terminal cap moiety (e.g., an inverted deoxy abasic moiety) is present at the 5'-end, the 3'-end, or both of the 5' and 3'

20 ends of the sense strand. In one embodiment, the antisense strand comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides and one or more 2'-O-methyl purine nucleotides. In one embodiment, the pyrimidine nucleotides present in the antisense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and any purine nucleotides present in the antisense strand are 2'-O-methyl purine nucleotides. In one embodiment, the

25 antisense strand comprises a phosphorothioate internucleotide linkage at the 3' end of the antisense strand. In another embodiment, the antisense strand comprises a glyceryl modification at the 3' end. In another embodiment, the 5'-end of the antisense strand optionally includes a phosphate group. In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates

30 expression of a gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of RNA encoded by a gene or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide

sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the nucleotide sequence of the antisense strand is complementary to a nucleotide sequence of the 5'-untranslated region or a portion thereof of the RNA. In another
5 embodiment, the nucleotide sequence of the antisense strand is complementary to a nucleotide sequence of the RNA or a portion thereof.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises
10 nucleotide sequence that is complementary to nucleotide sequence of a RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein each of the two strands of the siNA molecule comprises
15 21 nucleotides. In one embodiment, about 19 nucleotides of each strand of the siNA molecule are base-paired to the complementary nucleotides of the other strand of the siNA molecule and at least two 3' terminal nucleotides of each strand of the siNA molecule are not base-paired to the nucleotides of the other strand of the siNA molecule. In one embodiment, each of the two 3' terminal nucleotides of each fragment of the siNA
20 molecule are 2'-deoxy-pyrimidines, such as 2'-deoxy-thymidine. In another embodiment, each strand of the siNA molecule is base-paired to the complementary nucleotides of the other strand of the siNA molecule. In one embodiment, about 19 nucleotides of the antisense strand are base-paired to the nucleotide sequence of the RNA or a portion thereof. In another embodiment, 21 nucleotides of the antisense strand are base-paired to
25 the nucleotide sequence of the RNA or a portion thereof.

In one embodiment, the invention features a composition comprising a siNA molecule of the invention and a pharmaceutically acceptable carrier or diluent.

In one embodiment, the invention features a method of increasing the stability of a siNA molecule against cleavage by ribonucleases comprising introducing at least one
30 modified nucleotide into the siNA molecule, wherein the modified nucleotide is a 2'-deoxy-2'-fluoro nucleotide. In another embodiment, all pyrimidine nucleotides present in

the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides. In another embodiment, the modified nucleotides in the siNA include at least one 2'-deoxy-2'-fluoro cytidine or 2'-deoxy-2'-fluoro uridine nucleotide. In another embodiment, the modified nucleotides in the siNA include at least one 2'-fluoro cytidine and at least one 2'-deoxy-2'-fluoro uridine nucleotides. In another embodiment, all uridine nucleotides present in the siNA are 2'-deoxy-2'-fluoro uridine nucleotides. In another embodiment, all cytidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro cytidine nucleotides. In another embodiment, all adenosine nucleotides present in the siNA are 2'-deoxy-2'-fluoro adenosine nucleotides. In another embodiment, all guanosine nucleotides present in the siNA are 2'-deoxy-2'-fluoro guanosine nucleotides. The siNA can further comprise at least one modified internucleotidic linkage, such as phosphorothioate linkage. In another embodiment, the 2'-deoxy-2'-fluoronucleotides are present at specifically selected locations in the siNA that are sensitive to cleavage by ribonucleases, such as locations having pyrimidine nucleotides.

15 In one embodiment, the invention features the use of a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of a RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification.

In one embodiment, the invention features a short interfering nucleic acid (siNA) molecule comprising a double-stranded structure that down-regulates expression of a target nucleic acid, wherein the siNA molecule does not require a 2'-hydroxyl group containing ribonucleotide, each strand of the double-stranded structure of the siNA molecule comprises about 21 nucleotides and the siNA molecule comprises nucleotide sequence having complementarity to nucleotide sequence of the target nucleic acid or a portion thereof. The target nucleic acid can be an endogenous gene, an exogenous gene, a viral nucleic acid, or a RNA, such as a mammalian gene, plant gene, viral gene, fungal gene, bacterial gene, plant viral gene, or mammalian viral gene. Examples of mammalian viral gene include hepatitis C virus, human immunodeficiency virus, hepatitis B virus,

herpes simplex virus, cytomegalovirus, human papilloma virus, respiratory syncytial virus, influenza virus, and severe acute respiratory syndrome virus (SARS).

In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region wherein the antisense region comprises the nucleotide sequence that is
5 complementary to a nucleotide sequence or a portion thereof of the target nucleic acid and the sense region comprises a nucleotide sequence that is complementary to nucleotide sequence of the antisense region or a portion thereof.

In one embodiment, a siNA molecule of the invention is assembled from two separate oligonucleotide fragments wherein one fragment comprises the sense region and
10 the second fragment comprises the antisense region of the siNA molecule. The sense region can be connected to the antisense region via a linker molecule, such as a polynucleotide linker or non-nucleotide linker. In another embodiment, each sense region and antisense region comprise about 21 nucleotides in length. In another embodiment, about 19 nucleotides of each fragment of the siNA molecule are base-paired to the
15 complementary nucleotides of the other fragment of the siNA molecule and at least two 3' terminal nucleotides of each fragment of the siNA molecule are not base-paired to the nucleotides of the other fragment of the siNA molecule. In another embodiment, each of the two 3' terminal nucleotides of each fragment of the siNA molecule are 2'-deoxy-pyrimidines, such as the thymidine. In another embodiment, all 21 nucleotides of each
20 fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule. In another embodiment, about 19 nucleotides of the antisense region of the siNA molecule are base-paired to the nucleotide sequence or a portion thereof of the target nucleic acid. In another embodiment, 21 nucleotides of the antisense region of the siNA molecule are base-paired to the nucleotide sequence or a
25 portion thereof of the target nucleic acid. In another embodiment, the 5'-end of the fragment comprising the antisense region optionally includes a phosphate group.

In one embodiment, a siNA molecule of the invention comprises nucleotide sequence having complementarity to nucleotide sequence of RNA or a portion thereof encoded by the target nucleic acid or a portion thereof.

30 In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region, wherein the pyrimidine nucleotides when present in the sense region

are 2'-O-methyl pyrimidine nucleotides and wherein the purine nucleotides when present in the sense region are 2'-deoxy purine nucleotides.

In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region, wherein the pyrimidine nucleotides when present in the sense region
5 are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein the purine nucleotides when present in the sense region are 2'-deoxy purine nucleotides.

In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region, wherein the sense region includes a terminal cap moiety at the 5'-end, the 3'-end, or both of the 5' and 3' ends. The cap moiety can be an inverted deoxy abasic
10 moiety, an inverted deoxy thymidine moiety, or a thymidine moiety.

In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region, wherein the pyrimidine nucleotides when present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides and the purine nucleotides when present in the antisense region are 2'-O-methyl purine nucleotides.

15 In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region, wherein the pyrimidine nucleotides when present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein the purine nucleotides when present in the antisense region comprise 2'-deoxy- purine nucleotides.

In one embodiment, a siNA molecule of the invention comprises a sense region and
20 an antisense region, wherein the antisense region comprises a phosphate backbone modification at the 3' end of the antisense region. The phosphate backbone modification can be a phosphorothioate.

In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region, wherein the antisense region comprises a glyceryl modification at the
25 3' end of the antisense region.

In one embodiment, a siNA molecule of the invention comprises a sense region and an antisense region, wherein each of sense and the antisense regions of the siNA molecule comprise about 21 nucleotides.


In a non-limiting example, the introduction of chemically-modified nucleotides into nucleic acid molecules provides a powerful tool in overcoming potential limitations of *in vivo* stability and bioavailability inherent to native RNA molecules that are delivered exogenously. For example, the use of chemically-modified nucleic acid molecules can enable a lower dose of a particular nucleic acid molecule for a given therapeutic effect since chemically-modified nucleic acid molecules tend to have a longer half-life in serum. Furthermore, certain chemical modifications can improve the bioavailability of nucleic acid molecules by targeting particular cells or tissues and/or improving cellular uptake of the nucleic acid molecule. Therefore, even if the activity of a chemically-modified nucleic acid molecule is reduced as compared to a native nucleic acid molecule, for example, when compared to an all-RNA nucleic acid molecule, the overall activity of the modified nucleic acid molecule can be greater than that of the native molecule due to improved stability and/or delivery of the molecule. Unlike native unmodified siNA, chemically-modified siNA can also minimize the possibility of activating interferon activity in humans.

In any of the embodiments of siNA molecules described herein, the antisense region of a siNA molecule of the invention can comprise a phosphorothioate internucleotide linkage at the 3'-end of said antisense region. In any of the embodiments of siNA molecules described herein, the antisense region can comprise about one to about five phosphorothioate internucleotide linkages at the 5'-end of said antisense region. In any of the embodiments of siNA molecules described herein, the 3'-terminal nucleotide overhangs of a siNA molecule of the invention can comprise ribonucleotides or deoxyribonucleotides that are chemically-modified at a nucleic acid sugar, base, or backbone. In any of the embodiments of siNA molecules described herein, the 3'-terminal nucleotide overhangs can comprise one or more universal base ribonucleotides. In any of the embodiments of siNA molecules described herein, the 3'-terminal nucleotide overhangs can comprise one or more acyclic nucleotides.

One embodiment of the invention provides an expression vector comprising a nucleic acid sequence encoding at least one siNA molecule of the invention in a manner that allows expression of the nucleic acid molecule. Another embodiment of the invention provides a mammalian cell comprising such an expression vector. The mammalian cell can be a human cell. The siNA molecule of the expression vector can

comprise a sense region and an antisense region. The antisense region can comprise sequence complementary to an RNA or DNA sequence encoding a protein or polypeptide and the sense region can comprise sequence complementary to the antisense region. The siNA molecule can comprise two distinct strands having complementary sense and antisense regions. The siNA molecule can comprise a single strand having complementary sense and antisense regions.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides comprising a backbone modified internucleotide linkage having Formula I:




wherein each R1 and R2 is independently any nucleotide, non-nucleotide, or polynucleotide which can be naturally-occurring or chemically-modified, each X and Y is independently O, S, N, alkyl, or substituted alkyl, each Z and W is independently O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, or aralkyl, and wherein W, X, Y, and Z are optionally not all O. In another embodiment, a backbone modification of the invention comprises a phosphonoacetate and/or thiophosphonoacetate internucleotide linkage (see for example Sheehan et al., 2003, *Nucleic Acids Research*, 31, 4109-4118).

The chemically-modified internucleotide linkages having Formula I, for example, wherein any Z, W, X, and/or Y independently comprises a sulphur atom, can be present in one or both oligonucleotide strands of the siNA duplex, for example, in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) chemically-modified internucleotide linkages having Formula I at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified internucleotide linkages having

Formula I at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) pyrimidine nucleotides with chemically-modified internucleotide linkages having Formula I in the sense strand, the antisense strand, or both strands. In yet another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) purine nucleotides with chemically-modified internucleotide linkages having Formula I in the sense strand, the antisense strand, or both strands. In another embodiment, a siNA molecule of the invention having internucleotide linkage(s) of Formula I also comprises a chemically-modified nucleotide or non-nucleotide having any of Formulae I-VII.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides or non-nucleotides having Formula II:




wherein each R3, R4, R5, R6, R7, R8, R10, R11 and R12 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I or II; R9 is O, S, CH2, S=O, CHF, or CF2, and B is a nucleosidic base such as adenine, guanine, uracil, cytosine, thymine, 2-aminoadenosine, 5-methylcytosine, 2,6-diaminopurine, or any other non-naturally

occurring base that can be complementary or non-complementary to target RNA or a non-nucleosidic base such as phenyl, naphthyl, 3-nitropyrrole, 5-nitroindole, nebularine, pyridone, pyridinone, or any other non-naturally occurring universal base that can be complementary or non-complementary to target RNA.

5 The chemically-modified nucleotide or non-nucleotide of Formula II can be present in one or both oligonucleotide strands of the siNA duplex, for example in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more chemically-modified nucleotide or non-nucleotide of Formula II at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified nucleotides or non-nucleotides of Formula II at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified nucleotides or non-nucleotides of Formula II at the 3'-end of the sense strand, the antisense strand, or both strands.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides or non-nucleotides having Formula III:




wherein each R3, R4, R5, R6, R7, R8, R10, R11 and R12 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH,

O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO₂, NO₂, N₃, NH₂, aminoalkyl, aminoacid, aminoacyl, ONH₂, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I or II; R₉ is O, S, CH₂, S=O, CHF, or CF₂,
 5 and B is a nucleosidic base such as adenine, guanine, uracil, cytosine, thymine, 2-aminoadenosine, 5-methylcytosine, 2,6-diaminopurine, or any other non-naturally occurring base that can be employed to be complementary or non-complementary to target RNA or a non-nucleosidic base such as phenyl, naphthyl, 3-nitropyrrole, 5-nitroindole, nebularine, pyridone, pyridinone, or any other non-naturally occurring
 10 universal base that can be complementary or non-complementary to target RNA.

The chemically-modified nucleotide or non-nucleotide of Formula III can be present in one or both oligonucleotide strands of the siNA duplex, for example, in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more chemically-modified nucleotide or non-nucleotide of Formula
 15 III at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified nucleotide(s) or non-nucleotide(s) of Formula III at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an
 20 exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified nucleotide or non-nucleotide of Formula III at the 3'-end of the sense strand, the antisense strand, or both strands.

In another embodiment, a siNA molecule of the invention comprises a nucleotide having Formula II or III, wherein the nucleotide having Formula II or III is in an inverted
 25 configuration. For example, the nucleotide having Formula II or III is connected to the siNA construct in a 3'-3', 3'-2', 2'-3', or 5'-5' configuration, such as at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of one or both siNA strands.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) inside a cell
 30 or reconstituted *in vitro* system, wherein the chemical modification comprises a 5'-terminal phosphate group having Formula IV:



wherein each X and Y is independently O, S, N, alkyl, substituted alkyl, or alkylhalo; wherein each Z and W is independently O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, aralkyl, or alkylhalo or acetyl; and/or wherein W, X, Y and Z are not all O.

5 In one embodiment, the invention features a siNA molecule having a 5'-terminal phosphate group having Formula IV on the target-complementary strand, for example, a strand complementary to a target RNA, wherein the siNA molecule comprises an all RNA siNA molecule. In another embodiment, the invention features a siNA molecule having a 5'-terminal phosphate group having Formula IV on the target-complementary strand
 10 wherein the siNA molecule also comprises about 1 to about 3 (*e.g.*, about 1, 2, or 3) nucleotide 3'-terminal nucleotide overhangs having about 1 to about 4 (*e.g.*, about 1, 2, 3, or 4) deoxyribonucleotides on the 3'-end of one or both strands. In another embodiment, a 5'-terminal phosphate group having Formula IV is present on the target-complementary strand of a siNA molecule of the invention, for example a siNA molecule having
 15 chemical modifications having any of Formulae I-VII.

 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more phosphorothioate, phosphonoacetate, and/or thiophosphonoacetate internucleotide
 20 linkages. For example, in a non-limiting example, the invention features a chemically-modified short interfering nucleic acid (siNA) having about 1, 2, 3, 4, 5, 6, 7, 8 or more phosphorothioate internucleotide linkages in one siNA strand. In yet another embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) individually having about 1, 2, 3, 4, 5, 6, 7, 8 or more phosphorothioate
 25 internucleotide linkages in both siNA strands. The phosphorothioate internucleotide linkages can be present in one or both oligonucleotide strands of the siNA duplex, for example in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more phosphorothioate, phosphonoacetate, and/or

thiophosphonoacetate internucleotide linkages at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) consecutive phosphorothioate internucleotide linkages at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) pyrimidine phosphorothioate internucleotide linkages in the sense strand, the antisense strand, or both strands. In yet another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) purine phosphorothioate internucleotide linkages in the sense strand, the antisense strand, or both strands.

In one embodiment, the invention features a siNA molecule, wherein the sense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or about one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

In another embodiment, the invention features a siNA molecule, wherein the sense strand comprises about 1 to about 5, specifically about 1, 2, 3, 4, or 5 phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, or more) 2'-deoxy,

2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 5 or more, specifically about 1, 2, 3, 4, 5, or more
 5 phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4,
 10 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 5 or more, for example about 1, 2, 3, 4, 5, or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different
 15 strand.

In one embodiment, the invention features a siNA molecule, wherein the antisense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or about one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more
 20 (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more)
 25 more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-
 30 modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more

phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3' and 5'-ends, being present in the same or different strand.

In another embodiment, the invention features a siNA molecule, wherein the antisense strand comprises about 1 to about 5 or more, specifically about 1, 2, 3, 4, 5 or
 5 more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 5 or
 10 more, specifically about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment,
 15 one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 5, for example about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being
 20 present in the same or different strand.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule having about 1 to about 5, specifically about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages in each strand of the siNA molecule.

In another embodiment, the invention features a siNA molecule comprising 2'-5'
 25 internucleotide linkages. The 2'-5' internucleotide linkage(s) can be at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of one or both siNA sequence strands. In addition, the 2'-5' internucleotide linkage(s) can be present at various other positions within one or both siNA sequence strands, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a pyrimidine nucleotide in one or both strands of the
 30 siNA molecule can comprise a 2'-5' internucleotide linkage, or about 1, 2, 3, 4, 5, 6, 7, 8,

9, 10, or more including every internucleotide linkage of a purine nucleotide in one or both strands of the siNA molecule can comprise a 2'-5' internucleotide linkage.

In another embodiment, a chemically-modified siNA molecule of the invention comprises a duplex having two strands, one or both of which can be chemically-modified, wherein each strand is about 18 to about 27 (*e.g.*, about 18, 19, 20, 21, 22, 23, 24, 25, 26, or 27) nucleotides in length, wherein the duplex has about 18 to about 23 (*e.g.*, about 18, 19, 20, 21, 22, or 23) base pairs, and wherein the chemical modification comprises a structure having any of Formulae I-VII. For example, an exemplary chemically-modified siNA molecule of the invention comprises a duplex having two strands, one or both of which can be chemically-modified with a chemical modification having any of Formulae I-VII or any combination thereof, wherein each strand consists of about 21 nucleotides, each having a 2-nucleotide 3'-terminal nucleotide overhang, and wherein the duplex has about 19 base pairs. In another embodiment, a siNA molecule of the invention comprises a single stranded hairpin structure, wherein the siNA is about 36 to about 70 (*e.g.*, about 36, 40, 45, 50, 55, 60, 65, or 70) nucleotides in length having about 18 to about 23 (*e.g.*, about 18, 19, 20, 21, 22, or 23) base pairs, and wherein the siNA can include a chemical modification comprising a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises a linear oligonucleotide having about 42 to about 50 (*e.g.*, about 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides that is chemically-modified with a chemical modification having any of Formulae I-VII or any combination thereof, wherein the linear oligonucleotide forms a hairpin structure having about 19 base pairs and a 2-nucleotide 3'-terminal nucleotide overhang. In another embodiment, a linear hairpin siNA molecule of the invention contains a stem loop motif, wherein the loop portion of the siNA molecule is biodegradable. For example, a linear hairpin siNA molecule of the invention is designed such that degradation of the loop portion of the siNA molecule *in vivo* can generate a double-stranded siNA molecule with 3'-terminal overhangs, such as 3'-terminal nucleotide overhangs comprising about 2 nucleotides.

In another embodiment, a siNA molecule of the invention comprises a hairpin structure, wherein the siNA is about 25 to about 50 (*e.g.*, about 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides in length having about 3 to about 25 (*e.g.*, about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16,

17, 18, 19, 20, 21, 22, 23, 24, or 25) base pairs, and wherein the siNA can include one or more chemical modifications comprising a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises a linear oligonucleotide having about 25 to about 35 (*e.g.*, about 5 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35) nucleotides that is chemically-modified with one or more chemical modifications having any of Formulae I-VII or any combination thereof, wherein the linear oligonucleotide forms a hairpin structure having about 3 to about 23 (*e.g.*, about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23) base pairs and a 5'-terminal phosphate group that can be chemically modified as 10 described herein (for example a 5'-terminal phosphate group having Formula IV). In another embodiment, a linear hairpin siNA molecule of the invention contains a stem loop motif, wherein the loop portion of the siNA molecule is biodegradable. In another embodiment, a linear hairpin siNA molecule of the invention comprises a loop portion comprising a non-nucleotide linker.

15 In another embodiment, a siNA molecule of the invention comprises an asymmetric hairpin structure, wherein the siNA is about 25 to about 50 (*e.g.*, about 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides in length having about 3 to about 20 (*e.g.*, about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20) base pairs, and wherein the siNA can include one or 20 more chemical modifications comprising a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises a linear oligonucleotide having about 25 to about 35 (*e.g.*, about 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35) nucleotides that is chemically-modified with one or more chemical modifications having any of Formulae I-VII or any combination 25 thereof, wherein the linear oligonucleotide forms an asymmetric hairpin structure having about 3 to about 18 (*e.g.*, about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or 18) base pairs and a 5'-terminal phosphate group that can be chemically modified as described herein (for example a 5'-terminal phosphate group having Formula IV). In another embodiment, an asymmetric hairpin siNA molecule of the invention contains a stem loop 30 motif, wherein the loop portion of the siNA molecule is biodegradable. In another embodiment, an asymmetric hairpin siNA molecule of the invention comprises a loop portion comprising a non-nucleotide linker.


In another embodiment, a siNA molecule of the invention comprises an asymmetric double stranded structure having separate polynucleotide strands comprising sense and antisense regions, wherein the antisense region is about 16 to about 25 (*e.g.*, about 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25) nucleotides in length, wherein the sense region is about 3 to about 18 (*e.g.*, about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 18) nucleotides in length, wherein the sense region the antisense region have at least 3 complementary nucleotides, and wherein the siNA can include one or more chemical modifications comprising a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises an asymmetric double stranded structure having separate polynucleotide strands comprising sense and antisense regions, wherein the antisense region is about 18 to about 22 (*e.g.*, about 18, 19, 20, 21, or 22) nucleotides in length and wherein the sense region is about 3 to about 15 (*e.g.*, about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15) nucleotides in length, wherein the sense region the antisense region have at least 3 complementary nucleotides, and wherein the siNA can include one or more chemical modifications comprising a structure having any of Formulae I-VII or any combination thereof. In another embodiment, the asymmetric double stranded siNA molecule can also have a 5'-terminal phosphate group that can be chemically modified as described herein (for example a 5'-terminal phosphate group having Formula IV).

In another embodiment, a siNA molecule of the invention comprises a circular nucleic acid molecule, wherein the siNA is about 38 to about 70 (*e.g.*, about 38, 40, 45, 50, 55, 60, 65, or 70) nucleotides in length having about 18 to about 23 (*e.g.*, about 18, 19, 20, 21, 22, or 23) base pairs, and wherein the siNA can include a chemical modification, which comprises a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises a circular oligonucleotide having about 42 to about 50 (*e.g.*, about 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides that is chemically-modified with a chemical modification having any of Formulae I-VII or any combination thereof, wherein the circular oligonucleotide forms a dumbbell shaped structure having about 19 base pairs and 2 loops.

In another embodiment, a circular siNA molecule of the invention contains two loop motifs, wherein one or both loop portions of the siNA molecule is biodegradable.


For example, a circular siNA molecule of the invention is designed such that degradation of the loop portions of the siNA molecule *in vivo* can generate a double-stranded siNA molecule with 3'-terminal overhangs, such as 3'-terminal nucleotide overhangs comprising about 2 nucleotides.

- 5 In one embodiment, a siNA molecule of the invention comprises at least one (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) abasic moiety, for example a compound having Formula V:



- wherein each R3, R4, R5, R6, R7, R8, R10, R11, R12, and R13 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I or II; R9 is O, S, CH2, S=O, CHF, or CF2.
- 10
- 15

In one embodiment, a siNA molecule of the invention comprises at least one (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) inverted abasic moiety, for example a compound having Formula VI:



In another embodiment, a moiety having any of Formula V, VI or VII of the invention is at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of a siNA molecule of the invention. For example, a moiety having Formula V, VI or VII can be present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense strand, the sense strand, or both antisense and sense strands of the siNA molecule. In addition, a moiety having Formula VII can be present at the 3'-end or the 5'-end of a hairpin siNA molecule as described herein.

In another embodiment, a siNA molecule of the invention comprises an abasic residue having Formula V or VI, wherein the abasic residue having Formula V or VI is connected to the siNA construct in a 3-3', 3-2', 2-3', or 5-5' configuration, such as at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of one or both siNA strands.

In one embodiment, a siNA molecule of the invention comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) locked nucleic acid (LNA) nucleotides, for example at the 5'-end, the 3'-end, both of the 5' and 3'-ends, or any combination thereof, of the siNA molecule.

In another embodiment, a siNA molecule of the invention comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) acyclic nucleotides, for example at the 5'-end, the 3'-end, both of the 5' and 3'-ends, or any combination thereof, of the siNA molecule.

In one embodiment, the sense strand of a double stranded siNA molecule of the invention comprises a terminal cap moiety, (see for example **Figure 22**) such as an inverted deoxyabasic moiety or inverted nucleotide, at the 3'-end, 5'-end, or both 3' and 5'-ends of the sense strand.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises a sense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where any (*e.g.*, one or more or all) purine nucleotides present in the sense region are 2'-

deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides).

5 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises a sense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and
10 where any (*e.g.*, one or more or all) purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), wherein any nucleotides comprising a 3'-terminal nucleotide overhang that are present in said sense region are 2'-deoxy nucleotides.

15 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises a sense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and
20 where any (*e.g.*, one or more or all) purine nucleotides present in the sense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides).

25 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises a sense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and
30 where any (*e.g.*, one or more or all) purine nucleotides present in the sense region are 2'-

O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), wherein any nucleotides comprising a 3'-terminal nucleotide overhang that are present in said sense region are 2'-deoxy nucleotides.

5 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or
10 alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any (*e.g.*, one or more or all) purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides).

15 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or
20 alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any (*e.g.*, one or more or all) purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), wherein any nucleotides comprising a 3'-terminal
25 nucleotide overhang that are present in said antisense region are 2'-deoxy nucleotides.

 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*,
30 wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine

nucleotides), and where any (*e.g.*, one or more or all) purine nucleotides present in the antisense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides).

5 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system comprising a sense region and an antisense region. In one embodiment, the sense region comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-
10 fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides). The sense region can comprise inverted deoxy abasic modifications that are optionally present at the 3'-end, the
15 5'-end, or both of the 3' and 5'-ends of the sense region. The sense region can optionally further comprise a 3'-terminal overhang having about 1 to about 4 (*e.g.*, about 1, 2, 3, or 4) 2'-deoxyribonucleotides. The antisense region comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-
20 deoxy-2'-fluoro pyrimidine nucleotides), and one or more 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides). The antisense region can comprise a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3'
25 and 5'-ends of the antisense sequence. The antisense region optionally further comprises a 3'-terminal nucleotide overhang having about 1 to about 4 (*e.g.*, about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (*e.g.*, 1, 2, 3, or 4) phosphorothioate internucleotide linkages. Non-limiting examples of these chemically-modified siNAs are shown in **Figures 18 and 19** and **Table IV** herein.

30 In another embodiment of the chemically-modified short interfering nucleic acid comprising a sense region and an antisense region, the sense region comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides

are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more purine ribonucleotides (e.g., wherein all purine nucleotides are purine ribonucleotides or alternately a plurality of purine nucleotides are purine ribonucleotides). The sense region can also comprise inverted deoxy abasic modifications that are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region. The sense region optionally further comprises a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides. The antisense region comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides). The antisense region can also comprise a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence. The antisense region optionally further comprises a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages. Non-limiting examples of these chemically-modified siNAs are shown in **Figures 18 and 19** and **Table IV** herein.

In another embodiment of the chemically-modified short interfering nucleic acid comprising a sense region and an antisense region, the sense region comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more purine nucleotides selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides or alternately a plurality of purine nucleotides are selected from the group consisting of 2'-deoxy

nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides). The sense region can comprise inverted deoxy abasic modifications that are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region. The sense region can optionally further comprise a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides. The antisense region comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more purine nucleotides selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides or alternately a plurality of purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides). The antisense can also comprise a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence. The antisense region optionally further comprises a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages.

In another embodiment, any modified nucleotides present in the siNA molecules of the invention, preferably in the antisense strand of the siNA molecules of the invention, but also optionally in the sense and/or both antisense and sense strands, comprise modified nucleotides having properties or characteristics similar to naturally occurring ribonucleotides. For example, the invention features siNA molecules including modified nucleotides having a Northern conformation (e.g., Northern pseudorotation cycle, see for example Saenger, *Principles of Nucleic Acid Structure*, Springer-Verlag ed., 1984). As such, chemically modified nucleotides present in the siNA molecules of the invention, preferably in the antisense strand of the siNA molecules of the invention, but also

optionally in the sense and/or both antisense and sense strands, are resistant to nuclease degradation while at the same time maintaining the capacity to mediate RNAi. Non-limiting examples of nucleotides having a northern configuration include locked nucleic acid (LNA) nucleotides (e.g., 2'-O,4'-C-methylene-(D-ribofuranosyl) nucleotides); 2'-methoxyethoxy (MOE) nucleotides; 2'-methyl-thio-ethyl, 2'-deoxy-2'-fluoro nucleotides, 2'-deoxy-2'-chloro nucleotides, 2'-azido nucleotides, and 2'-O-methyl nucleotides.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid molecule (siNA) capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises a conjugate attached to the chemically-modified siNA molecule. The conjugate can be attached to the chemically-modified siNA molecule via a covalent attachment. In one embodiment, the conjugate is attached to the chemically-modified siNA molecule via a biodegradable linker. In one embodiment, the conjugate molecule is attached at the 3'-end of either the sense strand, the antisense strand, or both strands of the chemically-modified siNA molecule. In another embodiment, the conjugate molecule is attached at the 5'-end of either the sense strand, the antisense strand, or both strands of the chemically-modified siNA molecule. In yet another embodiment, the conjugate molecule is attached both the 3'-end and 5'-end of either the sense strand, the antisense strand, or both strands of the chemically-modified siNA molecule, or any combination thereof. In one embodiment, the conjugate molecule of the invention comprises a molecule that facilitates delivery of a chemically-modified siNA molecule into a biological system, such as a cell. In another embodiment, the conjugate molecule attached to the chemically-modified siNA molecule is a poly ethylene glycol, human serum albumin, or a ligand for a cellular receptor that can mediate cellular uptake. Examples of specific conjugate molecules contemplated by the instant invention that can be attached to chemically-modified siNA molecules are described in Vargeese *et al.*, U.S. Serial No. 10/201,394, incorporated by reference herein. The type of conjugates used and the extent of conjugation of siNA molecules of the invention can be evaluated for improved pharmacokinetic profiles, bioavailability, and/or stability of siNA constructs while at the same time maintaining the ability of the siNA to mediate RNAi activity. As such, one skilled in the art can screen siNA constructs that are modified with various conjugates to determine whether the siNA

conjugate complex possesses improved properties while maintaining the ability to mediate RNAi, for example in animal models as are generally known in the art.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system, wherein the chemically-modified siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where one or more purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and inverted deoxy abasic modifications that are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (*e.g.*, about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the chemically-modified short interfering nucleic acid molecule comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein one or more purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (*e.g.*, about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (*e.g.*, 1, 2, 3, or 4) phosphorothioate internucleotide linkages. Non-limiting examples of these chemically-modified siNAs are shown in **Figures 18 and 19** and **Table IV** herein.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference

(RNAi) inside a cell or reconstituted *in vitro* system, wherein the chemically-modified siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where one or more purine nucleotides present in the sense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and inverted deoxy abasic modifications that are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (*e.g.*, about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the chemically-modified short interfering nucleic acid molecule comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein one or more purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (*e.g.*, about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (*e.g.*, 1, 2, 3, or 4) phosphorothioate internucleotide linkages. Non-limiting examples of these chemically-modified siNAs are shown in **Figures 18 and 19** and **Table IV** herein.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system, wherein the siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides

are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where one or more purine nucleotides present in the sense region are purine ribonucleotides (e.g., wherein all purine nucleotides are purine ribonucleotides or alternately a plurality of purine nucleotides are purine ribonucleotides), and inverted deoxy abasic modifications that are optionally present at

5 the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the siNA comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-

10 deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as

15 any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide

20 linkages. Non-limiting examples of these chemically-modified siNAs are shown in **Figures 18 and 19** and **Table IV** herein.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) inside a cell or reconstituted *in vitro* system, wherein the chemically-modified

25 siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and for example where one or more purine nucleotides present in the sense region are selected from the

30 group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides,

locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides or alternately a plurality of purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides), and wherein inverted deoxy abasic modifications are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the chemically-modified short interfering nucleic acid molecule comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein one or more purine nucleotides present in the antisense region are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides or alternately a plurality of purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages.

In one embodiment, the invention features a short interfering nucleic acid (siNA) molecule of the invention, wherein the siNA further comprises a nucleotide, non-nucleotide, or mixed nucleotide/non-nucleotide linker that joins the sense region of the siNA to the antisense region of the siNA. In one embodiment, a nucleotide linker of the invention can be a linker of ≥ 2 nucleotides in length, for example 3, 4, 5, 6, 7, 8, 9, or 10

nucleotides in length. In another embodiment, the nucleotide linker can be a nucleic acid aptamer. By "aptamer" or "nucleic acid aptamer" as used herein is meant a nucleic acid molecule that binds specifically to a target molecule wherein the nucleic acid molecule has sequence that comprises a sequence recognized by the target molecule in its natural setting. Alternately, an aptamer can be a nucleic acid molecule that binds to a target molecule where the target molecule does not naturally bind to a nucleic acid. The target molecule can be any molecule of interest. For example, the aptamer can be used to bind to a ligand-binding domain of a protein, thereby preventing interaction of the naturally occurring ligand with the protein. This is a non-limiting example and those in the art will recognize that other embodiments can be readily generated using techniques generally known in the art (see, for example, Gold *et al.*, 1995, *Annu. Rev. Biochem.*, 64, 763; Brody and Gold, 2000, *J. Biotechnol.*, 74, 5; Sun, 2000, *Curr. Opin. Mol. Ther.*, 2, 100; Kusser, 2000, *J. Biotechnol.*, 74, 27; Hermann and Patel, 2000, *Science*, 287, 820; and Jayasena, 1999, *Clinical Chemistry*, 45, 1628.)

In yet another embodiment, a non-nucleotide linker of the invention comprises abasic nucleotide, polyether, polyamine, polyamide, peptide, carbohydrate, lipid, polyhydrocarbon, or other polymeric compounds (e.g. polyethylene glycols such as those having between 2 and 100 ethylene glycol units). Specific examples include those described by Seela and Kaiser, *Nucleic Acids Res.* 1990, 18:6353 and *Nucleic Acids Res.* 1987, 15:3113; Cload and Schepartz, *J. Am. Chem. Soc.* 1991, 113:6324; Richardson and Schepartz, *J. Am. Chem. Soc.* 1991, 113:5109; Ma *et al.*, *Nucleic Acids Res.* 1993, 21:2585 and *Biochemistry* 1993, 32:1751; Durand *et al.*, *Nucleic Acids Res.* 1990, 18:6353; McCurdy *et al.*, *Nucleosides & Nucleotides* 1991, 10:287; Jschke *et al.*, *Tetrahedron Lett.* 1993, 34:301; Ono *et al.*, *Biochemistry* 1991, 30:9914; Arnold *et al.*, International Publication No. WO 89/02439; Usman *et al.*, International Publication No. WO 95/06731; Dudycz *et al.*, International Publication No. WO 95/11910 and Ferentz and Verdine, *J. Am. Chem. Soc.* 1991, 113:4000, all hereby incorporated by reference herein. A "non-nucleotide" further means any group or compound that can be incorporated into a nucleic acid chain in the place of one or more nucleotide units, including either sugar and/or phosphate substitutions, and allows the remaining bases to exhibit their enzymatic activity. The group or compound can be abasic in that it does not

contain a commonly recognized nucleotide base, such as adenosine, guanine, cytosine, uracil or thymine, for example at the C1 position of the sugar.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid molecule (siNA) capable of mediating RNA interference (RNAi) against a target gene inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises a conjugate covalently attached to the chemically-modified siNA molecule. Non-limiting examples of conjugates contemplated by the invention include conjugates and ligands described in Vargeese *et al.*, USSN 10/427,160, filed April 30, 2003, incorporated by reference herein in its entirety, including the drawings. In another embodiment, the conjugate is covalently attached to the chemically-modified siNA molecule via a biodegradable linker. In one embodiment, the conjugate molecule is attached at the 3'-end of either the sense strand, the antisense strand, or both strands of the chemically-modified siNA molecule. In another embodiment, the conjugate molecule is attached at the 5'-end of either the sense strand, the antisense strand, or both strands of the chemically-modified siNA molecule. In yet another embodiment, the conjugate molecule is attached both the 3'-end and 5'-end of either the sense strand, the antisense strand, or both strands of the chemically-modified siNA molecule, or any combination thereof. In one embodiment, a conjugate molecule of the invention comprises a molecule that facilitates delivery of a chemically-modified siNA molecule into a biological system, such as a cell. In another embodiment, the conjugate molecule attached to the chemically-modified siNA molecule is a polyethylene glycol, human serum albumin, or a ligand for a cellular receptor that can mediate cellular uptake. Examples of specific conjugate molecules contemplated by the instant invention that can be attached to chemically-modified siNA molecules are described in Vargeese *et al.*, U.S. Serial No. 10/201,394, incorporated by reference herein. The type of conjugates used and the extent of conjugation of siNA molecules of the invention can be evaluated for improved pharmacokinetic profiles, bioavailability, and/or stability of siNA constructs while at the same time maintaining the ability of the siNA to mediate RNAi activity. As such, one skilled in the art can screen siNA constructs that are modified with various conjugates to determine whether the siNA conjugate complex possesses improved properties while maintaining the ability to mediate RNAi, for example in animal models as are generally known in the art.

In one embodiment, the invention features a short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) inside a cell or reconstituted in vitro system, wherein one or both strands of the siNA molecule that are assembled from two separate oligonucleotides do not comprise any ribonucleotides. For example, a siNA molecule can be assembled from a single oligonucleotide where the sense and antisense regions of the siNA comprise separate oligonucleotides that do not have any ribonucleotides (e.g., nucleotides having a 2'-OH group) present in the oligonucleotides. In another example, a siNA molecule can be assembled from a single oligonucleotide where the sense and antisense regions of the siNA are linked or circularized by a nucleotide or non-nucleotide linker as described herein, wherein the oligonucleotide does not have any ribonucleotides (e.g., nucleotides having a 2'-OH group) present in the oligonucleotide. Applicant has surprisingly found that the presense of ribonucleotides (e.g., nucleotides having a 2'-hydroxyl group) within the siNA molecule is not required or essential to support RNAi activity. As such, in one embodiment, all positions within the siNA can include chemically modified nucleotides and/or non-nucleotides such as nucleotides and or non-nucleotides having Formula I, II, III, IV, V, VI, or VII or any combination thereof to the extent that the ability of the siNA molecule to support RNAi activity in a cell is maintained.

In one embodiment, the invention features a siNA molecule that does not require the presence of a 2'-OH group (ribonucleotide) to be present within the siNA molecule to support RNA interference.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence. In another embodiment, the single stranded siNA molecule of the invention comprises a 5'-terminal phosphate group. In another embodiment, the single stranded siNA molecule of the invention comprises a 5'-terminal phosphate group and a 3'-terminal phosphate group (e.g., a 2',3'-cyclic phosphate). In another embodiment, the single stranded siNA molecule of the invention comprises about 19 to about 29 (e.g., about 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, or 29) nucleotides. In yet another embodiment, the single stranded siNA molecule of the invention comprises one or more chemically modified nucleotides or non-nucleotides described herein. For

example, all the positions within the siNA molecule can include chemically-modified nucleotides such as nucleotides having any of Formulae I-VII, or any combination thereof to the extent that the ability of the siNA molecule to support RNAi activity in a cell is maintained.

5 In one embodiment, the single stranded siNA molecule having complementarity to a target nucleic acid sequence comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more 2'-O-methyl purine nucleotides (e.g., wherein
10 all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides). In another embodiment, the single stranded siNA molecule comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro
15 pyrimidine nucleotides), and one or more 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides). In another embodiment, the single stranded siNA molecule comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or
20 alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), wherein any purine nucleotides present in the antisense region are locked nucleic acid (LNA) nucleotides (e.g., wherein all purine nucleotides are LNA nucleotides or alternately a plurality of purine nucleotides are LNA nucleotides). In another embodiment, the single stranded siNA molecule comprises one or more 2'-deoxy-2'-
25 fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more 2'-methoxyethyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-methoxyethyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-methoxyethyl purine nucleotides),
30 the single stranded siNA can comprise a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence. The single

stranded siNA optionally further comprises about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages. The single stranded siNA optionally further comprises a
 5 terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in
 10 the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a
 15 plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence. The siNA optionally further comprises about 1 to about 4 or more (e.g., about 1, 2, 3, 4 or more) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, 4 or
 20 more) phosphorothioate, phosphonoacetate, and/or thiophosphonoacetate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group. In any of these embodiments, any purine nucleotides present in the antisense region are alternatively 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a
 25 plurality of purine nucleotides are 2'-deoxy purine nucleotides). Also, in any of these embodiments, any purine nucleotides present in the siNA (i.e., purine nucleotides present in the sense and/or antisense region) can alternatively be locked nucleic acid (LNA) nucleotides (e.g., wherein all purine nucleotides are LNA nucleotides or alternately a
 30 plurality of purine nucleotides are LNA nucleotides). Also, in any of these embodiments, any purine nucleotides present in the siNA are alternatively 2'-methoxyethyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-methoxyethyl purine nucleotides

or alternately a plurality of purine nucleotides are 2'-methoxyethyl purine nucleotides). In another embodiment, any modified nucleotides present in the single stranded siNA molecules of the invention comprise modified nucleotides having properties or characteristics similar to naturally occurring ribonucleotides. For example, the invention features siNA molecules including modified nucleotides having a Northern conformation (e.g., Northern pseudorotation cycle, see for example Saenger, *Principles of Nucleic Acid Structure*, Springer-Verlag ed., 1984). As such, chemically modified nucleotides present in the single stranded siNA molecules of the invention are preferably resistant to nuclease degradation while at the same time maintaining the capacity to mediate RNAi

10 In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine
15 nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the siNA are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification,
20 such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate
25 internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to
30 a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of

pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the siNA are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and a terminal cap modification, such as any
5 modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide
10 linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to
15 a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the siNA are locked nucleic acid (LNA) nucleotides (e.g.,
20 wherein all purine nucleotides are LNA nucleotides or alternately a plurality of purine nucleotides are LNA nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-
25 deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA
30 molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in

the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the siNA are 2'-methoxyethyl purine nucleotides (e.g.,
5 wherein all purine nucleotides are 2'-methoxyethyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-methoxyethyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 22**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or
10 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In another embodiment, any modified nucleotides present in the single stranded
15 siNA molecules of the invention comprise modified nucleotides having properties or characteristics similar to naturally occurring ribonucleotides. For example, the invention features siNA molecules including modified nucleotides having a Northern conformation (e.g., Northern pseudorotation cycle, see for example Saenger, *Principles of Nucleic Acid Structure*, Springer-Verlag ed., 1984). As such, chemically modified nucleotides present
20 in the single stranded siNA molecules of the invention are preferably resistant to nuclease degradation while at the same time maintaining the capacity to mediate RNAi.

In one embodiment, the invention features a method for modulating the expression of a gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a
25 sequence complementary to RNA of the gene; and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the gene in the cell.

In one embodiment, the invention features a method for modulating the expression of a gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a
30 sequence complementary to RNA of the gene and wherein the sense strand sequence of the siNA comprises a sequence substantially similar to the sequence of the target RNA;

and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the gene in the cell.

In another embodiment, the invention features a method for modulating the expression of more than one gene within a cell comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the genes; and (b) introducing the siNA molecules into a cell under conditions suitable to modulate the expression of the genes in the cell.

In another embodiment, the invention features a method for modulating the expression of more than one gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the gene and wherein the sense strand sequence of the siNA comprises a sequence substantially similar to the sequence of the target RNA; and (b) introducing the siNA molecules into a cell under conditions suitable to modulate the expression of the genes in the cell.

In one embodiment, siNA molecules of the invention are used as reagents in ex vivo applications. For example, siNA reagents are introduced into tissue or cells that are transplanted into a subject for therapeutic effect. The cells and/or tissue can be derived from an organism or subject that later receives the explant, or can be derived from another organism or subject prior to transplantation. The siNA molecules can be used to modulate the expression of one or more genes in the cells or tissue, such that the cells or tissue obtain a desired phenotype or are able to perform a function when transplanted in vivo. In one embodiment, certain target cells from a patient are extracted. These extracted cells are contacted with siNAs targeting a specific nucleotide sequence within the cells under conditions suitable for uptake of the siNAs by these cells (e.g. using delivery reagents such as cationic lipids, liposomes and the like or using techniques such as electroporation to facilitate the delivery of siNAs into cells). The cells are then reintroduced back into the same patient or other patients. Non-limiting examples of ex vivo applications include use in organ/tissue transplant, tissue grafting, or treatment of pulmonary disease (e.g., restenosis) or prevent neointimal hyperplasia and atherosclerosis in vein grafts. Such ex vivo applications may also be used to treat conditions associated with

coronary and peripheral bypass graft failure, for example, such methods can be used in conjunction with peripheral vascular bypass graft surgery and coronary artery bypass graft surgery. Additional applications include transplants to treat CNS lesions or injury, including use in treatment of neurodegenerative conditions such as Alzheimer's disease,
5 Parkinson's Disease, Epilepsy, Dementia, Huntington's disease, or amyotrophic lateral sclerosis (ALS).

In one embodiment, the invention features a method of modulating the expression of a gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises
10 a sequence complementary to RNA of the gene; and (b) introducing the siNA molecule into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable
15 to modulate the expression of the gene in that organism.

In one embodiment, the invention features a method of modulating the expression of a gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the gene and wherein the sense strand sequence of
20 the siNA comprises a sequence substantially similar to the sequence of the target RNA; and (b) introducing the siNA molecule into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another
25 organism under conditions suitable to modulate the expression of the gene in that organism.

In another embodiment, the invention features a method of modulating the expression of more than one gene in a tissue explant comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA
30 strands comprises a sequence complementary to RNA of the genes; and (b) introducing the siNA molecules into a cell of the tissue explant derived from a particular organism

under conditions suitable to modulate the expression of the genes in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the genes in that organism.

5 In one embodiment, the invention features a method of modulating the expression of a gene in an organism comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the gene; and (b) introducing the siNA molecule into the organism under conditions suitable to modulate the expression of the gene in the
10 organism.

 In another embodiment, the invention features a method of modulating the expression of more than one gene in an organism comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the genes; and (b) introducing
15 the siNA molecules into the organism under conditions suitable to modulate the expression of the genes in the organism.

 In one embodiment, the invention features a method for modulating the expression of a gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded
20 sequence having complementarity to RNA of the gene; and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the gene in the cell.

 In one embodiment, the invention features a method of modulating the expression of a target gene in an tissue or organ comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single
25 stranded sequence having complementarity to RNA of the target gene; and (b) introducing the siNA molecule into the tissue or organ under conditions suitable to modulate the expression of the target gene in the organism. In another embodiment, the tissue is ocular tissue and the organ is the eye. In another embodiment, the tissue
30 comprises hepatocytes and/or hepatic tissue and the organ is the liver.

In another embodiment, the invention features a method for modulating the expression of more than one gene within a cell comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the gene; and
5 (b) contacting the siNA molecule with a cell in vitro or in vivo under conditions suitable to modulate the expression of the genes in the cell.

In one embodiment, the invention features a method of modulating the expression of a gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single
10 stranded sequence having complementarity to RNA of the gene; and (b) contacting the siNA molecule with a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable
15 to modulate the expression of the gene in that organism.

In another embodiment, the invention features a method of modulating the expression of more than one gene in a tissue explant comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the gene; and
20 (b) introducing the siNA molecules into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the genes in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the genes in that
25 organism.

In one embodiment, the invention features a method of modulating the expression of a gene in an organism comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the gene; and (b) introducing the siNA
30 molecule into the organism under conditions suitable to modulate the expression of the gene in the organism.

In another embodiment, the invention features a method of modulating the expression of more than one gene in an organism comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the gene; and
5 (b) introducing the siNA molecules into the organism under conditions suitable to modulate the expression of the genes in the organism.

In one embodiment, the invention features a method of modulating the expression of a gene in an organism comprising contacting the organism with a siNA molecule of the invention under conditions suitable to modulate the expression of the gene in the
10 organism.

In another embodiment, the invention features a method of modulating the expression of more than one gene in an organism comprising contacting the organism with one or more siNA molecules of the invention under conditions suitable to modulate the expression of the genes in the organism.

15 The siNA molecules of the invention can be designed to down regulate or inhibit target gene expression through RNAi targeting of a variety of RNA molecules. In one embodiment, the siNA molecules of the invention are used to target various RNAs corresponding to a target gene. Non-limiting examples of such RNAs include messenger RNA (mRNA), alternate RNA splice variants of target gene(s), post-transcriptionally
20 modified RNA of target gene(s), pre-mRNA of target gene(s), and/or RNA templates. If alternate splicing produces a family of transcripts that are distinguished by usage of appropriate exons, the instant invention can be used to inhibit gene expression through the appropriate exons to specifically inhibit or to distinguish among the functions of gene family members. For example, a protein that contains an alternatively spliced
25 transmembrane domain can be expressed in both membrane bound and secreted forms. Use of the invention to target the exon containing the transmembrane domain can be used to determine the functional consequences of pharmaceutical targeting of membrane bound as opposed to the secreted form of the protein. Non-limiting examples of applications of the invention relating to targeting these RNA molecules include therapeutic
30 pharmaceutical applications, pharmaceutical discovery applications, molecular diagnostic and gene function applications, and gene mapping, for example using single nucleotide

polymorphism mapping with siNA molecules of the invention. Such applications can be implemented using known gene sequences or from partial sequences available from an expressed sequence tag (EST).

In another embodiment, the siNA molecules of the invention are used to target
5 conserved sequences corresponding to a gene family or gene families. As such, siNA molecules targeting multiple gene targets can provide increased therapeutic effect. In addition, siNA can be used to characterize pathways of gene function in a variety of applications. For example, the present invention can be used to inhibit the activity of target gene(s) in a pathway to determine the function of uncharacterized gene(s) in gene
10 function analysis, mRNA function analysis, or translational analysis. The invention can be used to determine potential target gene pathways involved in various diseases and conditions toward pharmaceutical development. The invention can be used to understand pathways of gene expression involved in, for example, in development, such as prenatal development and postnatal development, and/or the progression and/or maintenance of
15 cancer, infectious disease, autoimmunity, inflammation, endocrine disorders, renal disease, pulmonary disease, cardiovascular disease, birth defects, ageing, any other disease or condition related to gene expression.

In one embodiment, siNA molecule(s) and/or methods of the invention are used to down-regulate or inhibit the expression of gene(s) that encode RNA referred to by
20 Genbank Accession, for example genes encoding RNA sequence(s) referred to herein by Genbank Accession number.

In one embodiment, the invention features a method comprising: (a) generating a library of siNA constructs having a predetermined complexity; and (b) assaying the siNA constructs of (a) above, under conditions suitable to determine RNAi target sites within
25 the target RNA sequence. In one embodiment, the siNA molecules of (a) have strands of a fixed length, for example, about 23 nucleotides in length. In another embodiment, the siNA molecules of (a) are of differing length, for example having strands of about 19 to about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In one embodiment, the assay can comprise a reconstituted *in vitro* siNA assay as described
30 herein. In another embodiment, the assay can comprise a cell culture system in which target RNA is expressed. In another embodiment, fragments of target RNA are analyzed

for detectable levels of cleavage, for example by gel electrophoresis, northern blot analysis, or RNase protection assays, to determine the most suitable target site(s) within the target RNA sequence. The target RNA sequence can be obtained as is known in the art, for example, by cloning and/or transcription for *in vitro* systems, and by cellular expression in *in vivo* systems.

In one embodiment, the invention features a method comprising: (a) generating a randomized library of siNA constructs having a predetermined complexity, such as of 4^N , where N represents the number of base paired nucleotides in each of the siNA construct strands (eg. for a siNA construct having 21 nucleotide sense and antisense strands with 19 base pairs, the complexity would be 4^{19}); and (b) assaying the siNA constructs of (a) above, under conditions suitable to determine RNAi target sites within the target RNA sequence. In another embodiment, the siNA molecules of (a) have strands of a fixed length, for example about 23 nucleotides in length. In yet another embodiment, the siNA molecules of (a) are of differing length, for example having strands of about 19 to about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In one embodiment, the assay can comprise a reconstituted *in vitro* siNA assay as described in Example 7 herein. In another embodiment, the assay can comprise a cell culture system in which target RNA is expressed. In another embodiment, fragments of target RNA are analyzed for detectable levels of cleavage, for example by gel electrophoresis, northern blot analysis, or RNase protection assays, to determine the most suitable target site(s) within the target RNA sequence. In another embodiment, the target RNA sequence can be obtained as is known in the art, for example, by cloning and/or transcription for *in vitro* systems, and by cellular expression in *in vivo* systems.

In another embodiment, the invention features a method comprising: (a) analyzing the sequence of a RNA target encoded by a target gene; (b) synthesizing one or more sets of siNA molecules having sequence complementary to one or more regions of the RNA of (a); and (c) assaying the siNA molecules of (b) under conditions suitable to determine RNAi targets within the target RNA sequence. In one embodiment, the siNA molecules of (b) have strands of a fixed length, for example about 23 nucleotides in length. In another embodiment, the siNA molecules of (b) are of differing length, for example having strands of about 19 to about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In one embodiment, the assay can comprise a reconstituted *in vitro*

siNA assay as described herein. In another embodiment, the assay can comprise a cell culture system in which target RNA is expressed. Fragments of target RNA are analyzed for detectable levels of cleavage, for example by gel electrophoresis, northern blot analysis, or RNase protection assays, to determine the most suitable target site(s) within the target RNA sequence. The target RNA sequence can be obtained as is known in the art, for example, by cloning and/or transcription for *in vitro* systems, and by expression in *in vivo* systems.

By "target site" is meant a sequence within a target RNA that is "targeted" for cleavage mediated by a siNA construct which contains sequences within its antisense region that are complementary to the target sequence.

By "detectable level of cleavage" is meant cleavage of target RNA (and formation of cleaved product RNAs) to an extent sufficient to discern cleavage products above the background of RNAs produced by random degradation of the target RNA. Production of cleavage products from 1-5% of the target RNA is sufficient to detect above the background for most methods of detection.

In one embodiment, the invention features a composition comprising a siNA molecule of the invention, which can be chemically-modified, in a pharmaceutically acceptable carrier or diluent. In another embodiment, the invention features a pharmaceutical composition comprising siNA molecules of the invention, which can be chemically-modified, targeting one or more genes in a pharmaceutically acceptable carrier or diluent. In another embodiment, the invention features a method for diagnosing a disease or condition in a subject comprising administering to the subject a composition of the invention under conditions suitable for the diagnosis of the disease or condition in the subject. In another embodiment, the invention features a method for treating or preventing a disease or condition in a subject, comprising administering to the subject a composition of the invention under conditions suitable for the treatment or prevention of the disease or condition in the subject, alone or in conjunction with one or more other therapeutic compounds. In yet another embodiment, the invention features a method for reducing or preventing tissue rejection in a subject comprising administering to the subject a composition of the invention under conditions suitable for the reduction or prevention of tissue rejection in the subject.

In another embodiment, the invention features a method for validating a gene target, comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands includes a sequence complementary to RNA of a target gene; (b) introducing the siNA molecule into a cell, tissue, or organism under conditions suitable for modulating expression of the target gene in the cell, tissue, or organism; and (c) determining the function of the gene by assaying for any phenotypic change in the cell, tissue, or organism.

In another embodiment, the invention features a method for validating a target gene comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands includes a sequence complementary to RNA of a target gene; (b) introducing the siNA molecule into a biological system under conditions suitable for modulating expression of the target gene in the biological system; and (c) determining the function of the gene by assaying for any phenotypic change in the biological system.

By "biological system" is meant, material, in a purified or unpurified form, from biological sources, including but not limited to human, animal, plant, insect, bacterial, viral or other sources, wherein the system comprises the components required for RNAi activity. The term "biological system" includes, for example, a cell, tissue, or organism, or extract thereof. The term biological system also includes reconstituted RNAi systems that can be used in an *in vitro* setting.

By "phenotypic change" is meant any detectable change to a cell that occurs in response to contact or treatment with a nucleic acid molecule of the invention (e.g., siNA). Such detectable changes include, but are not limited to, changes in shape, size, proliferation, motility, protein expression or RNA expression or other physical or chemical changes as can be assayed by methods known in the art. The detectable change can also include expression of reporter genes/molecules such as Green Florescent Protein (GFP) or various tags that are used to identify an expressed protein or any other cellular component that can be assayed.

In one embodiment, the invention features a kit containing a siNA molecule of the invention, which can be chemically-modified, that can be used to modulate the expression of a target gene in biological system, including, for example, in a cell, tissue, or organism.

In another embodiment, the invention features a kit containing more than one siNA molecule of the invention, which can be chemically-modified, that can be used to modulate the expression of more than one target gene in a biological system, including, for example, in a cell, tissue, or organism.

5 In one embodiment, the invention features a kit containing a siNA molecule of the invention, which can be chemically-modified, that can be used to modulate the expression of a target gene in a biological system. In another embodiment, the invention features a kit containing more than one siNA molecule of the invention, which can be chemically-modified, that can be used to modulate the expression of more than one target gene in a
10 biological system.

In one embodiment, the invention features a cell containing one or more siNA molecules of the invention, which can be chemically-modified. In another embodiment, the cell containing a siNA molecule of the invention is a mammalian cell. In yet another embodiment, the cell containing a siNA molecule of the invention is a human cell.

15 In one embodiment, the synthesis of a siNA molecule of the invention, which can be chemically-modified, comprises: (a) synthesis of two complementary strands of the siNA molecule; (b) annealing the two complementary strands together under conditions suitable to obtain a double-stranded siNA molecule. In another embodiment, synthesis of the two complementary strands of the siNA molecule is by solid phase oligonucleotide
20 synthesis. In yet another embodiment, synthesis of the two complementary strands of the siNA molecule is by solid phase tandem oligonucleotide synthesis.

In one embodiment, the invention features a method for synthesizing a siNA duplex molecule comprising: (a) synthesizing a first oligonucleotide sequence strand of the siNA molecule, wherein the first oligonucleotide sequence strand comprises a cleavable linker
25 molecule that can be used as a scaffold for the synthesis of the second oligonucleotide sequence strand of the siNA; (b) synthesizing the second oligonucleotide sequence strand of siNA on the scaffold of the first oligonucleotide sequence strand, wherein the second oligonucleotide sequence strand further comprises a chemical moiety than can be used to purify the siNA duplex; (c) cleaving the linker molecule of (a) under conditions suitable
30 for the two siNA oligonucleotide strands to hybridize and form a stable duplex; and (d) purifying the siNA duplex utilizing the chemical moiety of the second oligonucleotide

sequence strand. In one embodiment, cleavage of the linker molecule in (c) above takes place during deprotection of the oligonucleotide, for example, under hydrolysis conditions using an alkylamine base such as methylamine. In one embodiment, the method of synthesis comprises solid phase synthesis on a solid support such as controlled pore glass (CPG) or polystyrene, wherein the first sequence of (a) is synthesized on a cleavable linker, such as a succinyl linker, using the solid support as a scaffold. The cleavable linker in (a) used as a scaffold for synthesizing the second strand can comprise similar reactivity as the solid support derivatized linker, such that cleavage of the solid support derivatized linker and the cleavable linker of (a) takes place concomitantly. In another embodiment, the chemical moiety of (b) that can be used to isolate the attached oligonucleotide sequence comprises a trityl group, for example a dimethoxytrityl group, which can be employed in a trityl-on synthesis strategy as described herein. In yet another embodiment, the chemical moiety, such as a dimethoxytrityl group, is removed during purification, for example, using acidic conditions.

In a further embodiment, the method for siNA synthesis is a solution phase synthesis or hybrid phase synthesis wherein both strands of the siNA duplex are synthesized in tandem using a cleavable linker attached to the first sequence which acts a scaffold for synthesis of the second sequence. Cleavage of the linker under conditions suitable for hybridization of the separate siNA sequence strands results in formation of the double-stranded siNA molecule.

In another embodiment, the invention features a method for synthesizing a siNA duplex molecule comprising: (a) synthesizing one oligonucleotide sequence strand of the siNA molecule, wherein the sequence comprises a cleavable linker molecule that can be used as a scaffold for the synthesis of another oligonucleotide sequence; (b) synthesizing a second oligonucleotide sequence having complementarity to the first sequence strand on the scaffold of (a), wherein the second sequence comprises the other strand of the double-stranded siNA molecule and wherein the second sequence further comprises a chemical moiety that can be used to isolate the attached oligonucleotide sequence; (c) purifying the product of (b) utilizing the chemical moiety of the second oligonucleotide sequence strand under conditions suitable for isolating the full-length sequence comprising both siNA oligonucleotide strands connected by the cleavable linker and under conditions suitable for the two siNA oligonucleotide strands to hybridize and form a stable duplex.

In one embodiment, cleavage of the linker molecule in (c) above takes place during deprotection of the oligonucleotide, for example under hydrolysis conditions. In another embodiment, cleavage of the linker molecule in (c) above takes place after deprotection of the oligonucleotide. In another embodiment, the method of synthesis comprises solid
5 phase synthesis on a solid support such as controlled pore glass (CPG) or polystyrene, wherein the first sequence of (a) is synthesized on a cleavable linker, such as a succinyl linker, using the solid support as a scaffold. The cleavable linker in (a) used as a scaffold for synthesizing the second strand can comprise similar reactivity or differing reactivity as the solid support derivatized linker, such that cleavage of the solid support derivatized
10 linker and the cleavable linker of (a) takes place either concomitantly or sequentially. In one embodiment, the chemical moiety of (b) that can be used to isolate the attached oligonucleotide sequence comprises a trityl group, for example a dimethoxytrityl group.

In another embodiment, the invention features a method for making a double-stranded siNA molecule in a single synthetic process comprising: (a) synthesizing an
15 oligonucleotide having a first and a second sequence, wherein the first sequence is complementary to the second sequence, and the first oligonucleotide sequence is linked to the second sequence via a cleavable linker, and wherein a terminal 5'-protecting group, for example, a 5'-O-dimethoxytrityl group (5'-O-DMT) remains on the oligonucleotide having the second sequence; (b) deprotecting the oligonucleotide whereby the
20 deprotection results in the cleavage of the linker joining the two oligonucleotide sequences; and (c) purifying the product of (b) under conditions suitable for isolating the double-stranded siNA molecule, for example using a trityl-on synthesis strategy as described herein.

In another embodiment, the method of synthesis of siNA molecules of the invention
25 comprises the teachings of Scaringe *et al.*, US Patent Nos. 5,889,136; 6,008,400; and 6,111,086, incorporated by reference herein in their entirety.

In one embodiment, the invention features siNA constructs that mediate RNAi in a cell or reconstituted system, wherein the siNA construct comprises one or more chemical modifications, for example, one or more chemical modifications having any of Formulae
30 I-VII or any combination thereof that increases the nuclease resistance of the siNA construct.

In another embodiment, the invention features a method for generating siNA molecules with increased nuclease resistance comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased nuclease resistance.

In one embodiment, the invention features siNA constructs that mediate RNAi against a target gene, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the binding affinity between the sense and antisense strands of the siNA construct.

In one embodiment, the binding affinity between the sense and antisense strands of the siNA construct is modulated to increase the activity of the siNA molecule with regard to the ability of the siNA to mediate RNA interference. In another embodiment the binding affinity between the sense and antisense strands of the siNA construct is decreased. The binding affinity between the sense and antisense strands of the siNA construct can be decreased by introducing one or more chemically modified nucleotides in the siNA sequence that disrupts the duplex stability of the siNA (e.g., lowers the T_m of the duplex). The binding affinity between the sense and antisense strands of the siNA construct can be decreased by introducing one or more nucleotides in the siNA sequence that do not form Watson-Crick base pairs. The binding affinity between the sense and antisense strands of the siNA construct can be decreased by introducing one or more wobble base pairs in the siNA sequence. The binding affinity between the sense and antisense strands of the siNA construct can be decreased by modifying the nucleobase composition of the siNA, such as by altering the G-C content of the siNA sequence (e.g., decreasing the number of G-C base pairs in the siNA sequence). These modifications and alterations in sequence can be introduced selectively at pre-determined positions of the siNA sequence to increase siNA mediated RNAi activity. For example, such modifications and sequence alterations can be introduced to disrupt siNA duplex stability between the 5'-end of the antisense strand and the 3'-end of the sense strand, the 3'-end of the antisense strand and the 5'-end of the sense strand, or alternately the middle of the siNA duplex. In another embodiment, siNA molecules are screened for optimized RNAi activity by introducing such modifications and sequence alterations either by rational design based upon observed rules or trends in increasing siNA activity, or randomly via

combinatorial selection processes that cover either partial or complete sequence space of the siNA construct.

In another embodiment, the invention features a method for generating siNA molecules with increased binding affinity between the sense and antisense strands of the siNA molecule comprising (a) introducing nucleotides having any of Formula I-VII or
5 any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased binding affinity between the sense and antisense strands of the siNA molecule.

In one embodiment, the invention features siNA constructs that mediate RNAi in a
10 cell or reconstituted system, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the binding affinity between the antisense strand of the siNA construct and a complementary target RNA sequence within a cell.

In one embodiment, the invention features siNA constructs that mediate RNAi in a cell or reconstituted system, wherein the siNA construct comprises one or more chemical
15 modifications described herein that modulates the binding affinity between the antisense strand of the siNA construct and a complementary target DNA sequence within a cell.

In another embodiment, the invention features a method for generating siNA molecules with increased binding affinity between the antisense strand of the siNA molecule and a complementary target RNA sequence comprising (a) introducing
20 nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased binding affinity between the antisense strand of the siNA molecule and a complementary target RNA sequence.

In another embodiment, the invention features a method for generating siNA
25 molecules with increased binding affinity between the antisense strand of the siNA molecule and a complementary target DNA sequence comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased binding affinity between the antisense strand
30 of the siNA molecule and a complementary target DNA sequence.

In one embodiment, the invention features siNA constructs that mediate RNAi in a cell or reconstituted system, wherein the siNA construct comprises one or more chemical modifications described herein that modulate the polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to the chemically-modified siNA construct.

In another embodiment, the invention features a method for generating siNA molecules capable of mediating increased polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to a chemically-modified siNA molecule comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules capable of mediating increased polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to the chemically-modified siNA molecule. In one embodiment, the invention features chemically-modified siNA constructs that mediate RNAi in a cell or reconstituted system, wherein the chemical modifications do not significantly effect the interaction of siNA with a target RNA molecule, DNA molecule and/or proteins or other factors that are essential for RNAi in a manner that would decrease the efficacy of RNAi mediated by such siNA constructs.

In another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity.

In yet another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity against a target RNA comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity against the target RNA.

In yet another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity against a DNA target comprising (a) introducing

nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity against the DNA target, such as a gene, chromosome, or portion thereof.

5 In one embodiment, the invention features siNA constructs that mediate RNAi in a cell or reconstituted system, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the cellular uptake of the siNA construct.

 In another embodiment, the invention features a method for generating siNA molecules against a target gene with improved cellular uptake comprising (a) introducing
10 nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved cellular uptake.

 In one embodiment, the invention features siNA constructs that mediate RNAi against a target gene, wherein the siNA construct comprises one or more chemical
15 modifications described herein that increases the bioavailability of the siNA construct, for example, by attaching polymeric conjugates such as polyethyleneglycol or equivalent conjugates that improve the pharmacokinetics of the siNA construct, or by attaching conjugates that target specific tissue types or cell types *in vivo*. Non-limiting examples of such conjugates are described in Vargeese *et al.*, U.S. Serial No. 10/201,394 incorporated
20 by reference herein.

 In one embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability comprising (a) introducing a conjugate into the structure of a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved bioavailability. Such
25 conjugates can include ligands for cellular receptors, such as peptides derived from naturally occurring protein ligands; protein localization sequences, including cellular ZIP code sequences; antibodies; nucleic acid aptamers; vitamins and other co-factors, such as folate and N-acetylgalactosamine; polymers, such as polyethyleneglycol (PEG); phospholipids; cholesterol; polyamines, such as spermine or spermidine; and others.

In one embodiment, the invention features a double stranded short interfering nucleic acid (siNA) molecule that comprises a first nucleotide sequence complementary to a target RNA sequence or a portion thereof, and a second sequence having complementarity to said first sequence, wherein said second sequence is chemically
5 modified in a manner that it can no longer act as a guide sequence for efficiently mediating RNA interference and/or is recognized by cellular proteins that facilitate RNAi.

In one embodiment, the invention features a double stranded short interfering nucleic acid (siNA) molecule that comprises a first nucleotide sequence complementary to a target RNA sequence or a portion thereof, and a second sequence having
10 complementarity to said first sequence, wherein the second sequence is designed or modified in a manner that prevents its entry into the RNAi pathway as a guide sequence or as a sequence that is complementary to a target nucleic acid (e.g., RNA) sequence. Such design or modifications are expected to enhance the activity of siNA and/or improve the specificity of siNA molecules of the invention. These modifications are also expected
15 to minimize any off-target effects and/or associated toxicity.

In one embodiment, the invention features a double stranded short interfering nucleic acid (siNA) molecule that comprises a first nucleotide sequence complementary to a target RNA sequence or a portion thereof, and a second sequence having complementarity to said first sequence, wherein said second sequence is incapable of
20 acting as a guide sequence for mediating RNA interference.

In one embodiment, the invention features a double stranded short interfering nucleic acid (siNA) molecule that comprises a first nucleotide sequence complementary to a target RNA sequence or a portion thereof, and a second sequence having complementarity to said first sequence, wherein said second sequence does not have a
25 terminal 5'-hydroxyl (5'-OH) or 5'-phosphate group.

In one embodiment, the invention features a double stranded short interfering nucleic acid (siNA) molecule that comprises a first nucleotide sequence complementary to a target RNA sequence or a portion thereof, and a second sequence having complementarity to said first sequence, wherein said second sequence comprises a
30 terminal cap moiety at the 5'-end of said second sequence. In another embodiment, the terminal cap moiety comprises an inverted abasic, inverted deoxy abasic, inverted

nucleotide moiety, a group shown in **Figure 22**, an alkyl or cycloalkyl group, a heterocycle, or any other group that prevents RNAi activity in which the second sequence serves as a guide sequence or template for RNAi.

In one embodiment, the invention features a double stranded short interfering
5 nucleic acid (siNA) molecule that comprises a first nucleotide sequence complementary to a target RNA sequence or a portion thereof, and a second sequence having complementarity to said first sequence, wherein said second sequence comprises a terminal cap moiety at the 5'-end and 3'-end of said second sequence. In another embodiment, each terminal cap moiety individually comprises an inverted abasic,
10 inverted deoxy abasic, inverted nucleotide moiety, a group shown in **Figure 22**, an alkyl or cycloalkyl group, a heterocycle, or any other group that prevents RNAi activity in which the second sequence serves as a guide sequence or template for RNAi.

In one embodiment, the invention features a method for generating siNA molecules of the invention with improved specificity for down regulating or inhibiting the
15 expression of a target nucleic acid (e.g., a DNA or RNA such as a gene or its corresponding RNA), comprising (a) introducing one or more chemical modifications into the structure of a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved specificity. In another embodiment, the chemical modification used to improve specificity comprises
20 terminal cap modifications at the 5'-end, 3'-end, or both 5' and 3'-ends of the siNA molecule. The terminal cap modifications can comprise, for example, structures shown in **Figure 22** (e.g. inverted deoxyabasic moieties) or any other chemical modification that renders a portion of the siNA molecule (e.g. the sense strand) incapable of mediating RNA interference against an off target nucleic acid sequence. In a non-limiting example,
25 a siNA molecule is designed such that only the antisense sequence of the siNA molecule can serve as a guide sequence for RISC mediated degradation of a corresponding target RNA sequence. This can be accomplished by rendering the sense sequence of the siNA inactive by introducing chemical modifications to the sense strand that preclude recognition of the sense strand as a guide sequence by RNAi machinery. In one
30 embodiment, such chemical modifications comprise any chemical group at the 5'-end of the sense strand of the siNA, or any other group that serves to render the sense strand inactive as a guide sequence for mediating RNA interference. These modifications, for

example, can result in a molecule where the 5'-end of the sense strand no longer has a free 5'-hydroxyl (5'-OH) or a free 5'-phosphate group (e.g., phosphate, diphosphate, triphosphate, cyclic phosphate etc.). Non-limiting examples of such siNA constructs are described herein, such as "Stab 9/10" and "Stab 7/8" chemistries and variants thereof
5 wherein the 5'-end and 3'-end of the sense strand of the siNA do not comprise a hydroxyl group or phosphate group.

In one embodiment, the invention features a method for generating siNA molecules of the invention with improved specificity for down regulating or inhibiting the expression of a target nucleic acid (e.g., a DNA or RNA such as a gene or its
10 corresponding RNA), comprising (a) introducing one or more chemical modifications into the structure of a siNA molecule that prevent a strand or portion of the siNA molecule from acting as a template or guide sequence for RNAi activity. In another embodiment, the inactive strand or sense region of the siNA molecule is the sense strand or sense region of the siNA molecule, i.e. the strand or region of the siNA that does not
15 have complementarity to the target nucleic acid sequence. In one embodiment, such chemical modifications comprise any chemical group at the 5'-end of the sense strand or region of the siNA that does not comprise a 5'-hydroxyl (5'-OH) or 5'-phosphate group, or any other group that serves to render the sense strand or sense region inactive as a guide sequence for mediating RNA interference. Non-limiting examples of such siNA
20 constructs are described herein, such as "Stab 9/10" and "Stab 7/8" chemistries and variants thereof wherein the 5'-end and 3'-end of the sense strand of the siNA do not comprise a hydroxyl group or phosphate group.

In one embodiment, the invention features a method for screening siNA molecules against a target nucleic acid sequence comprising, (a) generating a plurality of
25 unmodified siNA molecules, (b) assaying the siNA molecules of step (a) under conditions suitable for isolating siNA molecules that are active in mediating RNA interference against the target nucleic acid sequence, (c) introducing chemical modifications (e.g. chemical modifications as described herein or as otherwise known in the art) into the active siNA molecules of (b), and (d) optionally re-screening the chemically modified
30 siNA molecules of (c) under conditions suitable for isolating chemically modified siNA molecules that are active in mediating RNA interference against the target nucleic acid sequence.

In one embodiment, the invention features a method for screening siNA molecules against a target nucleic acid sequence comprising, (a) generating a plurality of chemically modified siNA molecules (e.g. siNA molecules as described herein or as otherwise known in the art), and (b) assaying the siNA molecules of step (a) under conditions
5 suitable for isolating chemically modified siNA molecules that are active in mediating RNA interference against the target nucleic acid sequence.

In another embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability comprising (a) introducing an excipient formulation to a siNA molecule, and (b) assaying the siNA molecule of step (a)
10 under conditions suitable for isolating siNA molecules having improved bioavailability. Such excipients include polymers such as cyclodextrins, lipids, cationic lipids, polyamines, phospholipids, nanoparticles, receptors, ligands, and others.

In another embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability comprising (a) introducing an excipient formulation to a siNA molecule, and (b) assaying the siNA molecule of step (a)
15 under conditions suitable for isolating siNA molecules having improved bioavailability. Such excipients include polymers such as cyclodextrins, lipids, cationic lipids, polyamines, phospholipids, and others.

In another embodiment, the invention features a method for generating siNA
20 molecules of the invention with improved bioavailability comprising (a) introducing nucleotides having any of Formulae I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved bioavailability.

In another embodiment, polyethylene glycol (PEG) can be covalently attached to
25 siNA compounds of the present invention. The attached PEG can be any molecular weight, preferably from about 2,000 to about 50,000 daltons (Da).

The present invention can be used alone or as a component of a kit having at least one of the reagents necessary to carry out the *in vitro* or *in vivo* introduction of RNA to test samples and/or subjects. For example, preferred components of the kit include a
30 siNA molecule of the invention and a vehicle that promotes introduction of the siNA into

cells of interest as described herein (e.g., using lipids and other methods of transfection known in the art, see for example Beigelman *et al.*, US 6,395,713). The kit can be used for target validation, such as in determining gene function and/or activity, or in drug optimization, and in drug discovery (see for example Usman *et al.*, USSN 60/402,996).

5 Such a kit can also include instructions to allow a user of the kit to practice the invention.

The term "short interfering nucleic acid", "siNA", "short interfering RNA", "siRNA", "short interfering nucleic acid molecule", "short interfering oligonucleotide molecule", or "chemically-modified short interfering nucleic acid molecule" as used herein refers to any nucleic acid molecule capable of inhibiting or down regulating gene
 10 expression or viral replication, for example by mediating RNA interference "RNAi" or gene silencing in a sequence-specific manner; see for example Zamore *et al.*, 2000, *Cell*, 101, 25-33; Bass, 2001, *Nature*, 411, 428-429; Elbashir *et al.*, 2001, *Nature*, 411, 494-498; and Kreutzer *et al.*, International PCT Publication No. WO 00/44895; Zernicka-Goetz *et al.*, International PCT Publication No. WO 01/36646; Fire, International PCT
 15 Publication No. WO 99/32619; Plaetinck *et al.*, International PCT Publication No. WO 00/01846; Mello and Fire, International PCT Publication No. WO 01/29058; Deschamps-Depaillette, International PCT Publication No. WO 99/07409; and Li *et al.*, International PCT Publication No. WO 00/44914; Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 2002, *Science*, 297, 2232-2237; Hutvagner and Zamore, 2002, *Science*, 297, 2056-60; McManus *et al.*, 2002, *RNA*, 8, 842-850; Reinhart *et al.*, 2002, *Gene & Dev.*, 16, 1616-1626; and Reinhart & Bartel, 2002, *Science*, 297, 1831). Non limiting examples of siNA molecules of the invention are shown in **Figures 18-20**, and **Table I** herein. For example the siNA can be a double-stranded polynucleotide molecule comprising self-
 25 complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. The siNA can be assembled from two separate oligonucleotides, where one strand is the sense strand and
 30 the other is the antisense strand, wherein the antisense and sense strands are self-complementary (i.e. each strand comprises nucleotide sequence that is complementary to nucleotide sequence in the other strand; such as where the antisense strand and sense

strand form a duplex or double stranded structure, for example wherein the double stranded region is about 19 base pairs); the antisense strand comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense strand comprises nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. Alternatively, the siNA is assembled from a single oligonucleotide, where the self-complementary sense and antisense regions of the siNA are linked by means of a nucleic acid based or non-nucleic acid-based linker(s). The siNA can be a polynucleotide with a duplex, asymmetric duplex, hairpin or asymmetric hairpin secondary structure, having self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a separate target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. The siNA can be a circular single-stranded polynucleotide having two or more loop structures and a stem comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof, and wherein the circular polynucleotide can be processed either *in vivo* or *in vitro* to generate an active siNA molecule capable of mediating RNAi. The siNA can also comprise a single stranded polynucleotide having nucleotide sequence complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof (for example, where such siNA molecule does not require the presence within the siNA molecule of nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof), wherein the single stranded polynucleotide can further comprise a terminal phosphate group, such as a 5'-phosphate (see for example Martinez *et al.*, 2002, *Cell.*, 110, 563-574 and Schwarz *et al.*, 2002, *Molecular Cell*, 10, 537-568), or 5',3'-diphosphate. In certain embodiments, the siNA molecule of the invention comprises separate sense and antisense sequences or regions, wherein the sense and antisense regions are covalently linked by nucleotide or non-nucleotide linker molecules as is known in the art, or are alternately non-covalently linked by ionic interactions, hydrogen bonding, van der Waals interactions, hydrophobic interactions, and/or stacking interactions. In certain embodiments, the siNA molecules of the invention comprise nucleotide sequence that is

complementary to nucleotide sequence of a target gene. In another embodiment, the siNA molecule of the invention interacts with nucleotide sequence of a target gene in a manner that causes inhibition of expression of the target gene. As used herein, siNA molecules need not be limited to those molecules containing only RNA, but further encompasses chemically-modified nucleotides and non-nucleotides. In certain embodiments, the short interfering nucleic acid molecules of the invention lack 2'-hydroxy (2'-OH) containing nucleotides. Applicant describes in certain embodiments short interfering nucleic acids that do not require the presence of nucleotides having a 2'-hydroxy group for mediating RNAi and as such, short interfering nucleic acid molecules of the invention optionally do not include any ribonucleotides (e.g., nucleotides having a 2'-OH group). Such siNA molecules that do not require the presence of ribonucleotides within the siNA molecule to support RNAi can however have an attached linker or linkers or other attached or associated groups, moieties, or chains containing one or more nucleotides with 2'-OH groups. Optionally, siNA molecules can comprise ribonucleotides at about 5, 10, 20, 30, 40, or 50% of the nucleotide positions. The modified short interfering nucleic acid molecules of the invention can also be referred to as short interfering modified oligonucleotides "siMON." As used herein, the term siNA is meant to be equivalent to other terms used to describe nucleic acid molecules that are capable of mediating sequence specific RNAi, for example short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), short hairpin RNA (shRNA), short interfering oligonucleotide, short interfering nucleic acid, short interfering modified oligonucleotide, chemically-modified siRNA, post-transcriptional gene silencing RNA (ptgsRNA), and others. In addition, as used herein, the term RNAi is meant to be equivalent to other terms used to describe sequence specific RNA interference, such as post transcriptional gene silencing, translational inhibition, or epigenetics. For example, siNA molecules of the invention can be used to epigenetically silence genes at both the post-transcriptional level or the pre-transcriptional level. In a non-limiting example, epigenetic regulation of gene expression by siNA molecules of the invention can result from siNA mediated modification of chromatin structure to alter gene expression (see, for example, Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 2002, *Science*, 297, 2232-2237).

By “asymmetric hairpin” as used herein is meant a linear siNA molecule comprising an antisense region, a loop portion that can comprise nucleotides or non-nucleotides, and a sense region that comprises fewer nucleotides than the antisense region to the extent that the sense region has enough complimentary nucleotides to base pair with the antisense region and form a duplex with loop. For example, an asymmetric hairpin siNA molecule of the invention can comprise an antisense region having length sufficient to mediate RNAi in a cell or in vitro system (e.g. about 19 to about 22 nucleotides) and a loop region comprising about 4 to about 8 nucleotides, and a sense region having about 3 to about 18 nucleotides that are complementary to the antisense region (see for example **Figure 74**). The asymmetric hairpin siNA molecule can also comprise a 5'-terminal phosphate group that can be chemically modified (for example as shown in **Figure 75**). The loop portion of the asymmetric hairpin siNA molecule can comprise nucleotides, non-nucleotides, linker molecules, or conjugate molecules as described herein.

By “asymmetric duplex” as used herein is meant a siNA molecule having two separate strands comprising a sense region and an antisense region, wherein the sense region comprises fewer nucleotides than the antisense region to the extent that the sense region has enough complimentary nucleotides to base pair with the antisense region and form a duplex. For example, an asymmetric duplex siNA molecule of the invention can comprise an antisense region having length sufficient to mediate RNAi in a cell or in vitro system (e.g. about 19 to about 22 nucleotides) and a sense region having about 3 to about 18 nucleotides that are complementary to the antisense region (see for example **Figure 74**).

By “modulate” is meant that the expression of the gene, or level of RNA molecule or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity of one or more proteins or protein subunits is up regulated or down regulated, such that expression, level, or activity is greater than or less than that observed in the absence of the modulator. For example, the term “modulate” can mean “inhibit,” but the use of the word “modulate” is not limited to this definition.

By “inhibit”, “down-regulate”, or “reduce”, it is meant that the expression of the gene, or level of RNA molecules or equivalent RNA molecules encoding one or more

proteins or protein subunits, or activity of one or more proteins or protein subunits, is reduced below that observed in the absence of the nucleic acid molecules (e.g., siNA) of the invention. In one embodiment, inhibition, down-regulation or reduction with an siNA molecule is below that level observed in the presence of an inactive or attenuated molecule. In another embodiment, inhibition, down-regulation, or reduction with siNA molecules is below that level observed in the presence of, for example, an siNA molecule with scrambled sequence or with mismatches. In another embodiment, inhibition, down-regulation, or reduction of gene expression with a nucleic acid molecule of the instant invention is greater in the presence of the nucleic acid molecule than in its absence.

By "gene", or "target gene", is meant, a nucleic acid that encodes an RNA, for example, nucleic acid sequences including, but not limited to, structural genes encoding a polypeptide. A gene or target gene can also encode a functional RNA (fRNA) or non-coding RNA (ncRNA), such as small temporal RNA (stRNA), micro RNA (miRNA), small nuclear RNA (snRNA), short interfering RNA (siRNA), small nucleolar RNA (snRNA), ribosomal RNA (rRNA), transfer RNA (tRNA) and precursor RNAs thereof. Such non-coding RNAs can serve as target nucleic acid molecules for siNA mediated RNA interference in modulating the activity of fRNA or ncRNA involved in functional or regulatory cellular processes. Abberant fRNA or ncRNA activity leading to disease can therefore be modulated by siNA molecules of the invention. siNA molecules targeting fRNA and ncRNA can also be used to manipulate or alter the genotype or phenotype of an organism or cell, by intervening in cellular processes such as genetic imprinting, transcription, translation, or nucleic acid processing (e.g., transamination, methylation etc.). The target gene can be a gene derived from a cell, an endogenous gene, a transgene, or exogenous genes such as genes of a pathogen, for example a virus, which is present in the cell after infection thereof. The cell containing the target gene can be derived from or contained in any organism, for example a plant, animal, protozoan, virus, bacterium, or fungus. Non-limiting examples of plants include monocots, dicots, or gymnosperms. Non-limiting examples of animals include vertebrates or invertebrates. Non-limiting examples of fungi include molds or yeasts.

By "highly conserved sequence region" is meant, a nucleotide sequence of one or more regions in a target gene does not vary significantly from one generation to the other or from one biological system to the other.

By "cancer" is meant a group of diseases characterized by uncontrolled growth and spread of abnormal cells.

By "sense region" is meant a nucleotide sequence of a siNA molecule having complementarity to an antisense region of the siNA molecule. In addition, the sense region of a siNA molecule can comprise a nucleic acid sequence having homology with a target nucleic acid sequence.

By "antisense region" is meant a nucleotide sequence of a siNA molecule having complementarity to a target nucleic acid sequence. In addition, the antisense region of a siNA molecule can optionally comprise a nucleic acid sequence having complementarity to a sense region of the siNA molecule.

By "target nucleic acid" is meant any nucleic acid sequence whose expression or activity is to be modulated. The target nucleic acid can be DNA or RNA, such as endogenous DNA or RNA, viral DNA or viral RNA, or other RNA encoded by a gene, virus, bacteria, fungus, mammal, or plant.

By "complementarity" is meant that a nucleic acid can form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick or other non-traditional types. In reference to the nucleic molecules of the present invention, the binding free energy for a nucleic acid molecule with its complementary sequence is sufficient to allow the relevant function of the nucleic acid to proceed, e.g., RNAi activity. Determination of binding free energies for nucleic acid molecules is well known in the art (see, e.g., Turner *et al.*, 1987, *CSH Symp. Quant. Biol.* LII pp.123-133; Frier *et al.*, 1986, *Proc. Nat. Acad. Sci. USA* 83:9373-9377; Turner *et al.*, 1987, *J. Am. Chem. Soc.* 109:3783-3785). A percent complementarity indicates the percentage of contiguous residues in a nucleic acid molecule that can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, or 10 nucleotides out of a total of 10 nucleotides in the first oligonucleotide being base paired to a second nucleic acid sequence having 10 nucleotides represents 50%, 60%, 70%, 80%, 90%, and 100% complementary respectively). "Perfectly complementary" means that all the contiguous residues of a nucleic acid sequence will hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence.

The siNA molecules of the invention represent a novel therapeutic approach to a broad spectrum of diseases and conditions, including cancer or cancerous disease, infectious disease, cardiovascular disease, neurological disease, prion disease, inflammatory disease, autoimmune disease, pulmonary disease, renal disease, liver
 5 disease, mitochondrial disease, endocrine disease, reproduction related diseases and conditions, and any other indications that can respond to the level of an expressed gene product in a cell or organism.

In one embodiment of the present invention, each sequence of a siNA molecule of the invention is independently about 18 to about 24 nucleotides in length, in specific
 10 embodiments about 18, 19, 20, 21, 22, 23, or 24 nucleotides in length. In another embodiment, the siNA duplexes of the invention independently comprise about 17 to about 23 base pairs (*e.g.*, about 17, 18, 19, 20, 21, 22 or 23). In yet another embodiment, siNA molecules of the invention comprising hairpin or circular structures are about 35 to about 55 (*e.g.*, about 35, 40, 45, 50 or 55) nucleotides in length, or about 38 to about 44
 15 (*e.g.*, 38, 39, 40, 41, 42, 43 or 44) nucleotides in length and comprising about 16 to about 22 (*e.g.*, about 16, 17, 18, 19, 20, 21 or 22) base pairs. Exemplary siNA molecules of the invention are shown in **Table I.** and/or **Figures 18-19.**

As used herein "cell" is used in its usual biological sense, and does not refer to an entire multicellular organism, *e.g.*, specifically does not refer to a human. The cell can be
 20 present in an organism, *e.g.*, birds, plants and mammals such as humans, cows, sheep, apes, monkeys, swine, dogs, and cats. The cell can be prokaryotic (*e.g.*, bacterial cell) or eukaryotic (*e.g.*, mammalian or plant cell). The cell can be of somatic or germ line origin, totipotent or pluripotent, dividing or non-dividing. The cell can also be derived from or can comprise a gamete or embryo, a stem cell, or a fully differentiated cell.

The siNA molecules of the invention are added directly, or can be complexed with cationic lipids, packaged within liposomes, or otherwise delivered to target cells or tissues. The nucleic acid or nucleic acid complexes can be locally administered to relevant tissues *ex vivo*, or *in vivo* through injection, infusion pump or stent, with or without their incorporation in biopolymers. In particular embodiments, the nucleic acid
 25 molecules of the invention comprise sequences shown in **Table I** and/or **Figures 18-19.** Examples of such nucleic acid molecules consist essentially of sequences defined in these
 30

tables and figures. Furthermore, the chemically modified constructs described in **Table IV** can be applied to any siNA sequence of the invention.

In another aspect, the invention provides mammalian cells containing one or more siNA molecules of this invention. The one or more siNA molecules can independently be
5 targeted to the same or different sites.

By "RNA" is meant a molecule comprising at least one ribonucleotide residue. By "ribonucleotide" is meant a nucleotide with a hydroxyl group at the 2' position of a β -D-ribo-furanose moiety. The terms include double-stranded RNA, single-stranded RNA, isolated RNA such as partially purified RNA, essentially pure RNA, synthetic RNA,
10 recombinantly produced RNA, as well as altered RNA that differs from naturally occurring RNA by the addition, deletion, substitution and/or alteration of one or more nucleotides. Such alterations can include addition of non-nucleotide material, such as to the end(s) of the siNA or internally, for example at one or more nucleotides of the RNA. Nucleotides in the RNA molecules of the instant invention can also comprise non-
15 standard nucleotides, such as non-naturally occurring nucleotides or chemically synthesized nucleotides or deoxynucleotides. These altered RNAs can be referred to as analogs or analogs of naturally-occurring RNA.

By "subject" is meant an organism, which is a donor or recipient of explanted cells or the cells themselves. "Subject" also refers to an organism to which the nucleic acid
20 molecules of the invention can be administered. A subject can be a mammal or mammalian cells, including a human or human cells.

The term "ligand" refers to any compound or molecule, such as a drug, peptide, hormone, or neurotransmitter, that is capable of interacting with another compound, such as a receptor, either directly or indirectly. The receptor that interacts with a ligand can be
25 present on the surface of a cell or can alternately be an intercellular receptor. Interaction of the ligand with the receptor can result in a biochemical reaction, or can simply be a physical interaction or association.

The term "phosphorothioate" as used herein refers to an internucleotide linkage having Formula I, wherein Z and/or W comprise a sulfur atom. Hence, the term

phosphorothioate refers to both phosphorothioate and phosphorodithioate internucleotide linkages.

The term "phosphonoacetate" as used herein refers to an internucleotide linkage having Formula I, wherein Z and/or W comprise an acetyl or protected acetyl group.

- 5 The term "thiophosphonoacetate" as used herein refers to an internucleotide linkage having Formula I, wherein Z comprises an acetyl or protected acetyl group and W comprises a sulfur atom or alternately W comprises an acetyl or protected acetyl group and Z comprises a sulfur atom.

- 10 The term "universal base" as used herein refers to nucleotide base analogs that form base pairs with each of the natural DNA/RNA bases with little discrimination between them. Non-limiting examples of universal bases include C-phenyl, C-naphthyl and other aromatic derivatives, inosine, azole carboxamides, and nitroazole derivatives such as 3-nitropyrrole, 4-nitroindole, 5-nitroindole, and 6-nitroindole as known in the art (see for example Loakes, 2001, *Nucleic Acids Research*, 29, 2437-2447).

- 15 The term "acyclic nucleotide" as used herein refers to any nucleotide having an acyclic ribose sugar, for example where any of the ribose carbons (C1, C2, C3, C4, or C5), are independently or in combination absent from the nucleotide.

- 20 The nucleic acid molecules of the instant invention, individually, or in combination or in conjunction with other drugs, can be used to treat diseases or conditions discussed herein (e.g., cancers and othe proliferative conditions, viral infection, inflammatory disease, autoimmunity, pulmonary disease, renal disease, ocular disease, etc.). For example, to treat a particular disease or condition, the siNA molecules can be administered to a subject or can be administered to other appropriate cells evident to those skilled in the art, individually or in combination with one or more drugs under conditions
25 suitable for the treatment.

In one embodiment, the invention features a method for treating or preventing a disease or condition in a subject, wherein the disease or condition is related to angiogenesis or neovascularization, comprising administering to the subject a siNA molecule of the invention under conditions suitable for the treatment or prevention of the

disease or condition in the subject, alone or in conjunction with one or more other therapeutic compounds. In another embodiment, the disease or condition comprises tumor angiogenesis and cancer, including but not limited to breast cancer, lung cancer (including non-small cell lung carcinoma), prostate cancer, colorectal cancer, brain cancer, esophageal cancer, bladder cancer, pancreatic cancer, cervical cancer, head and neck cancer, skin cancers, nasopharyngeal carcinoma, liposarcoma, epithelial carcinoma, renal cell carcinoma, gallbladder adeno carcinoma, parotid adenocarcinoma, ovarian cancer, melanoma, lymphoma, glioma, endometrial sarcoma, multidrug resistant cancers, diabetic retinopathy, macular degeneration, age related macular degeneration, neovascular glaucoma, myopic degeneration, arthritis, psoriasis, endometriosis, female reproduction, verruca vulgaris, angiofibroma of tuberous sclerosis, pot-wine stains, Sturge Weber syndrome, Kippel-Trenaunay-Weber syndrome, Osler-Weber-Rendu syndrome, renal disease such as Autosomal dominant polycystic kidney disease (ADPKD), restenosis, arteriosclerosis, and any other diseases or conditions that are related to gene expression or will respond to RNA interference in a cell or tissue, alone or in combination with other therapies.

In one embodiment, the invention features a method for treating or preventing an ocular disease or condition in a subject, wherein the ocular disease or condition is related to angiogenesis or neovascularization, comprising administering to the subject a siNA molecule of the invention under conditions suitable for the treatment or prevention of the disease or condition in the subject, alone or in conjunction with one or more other therapeutic compounds. In another embodiment, the ocular disease or condition comprises macular degeneration, age related macular degeneration, diabetic retinopathy, neovascular glaucoma, myopic degeneration, trachoma, scarring of the eye, cataract, ocular inflammation and/or ocular infections.

In one embodiment, the invention features a method for treating or preventing tumor angiogenesis in a subject, comprising administering to the subject a siNA molecule of the invention under conditions suitable for the treatment or prevention of tumor angiogenesis in the subject, alone or in conjunction with one or more other therapeutic compounds.

In one embodiment, the invention features a method for treating or preventing viral infection or replication in a subject, comprising administering to the subject a siNA molecule of the invention under conditions suitable for the treatment or prevention of viral infection or replication in the subject, alone or in conjunction with one or more other therapeutic compounds.

In one embodiment, the invention features a method for treating or preventing autoimmune disease in a subject, comprising administering to the subject a siNA molecule of the invention under conditions suitable for the treatment or prevention of autoimmune disease in the subject, alone or in conjunction with one or more other therapeutic compounds.

In one embodiment, the invention features a method for treating or preventing inflammation in a subject, comprising administering to the subject a siNA molecule of the invention under conditions suitable for the treatment or prevention of inflammation in the subject, alone or in conjunction with one or more other therapeutic compounds.

In a further embodiment, the siNA molecules can be used in combination with other known treatments to treat conditions or diseases discussed above. For example, the described molecules could be used in combination with one or more known therapeutic agents to treat a disease or condition. Non-limiting examples of other therapeutic agents that can be readily combined with a siNA molecule of the invention are enzymatic nucleic acid molecules, allosteric nucleic acid molecules, antisense, decoy, or aptamer nucleic acid molecules, antibodies such as monoclonal antibodies, small molecules, and other organic and/or inorganic compounds including metals, salts and ions.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a non-limiting example of a scheme for the synthesis of siNA molecules. The complementary siNA sequence strands, strand 1 and strand 2, are synthesized in tandem and are connected by a cleavable linkage, such as a nucleotide succinate or abasic succinate, which can be the same or different from the cleavable

linker used for solid phase synthesis on a solid support. The synthesis can be either solid phase or solution phase, in the example shown, the synthesis is a solid phase synthesis. The synthesis is performed such that a protecting group, such as a dimethoxytrityl group, remains intact on the terminal nucleotide of the tandem oligonucleotide. Upon cleavage and deprotection of the oligonucleotide, the two siNA strands spontaneously hybridize to form a siNA duplex, which allows the purification of the duplex by utilizing the properties of the terminal protecting group, for example by applying a trityl on purification method wherein only duplexes/oligonucleotides with the terminal protecting group are isolated.

10 **Figure 2** shows a MALDI-TOF mass spectrum of a purified siNA duplex synthesized by a method of the invention. The two peaks shown correspond to the predicted mass of the separate siNA sequence strands. This result demonstrates that the siNA duplex generated from tandem synthesis can be purified as a single entity using a simple trityl-on purification methodology.

15 **Figure 3** shows the results of a stability assay used to determine the serum stability of chemically modified siNA constructs compared to a siNA control consisting of all RNA with 3'-TT termini. T $\frac{1}{2}$ values are shown for duplex stability.

Figure 4 shows the results of an RNAi activity screen of several phosphorothioate modified siNA constructs using a luciferase reporter system.

20 **Figure 5** shows the results of an RNAi activity screen of several phosphorothioate and universal base modified siNA constructs using a luciferase reporter system.

Figure 6 shows the results of an RNAi activity screen of several 2'-O-methyl modified siNA constructs using a luciferase reporter system.

25 **Figure 7** shows the results of an RNAi activity screen of several 2'-O-methyl and 2'-deoxy-2'-fluoro modified siNA constructs using a luciferase reporter system.

Figure 8 shows the results of an RNAi activity screen of a phosphorothioate modified siNA construct using a luciferase reporter system.

Figure 9 shows the results of an RNAi activity screen of an inverted deoxyabasic modified siNA construct generated via tandem synthesis using a luciferase reporter system.

Figure 10 shows the results of an RNAi activity screen of chemically modified siNA constructs including 3'-glyceryl modified siNA constructs compared to an all RNA control siNA construct using a luciferase reporter system. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I.

Figure 11 shows the results of an RNAi activity screen of chemically modified siNA constructs. The screen compared various combinations of sense strand chemical modifications and antisense strand chemical modifications. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I.

Figure 12 shows the results of an RNAi activity screen of chemically modified siNA constructs. The screen compared various combinations of sense strand chemical modifications and antisense strand chemical modifications. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number

(sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I. In addition, the antisense strand alone (Sirna/RPI 30430) and an inverted control (Sirna/RPI 30227/30229, having matched chemistry to Sirna/RPI (30063/30224) was compared to the siNA duplexes described above.

5 **Figure 13** shows the results of an RNAi activity screen of chemically modified siNA constructs. The screen compared various combinations of sense strand chemical modifications and antisense strand chemical modifications. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine
10 (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I. In addition, an inverted control (Sirna/RPI 30226/30229), having
15 matched chemistry to Sirna/RPI (30222/30224) was compared to the siNA duplexes described above.

Figure 14 shows the results of an RNAi activity screen of chemically modified siNA constructs including various 3'-terminal modified siNA constructs compared to an all RNA control siNA construct using a luciferase reporter system. These chemically
20 modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by
25 Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I.

Figure 15 shows the results of an RNAi activity screen of chemically modified siNA constructs. The screen compared various combinations of sense strand chemistries compared to a fixed antisense strand chemistry. These chemically modified siNAs were
30 compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its

corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the “cells” column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I.

Figure 16 shows the results of a siNA titration study using a luciferase reporter system, wherein the RNAi activity of a phosphorothioate modified siNA construct is compared to that of a siNA construct consisting of all ribonucleotides except for two terminal thymidine residues.

Figure 17 shows a non-limiting proposed mechanistic representation of target RNA degradation involved in RNAi. Double-stranded RNA (dsRNA), which is generated by RNA-dependent RNA polymerase (RdRP) from foreign single-stranded RNA, for example viral, transposon, or other exogenous RNA, activates the DICER enzyme that in turn generates siNA duplexes. Alternately, synthetic or expressed siNA can be introduced directly into a cell by appropriate means. An active siNA complex forms which recognizes a target RNA, resulting in degradation of the target RNA by the RISC endonuclease complex or in the synthesis of additional RNA by RNA-dependent RNA polymerase (RdRP), which can activate DICER and result in additional siNA molecules, thereby amplifying the RNAi response.

Figure 18A-F shows non-limiting examples of chemically-modified siNA constructs of the present invention. In the figure, N stands for any nucleotide (adenosine, guanosine, cytosine, uridine, or optionally thymidine, for example thymidine can be substituted in the overhanging regions designated by parenthesis (N N)). Various modifications are shown for the sense and antisense strands of the siNA constructs.

Figure 18A: The sense strand comprises 21 nucleotides wherein the two terminal 3'-nucleotides are optionally base paired and wherein all nucleotides present are ribonucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all nucleotides present are ribonucleotides except for (N N)

nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. A modified internucleotide linkage, such as a phosphorothioate, phosphorodithioate or other modified internucleotide linkage as described herein, shown as “s” connects the (N N) nucleotides in the antisense strand.

5 **Figure 18B:** The sense strand comprises 21 nucleotides wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-O-methyl modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical
10 modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-O-methyl modified nucleotides
15 except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. A modified internucleotide linkage, such as a phosphorothioate, phosphorodithioate or other modified internucleotide linkage as described herein, shown as “s” connects the (N N) nucleotides in the sense and antisense strand.

20 **Figure 18C:** The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical
25 modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical
30 modifications described herein. A modified internucleotide linkage, such as a phosphorothioate, phosphorodithioate or other modified internucleotide linkage as described herein, shown as “s” connects the (N N) nucleotides in the antisense strand.

Figure 18D: The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein and wherein all purine nucleotides that may be present are 2'-deoxy nucleotides. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-O-methyl modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. A modified internucleotide linkage, such as a phosphorothioate, phosphorodithioate or other modified internucleotide linkage as described herein, shown as "s" connects the (N N) nucleotides in the antisense strand.

Figure 18E: The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-O-methyl modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. A modified internucleotide linkage, such as a phosphorothioate, phosphorodithioate or other modified internucleotide linkage as described herein, shown as "s" connects the (N N) nucleotides in the antisense strand.

Figure 18F: The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified

nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein and wherein all purine nucleotides that may be present are 2'-deoxy nucleotides. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-deoxy nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. A modified internucleotide linkage, such as a phosphorothioate, phosphorodithioate or other modified internucleotide linkage as described herein, shown as "s" connects the (N N) nucleotides in the antisense strand. The antisense strand of constructs A-F comprise sequence complementary to any target nucleic acid sequence of the invention. Furthermore, when a glyceryl moiety (L) is present at the 3'-end of the antisense strand for any construct shown in Figure 4 A-F, the modified internucleotide linkage is optional.

Figure 19 shows non-limiting examples of specific chemically modified siNA sequences of the invention. A-F applies the chemical modifications described in **Figure 18A-F** to a representative siNA sequence targeting the hepatitis C virus (HCV).

Figure 20 shows non-limiting examples of different siNA constructs of the invention. The examples shown (constructs 1, 2, and 3) have 19 representative base pairs; however, different embodiments of the invention include any number of base pairs described herein. Bracketed regions represent nucleotide overhangs, for example comprising about 1, 2, 3, or 4 nucleotides in length when present, preferably about 2 nucleotides. Such overhangs can be present or absent (i.e., blunt ends). Such blunt ends can be present on one end or both ends of the siNA molecule, for example where all nucleotides present in a siNA duplex are base paired. Constructs 1 and 2 can be used independently for RNAi activity. Construct 2 can comprise a polynucleotide or non-nucleotide linker, which can optionally be designed as a biodegradable linker. In one embodiment, the loop structure shown in construct 2 can comprise a biodegradable linker that results in the formation of construct 1 *in vivo* and/or *in vitro*. In another example, construct 3 can be used to generate construct 2 under the same principle wherein a linker is used to generate the active siNA construct 2 *in vivo* and/or *in vitro*, which can

optionally utilize another biodegradable linker to generate the active siNA construct 1 *in vivo* and/or *in vitro*. As such, the stability and/or activity of the siNA constructs can be modulated based on the design of the siNA construct for use *in vivo* or *in vitro* and/or *in vitro*.

5 **Figure 21** is a diagrammatic representation of a method used to determine target sites for siNA mediated RNAi within a particular target nucleic acid sequence, such as messenger RNA. (A) A pool of siNA oligonucleotides are synthesized wherein the antisense region of the siNA constructs has complementarity to target sites across the target nucleic acid sequence, and wherein the sense region comprises sequence
10 complementary to the antisense region of the siNA. (B) The sequences are transfected into cells. (C) Cells are selected based on phenotypic change that is associated with modulation of the target nucleic acid sequence. (D) The siNA is isolated from the selected cells and is sequenced to identify efficacious target sites within the target nucleic acid sequence.

15 **Figure 22** shows non-limiting examples of different stabilization chemistries (1-10) that can be used, for example, to stabilize the 3'-end of siNA sequences of the invention, including (1) [3'-3']-inverted deoxyribose; (2) deoxyribonucleotide; (3) [5'-3']-3'-deoxyribonucleotide; (4) [5'-3']-ribonucleotide; (5) [5'-3']-3'-O-methyl ribonucleotide; (6) 3'-glyceryl; (7) [3'-5']-3'-deoxyribonucleotide; (8) [3'-3']-deoxyribonucleotide; (9) [5'-2']-
20 deoxyribonucleotide; and (10) [5'-3']-dideoxyribonucleotide. In addition to modified and unmodified backbone chemistries indicated in the figure, these chemistries can be combined with different backbone modifications as described herein, for example, backbone modifications having Formula I. In addition, the 2'-deoxy nucleotide shown 5' to the terminal modifications shown can be another modified or unmodified nucleotide or
25 non-nucleotide described herein, for example modifications having any of Formulae I-VII or any combination thereof.

30 **Figure 23** shows a non-limiting example of siNA mediated inhibition of VEGF-induced angiogenesis using the rat corneal model of angiogenesis. siNA targeting site 2340 of VEGFR1 RNA (shown as Sirna/RPI No. 29695/29699) were compared to inverted controls (shown as Sirna/RPI No. 29983/29984) at three different concentrations and compared to a VEGF control in which no siNA was administered.

Figure 24 is a non-limiting example of a HBsAg screen of stabilized siNA constructs (“stab 4/5”, see **Table IV**) targeting HBV pregenomic RNA in HepG2 cells at 25 nM compared to untreated and matched chemistry inverted sequence controls. The siNA sense and antisense strands are shown by Sima/RPI number (sense/antisense).

5 **Figure 25** is a non-limiting example of a dose response HBsAg screen of stabilized siNA constructs (“stab 4/5”, see **Table IV**) targeting sites 262 and 1580 of the HBV pregenomic RNA in HepG2 cells at 0.5, 5, 10 and 25 nM compared to untreated and matched chemistry inverted sequence controls. The siNA sense and antisense strands are shown by Sima/RPI number (sense/antisense).

10 **Figure 26** shows a dose response comparison of two different stabilization chemistries (“stab 7/8” and “stab 7/11”, see **Table IV**) targeting site 1580 of the HBV pregenomic RNA in HepG2 cells at 5, 10, 25, 50 and 100 nM compared to untreated and matched chemistry inverted sequence controls. The siNA sense and antisense strands are shown by Sima/RPI number (sense/antisense).

15 **Figure 27** shows a non-limiting example of a strategy used to identify chemically modified siNA constructs of the invention that are nuclease resistance while preserving the ability to mediate RNAi activity. Chemical modifications are introduced into the siNA construct based on educated design parameters (e.g. introducing 2'-modifications, base modifications, backbone modifications, terminal cap modifications etc). The modified construct is tested in an appropriate system (e.g human serum for nuclease resistance, shown, or an animal model for PK/delivery parameters). In parallel, the siNA construct is tested for RNAi activity, for example in a cell culture system such as a luciferase reporter assay). Lead siNA constructs are then identified which possess a particular characteristic while maintaining RNAi activity, and can be further modified and
20 assayed once again. This same approach can be used to identify siNA-conjugate molecules with improved pharmacokinetic profiles, delivery, and RNAi activity.
25

30 **Figure 28** shows representative data of a chemically modified siNA construct (Stab 4/5, **Table IV**) targeting HBV site 1580 RNA compared to an unstabilized siRNA construct in a dose response time course HBsAg assay. The constructs were compared at different concentrations (5nM, 10 nM, 25 nM, 50 nM, and 100 nM) over the course of nine days. Activity based on HBsAg levels was determined at day 3, day 6, and day 9.

Figure 29 shows representative data of a chemically modified siNA construct (Stab 7/8, **Table IV**) targeting HBV site 1580 RNA compared to an unstabilized siRNA construct in a dose response time course HBsAg assay. The constructs were compared at different concentrations (5nM, 10 nM, 25 nM, 50 nM, and 100 nM) over the course of nine days. SiNA activity based on HBsAg levels was determined at day 3, day 6, and day 9.

Figure 30 shows representative data of a chemically modified siNA construct (Stab 7/11, **Table IV**) targeting HBV site 1580 RNA compared to an unstabilized siRNA construct in a dose response time course HBsAg assay. The constructs were compared at different concentrations (5nM, 10 nM, 25 nM, 50 nM, and 100 nM) over the course of nine days. SiNA activity based on HBsAg levels was determined at day 3, day 6, and day 9.

Figure 31 shows representative data of a chemically modified siNA construct (Stab 9/10, **Table IV**) targeting HBV site 1580 RNA compared to an unstabilized siRNA construct in a dose response time course HBsAg assay. The constructs were compared at different concentrations (5nM, 10 nM, 25 nM, 50 nM, and 100 nM) over the course of nine days. SiNA activity based on HBsAg levels was determined at day 3, day 6, and day 9.

Figure 32 shows non-limiting examples of inhibition of viral replication of a HCV/poliovirus chimera by siNA constructs targeted to HCV chimera (29579/29586; 29578/29585) compared to control (29593/29600).

Figure 33 shows a non-limiting example of a dose response study demonstrating the inhibition of viral replication of a HCV/poliovirus chimera by siNA construct (29579/29586) at various concentrations (1nM, 5nM, 10nM, and 25nM) compared to control (29593/29600).

Figure 34 shows a non-limiting example demonstrating the inhibition of viral replication of a HCV/poliovirus chimera by a chemically modified siRNA construct (30051/30053) compared to control construct (30052/30054).

Figure 35 shows a non-limiting example demonstrating the inhibition of viral replication of a HCV/poliovirus chimera by a chemically modified siRNA construct (30055/30057) compared to control construct (30056/30058).

Figure 36 shows a non-limiting example of several chemically modified siRNA constructs targeting viral replication of an HCV/poliovirus chimera at 10 nM treatment in comparison to a lipid control and an inverse siNA control construct 29593/ 29600.

Figure 37 shows a non-limiting example of several chemically modified siRNA constructs targeting viral replication of a HCV/poliovirus chimera at 25 nM treatment in comparison to a lipid control and an inverse siNA control construct 29593/ 29600.

Figure 38 shows a non-limiting example of several chemically modified siRNA constructs targeting viral replication of a Huh7 HCV replicon system at 25 nM treatment in comparison to untreated cells (“cells”), cells transfected with lipofectamine (“LFA2K”) and inverse siNA control constructs.

Figure 39 shows a non-limiting example of a dose response study using chemically modified siNA molecules (Stab 4/5, see **Table IV**) targeting HCV RNA sites 291, 300, and 303 in a Huh7 HCV replicon system at 5, 10, 25, and 100 nM treatment comparison to untreated cells (“cells”), cells transfected with lipofectamine (“LFA”) and inverse siNA control constructs.

Figure 40 shows a non-limiting example of several chemically modified siNA constructs (Stab 7/8, see **Table IV**) targeting viral replication in a Huh7 HCV replicon system at 25 nM treatment in comparison to untreated cells (“cells”), cells transfected with lipofectamine (“Lipid”) and inverse siNA control constructs.

Figure 41 shows a non-limiting example of a dose response study using chemically modified siNA molecules (Stab 7/8, see **Table IV**) targeting HCV site 327 in a Huh7 HCV replicon system at 5, 10, 25, 50, and 100 nM treatment in comparison to inverse siNA control constructs.

Figure 42 shows a synthetic scheme for post-synthetic modification of a nucleic acid molecule to produce a folate conjugate.

Figure 43 shows a synthetic scheme for generating an oligonucleotide or nucleic acid-folate conjugate.

Figure 44 shows an alternative synthetic scheme for generating an oligonucleotide or nucleic acid-folate conjugate.

5 **Figure 45** shows an alternative synthetic scheme for post-synthetic modification of a nucleic acid molecule to produce a folate conjugate.

Figure 46 shows a non-limiting example of a synthetic scheme for the synthesis of a N-acetyl-D-galactosamine-2'-aminouridine phosphoramidite conjugate of the invention.

10 **Figure 47** shows a non-limiting example of a synthetic scheme for the synthesis of a N-acetyl-D-galactosamine-D-threoninol phosphoramidite conjugate of the invention.

15 **Figure 48** shows a non-limiting example of a N-acetyl-D-galactosamine siNA nucleic acid conjugate of the invention. W shown in the example refers to a biodegradable linker, for example a nucleic acid dimer, trimer, or tetramer comprising ribonucleotides and/or deoxyribonucleotides. The siNA can be conjugated at the 3', 5' or both 3' and 5' ends of the sense strand of a double stranded siNA and/or the 3'-end of the antisense strand of the siNA. A single stranded siNA molecule can be conjugated at the 3'-end of the siNA.

Figure 49 shows a non-limiting example of a synthetic scheme for the synthesis of a dodecanoic acid derived conjugate linker of the invention.

20 **Figure 50** shows a non-limiting example of a synthetic scheme for the synthesis of an oxime linked nucleic acid/peptide conjugate of the invention.

25 **Figure 51** shows non-limiting examples of phospholipid derived siNA conjugates of the invention. CL shown in the examples refers to a biodegradable linker, for example a nucleic acid dimer, trimer, or tetramer comprising ribonucleotides and/or deoxyribonucleotides. The siNA can be conjugated at the 3', 5' or both 3' and 5' ends of the sense strand of a double stranded siNA and/or the 3'-end of the antisense strand of the siNA. A single stranded siNA molecule can be conjugated at the 3'-end of the siNA.

Figure 52 shows a non-limiting example of a synthetic scheme for preparing a phospholipid derived siNA conjugates of the invention.

30 **Figure 53** shows a non-limiting example of a synthetic scheme for preparing a poly-N-acetyl-D-galactosamine nucleic acid conjugate of the invention.

Figure 54 shows a non-limiting example of the synthesis of siNA cholesterol conjugates of the invention using a phosphoramidite approach.

Figure 55 shows a non-limiting example of the synthesis of siNA PEG conjugates of the invention using NHS ester coupling.

5 **Figure 56** shows a non-limiting example of the synthesis of siNA cholesterol conjugates of the invention using NHS ester coupling.

Figure 57 shows a non-limiting example of various siNA cholesterol conjugates of the invention.

10 **Figure 58** shows a non-limiting example of various siNA cholesterol conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a double stranded siNA molecule.

Figure 59 shows a non-limiting example of various siNA cholesterol conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a double stranded siNA molecule.

15 **Figure 60** shows a non-limiting example of various siNA cholesterol conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a single stranded siNA molecule.

20 **Figure 61** shows a non-limiting example of various siNA phospholipid conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a double stranded siNA molecule.

Figure 62 shows a non-limiting example of various siNA phospholipid conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a single stranded siNA molecule.

25 **Figure 63** shows a non-limiting example of various siNA galactosamine conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a double stranded siNA molecule.

Figure 64 shows a non-limiting example of various siNA galactosamine conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a single stranded siNA molecule.

Figure 65 shows a non-limiting example of various generalized siNA conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a double stranded siNA molecule. CONJ in the figure refers to any biologically active compound or any other conjugate compound as described
5 herein and in the Formulae herein.

Figure 66 shows a non-limiting example of various generalized siNA conjugates of the invention in which various linker chemistries and/or cleavable linkers can be utilized at different positions of a single stranded siNA molecule. CONJ in the figure refers to any biologically active compound or any other conjugate compound as described
10 herein and in the Formulae herein.

Figure 67 shows a non-limiting example of the pharmacokinetic distribution of intact siNA in liver after administration of conjugated or unconjugated siNA molecules in mice.

Figure 68 shows a non-limiting example of the activity of conjugated siNA
15 constructs compared to matched chemistry unconjugated siNA constructs in an HBV cell culture system without the use of transfection lipid. As shown in the Figure, siNA conjugates provide efficacy in cell culture without the need for transfection reagent.

Figure 69 shows a non-limiting example of a scheme for the synthesis of a mono-galactosamine phosphoramidite of the invention that can be used to generate
20 galactosamine conjugated nucleic acid molecules.

Figure 70 shows a non-limiting example of a scheme for the synthesis of a tri-galactosamine phosphoramidite of the invention that can be used to generate tri-galactosamine conjugated nucleic acid molecules.

Figure 71 shows a non-limiting example of a scheme for the synthesis of another
25 tri-galactosamine phosphoramidite of the invention that can be used to generate tri-galactosamine conjugated nucleic acid molecules.

Figure 72 shows a non-limiting example of an alternate scheme for the synthesis of a tri-galactosamine phosphoramidite of the invention that can be used to generate tri-galactosamine conjugated nucleic acid molecules.

Figure 73 shows a non-limiting example of a scheme for the synthesis of a cholesterol NHS ester of the invention that can be used to generate cholesterol conjugated
30 nucleic acid molecules.

Figure 74 shows non-limiting examples of phosphorylated siNA molecules of the invention, including linear and duplex constructs and asymmetric derivatives thereof.

Figure 75 shows non-limiting examples of a chemically modified terminal phosphate groups of the invention.

5 **Figure 76** shows a non-limiting example of inhibition of VEGF induced neovascularization in the rat corneal model. VEGFr1 site 349 active siNA having “Stab 9/10” chemistry (Sirna # 31270/31273) was tested for inhibition of VEGF-induced angiogenesis at three different concentrations (2.0 ug, 1.0 ug, and 0.1 μ g dose response) as compared to a matched chemistry inverted control siNA construct (Sirna #
10 31276/31279) at each concentration and a VEGF control in which no siNA was administered. As shown in the figure, the active siNA construct having “Stab 9/10” chemistry (Sirna # 31270/31273) is highly effective in inhibiting VEGF-induced angiogenesis in the rat corneal model compared to the matched chemistry inverted control siNA at concentrations from 0.1 μ g to 2.0 ug.

15 **Figure 77** shows activity of modified siNA constructs having stab 4/5 (Sirna 30355/30366), stab 7/8 (Sirna 30612/30620), and stab 7/11 (Sirna 30612/31175) chemistries and an all ribo siNA construct (Sirna 30287/30298) in the reduction of HBsAg levels compared to matched inverted controls at **A.** 3 days, **B.** 9 days, and **C.** 21 days post transfection. Also shown is the corresponding percent inhibition as function of
20 time at siNA concentrations of **D.** 100 nM, **E.** 50 nM, and **F.** 25 nM.

Figure 78 shows non-limiting examples of phosphorylated siNA molecules of the invention, including linear and duplex constructs and asymmetric derivatives thereof.

Figure 79 shows non-limiting examples of chemically modified terminal phosphate groups of the invention.

25 **Figure 80** shows a non-limiting example of reduction of serum HBV DNA in mice treated with hydrodynamically administered chemically modified siNA (Stab 7/8 and Stab 9/10) targeting HBV RNA compared to matched chemistry inverted controls and a saline control.

30 **Figure 81** shows a non-limiting example of reduction of serum HBV S antigen (HBsAg) in mice treated with hydrodynamically administered chemically modified siNA (Stab 7/8 and Stab 9/10) targeting HBV RNA compared to matched chemistry inverted controls and a saline control.

Figure 82 shows a non-limiting example of reduction of serum HBV RNA in mice treated with hydrodynamically administered chemically modified siNA (Stab 7/8 and Stab 9/10) targeting HBV RNA compared to matched chemistry inverted controls and a saline control.

5 **Figure 83** shows a non-limiting example of reduction of serum HBV DNA in mice treated with hydrodynamically administered chemically modified siNA (Stab 7/8 and Stab 9/10) targeting HBV RNA at 5 days and 7 days post administration.

10 **Figure 84** shows a non-limiting example of an assay for dose dependent reduction of Luciferase expression utilizing Stab 7/8 chemically modified siNA constructs targeting luciferase RNA sites 80, 237, and 1478 that were selected from a screen using all Stab 7/8 chemically modified siNA constructs.

15 **Figure 85** shows a non-limiting example of an assay for dose dependent reduction of Luciferase expression utilizing Stab 7/8 chemically modified siNA constructs targeting luciferase RNA sites 1544 and 1607 that were selected from a screen using all Stab 7/8 chemically modified siNA constructs.

20 **Figure 86** shows a non-limiting example of an assay screen of Stab 7/8 siNA constructs targeting various sites of HCV RNA in a replicon system compared to untreated, lipid, and an inverted control. As shown in the figure, several Stab 7/8 constructs were identified with potent anti-HCV activity as shown by reduction in HCV RNA levels.

Figure 87 shows a non-limiting example of an assay screen of Stab 7/8 siNA constructs targeting various sites of HBV RNA in HEpG2 cells compared to untreated cells and an inverted control. As shown in the figure, several Stab 7/8 constructs were identified with potent anti-HBV activity as shown by reduction in HBV S antigen levels.

25 **Figure 88** shows a non-limiting example of an assay screen of various combinations of chemically modified siNA constructs (e.g., Stab 7/8, 7/10, 7/11, 9/8, and 9/10) targeting site 1580 of HBV RNA in HEpG2 cells compared to untreated cells and an matched chemistry inverted controls. As shown in the figure, the combination

chemistries tested demonstrated potent anti-HBV activity as shown by reduction in HBV S antigen levels.

Figure 89 shows a non-limiting example of an assay screen of various combinations of chemically modified siNA constructs (e.g., Stab 7/8, 9/10, 6/10, 16/8, 16/10, 18/8, and 18/10) targeting site 1580 of HBV RNA in HEpG2 cells compared to untreated cells and an matched chemistry inverted controls. As shown in the figure, the combination chemistries tested demonstrated potent anti-HBV activity as shown by reduction in HBV S antigen levels.

Figure 90 shows a non-limiting example of an assay screen of various combinations of chemically modified siNA constructs (e.g., Stab 4/8, 4/10, 7/5, 7/10, 9/5, 9/8, and 9/11) targeting site 1580 of HBV RNA in HEpG2 cells compared to untreated cells and an matched chemistry inverted controls. As shown in the figure, the combination chemistries tested demonstrated potent anti-HBV activity as shown by reduction in HBV S antigen levels.

15

DETAILED DESCRIPTION OF THE INVENTION

Mechanism of action of Nucleic Acid Molecules of the Invention

The discussion that follows discusses the proposed mechanism of RNA interference mediated by short interfering RNA as is presently known, and is not meant to be limiting and is not an admission of prior art. Applicant demonstrates herein that chemically-modified short interfering nucleic acids possess similar or improved capacity to mediate RNAi as do siRNA molecules and are expected to possess improved stability and activity *in vivo*; therefore, this discussion is not meant to be limited to siRNA only and can be applied to siNA as a whole. By "improved capacity to mediate RNAi" or "improved RNAi activity" is meant to include RNAi activity measured *in vitro* and/or *in vivo* where the RNAi activity is a reflection of both the ability of the siNA to mediate RNAi and the stability of the siNAs of the invention. In this invention, the product of these activities can be increased *in vitro* and/or *in vivo* compared to an all RNA siRNA or a siNA containing a plurality of ribonucleotides. In some cases, the activity or stability of the

siNA molecule can be decreased (i.e., less than ten-fold), but the overall activity of the siNA molecule is enhanced *in vitro* and/or *in vivo*.

RNA interference refers to the process of sequence specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Zamore *et al.*, 2000, *Cell*, 101, 25-33; Fire *et al.*, 1998, *Nature*, 391, 806). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-conserved cellular defense mechanism used to prevent the expression of foreign genes which is commonly shared by diverse flora and phyla (Fire *et al.*, 1999, *Trends Genet.*, 15, 358). Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or the random integration of transposon elements into a host genome via a cellular response that specifically destroys homologous single-stranded RNA or viral genomic RNA. The presence of dsRNA in cells triggers the RNAi response though a mechanism that has yet to be fully characterized. This mechanism appears to be different from the interferon response that results from dsRNA-mediated activation of protein kinase PKR and 2', 5'-oligoadenylate synthetase resulting in non-specific cleavage of mRNA by ribonuclease L.

The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as Dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Berstein *et al.*, 2001, *Nature*, 409, 363). Short interfering RNAs derived from Dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes. Dicer has also been implicated in the excision of 21- and 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner *et al.*, 2001, *Science*, 293, 834). The RNAi response also features an endonuclease complex containing a siRNA, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having sequence homologous to the siRNA. Cleavage of the target RNA takes place in the middle of the region complementary to the guide sequence of the siRNA duplex (Elbashir *et al.*, 2001, *Genes Dev.*, 15, 188). In addition, RNA interference can also involve small RNA (e.g., micro-RNA or miRNA) mediated gene silencing, presumably through cellular

mechanisms that regulate chromatin structure and thereby prevent transcription of target gene sequences (see for example Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 2002, *Science*, 297, 2232-2237). As such, siNA molecules of the invention can be used to
 5 mediate gene silencing via interaction with RNA transcripts or alternately by interaction with particular gene sequences, wherein such interaction results in gene silencing either at the transcriptional level or post-transcriptional level.

RNAi has been studied in a variety of systems. Fire *et al.*, 1998, *Nature*, 391, 806, were the first to observe RNAi in *C. elegans*. Wianny and Goetz, 1999, *Nature Cell*
 10 *Biol.*, 2, 70, describe RNAi mediated by dsRNA in mouse embryos. Hammond *et al.*, 2000, *Nature*, 404, 293, describe RNAi in *Drosophila* cells transfected with dsRNA. Elbashir *et al.*, 2001, *Nature*, 411, 494, describe RNAi induced by introduction of duplexes of synthetic 21-nucleotide RNAs in cultured mammalian cells including human embryonic kidney and HeLa cells. Recent work in *Drosophila* embryonic lysates has
 15 revealed certain requirements for siRNA length, structure, chemical composition, and sequence that are essential to mediate efficient RNAi activity. These studies have shown that 21 nucleotide siRNA duplexes are most active when containing two 2-nucleotide 3'-terminal nucleotide overhangs. Furthermore, substitution of one or both siRNA strands with 2'-deoxy or 2'-O-methyl nucleotides abolishes RNAi activity, whereas substitution of
 20 3'-terminal siRNA nucleotides with deoxy nucleotides was shown to be tolerated. Mismatch sequences in the center of the siRNA duplex were also shown to abolish RNAi activity. In addition, these studies also indicate that the position of the cleavage site in the target RNA is defined by the 5'-end of the siRNA guide sequence rather than the 3'-end (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). Other studies have indicated that a 5'-
 25 phosphate on the target-complementary strand of a siRNA duplex is required for siRNA activity and that ATP is utilized to maintain the 5'-phosphate moiety on the siRNA (Nykanen *et al.*, 2001, *Cell*, 107, 309); however, siRNA molecules lacking a 5'-phosphate are active when introduced exogenously, suggesting that 5'-phosphorylation of siRNA constructs may occur *in vivo*.

30 Synthesis of Nucleic acid Molecules

Synthesis of nucleic acids greater than 100 nucleotides in length is difficult using automated methods, and the therapeutic cost of such molecules is prohibitive. In this invention, small nucleic acid motifs ("small" refers to nucleic acid motifs no more than 100 nucleotides in length, preferably no more than 80 nucleotides in length, and most preferably no more than 50 nucleotides in length; *e.g.*, individual siNA oligonucleotide sequences or siNA sequences synthesized in tandem) are preferably used for exogenous delivery. The simple structure of these molecules increases the ability of the nucleic acid to invade targeted regions of protein and/or RNA structure. Exemplary molecules of the instant invention are chemically synthesized, and others can similarly be synthesized.

Oligonucleotides (*e.g.*, certain modified oligonucleotides or portions of oligonucleotides lacking ribonucleotides) are synthesized using protocols known in the art, for example as described in Caruthers *et al.*, 1992, *Methods in Enzymology* 211, 3-19, Thompson *et al.*, International PCT Publication No. WO 99/54459, Wincott *et al.*, 1995, *Nucleic Acids Res.* 23, 2677-2684, Wincott *et al.*, 1997, *Methods Mol. Bio.*, 74, 59, Brennan *et al.*, 1998, *Biotechnol Bioeng.*, 61, 33-45, and Brennan, U.S. Pat. No. 6,001,311. All of these references are incorporated herein by reference. The synthesis of oligonucleotides makes use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. In a non-limiting example, small scale syntheses are conducted on a 394 Applied Biosystems, Inc. synthesizer using a 0.2 μ mol scale protocol with a 2.5 min coupling step for 2'-O-methylated nucleotides and a 45 second coupling step for 2'-deoxy nucleotides or 2'-deoxy-2'-fluoro nucleotides. **Table V** outlines the amounts and the contact times of the reagents used in the synthesis cycle. Alternatively, syntheses at the 0.2 μ mol scale can be performed on a 96-well plate synthesizer, such as the instrument produced by Protogene (Palo Alto, CA) with minimal modification to the cycle. A 33-fold excess (60 μ L of 0.11 M = 6.6 μ mol) of 2'-O-methyl phosphoramidite and a 105-fold excess of S-ethyl tetrazole (60 μ L of 0.25 M = 15 μ mol) can be used in each coupling cycle of 2'-O-methyl residues relative to polymer-bound 5'-hydroxyl. A 22-fold excess (40 μ L of 0.11 M = 4.4 μ mol) of deoxy phosphoramidite and a 70-fold excess of S-ethyl tetrazole (40 μ L of 0.25 M = 10 μ mol) can be used in each coupling cycle of deoxy residues relative to polymer-bound 5'-hydroxyl. Average coupling yields on the 394 Applied Biosystems, Inc. synthesizer, determined by colorimetric quantitation of the trityl fractions, are typically 97.5-99%.

Other oligonucleotide synthesis reagents for the 394 Applied Biosystems, Inc. synthesizer include the following: detritylation solution is 3% TCA in methylene chloride (ABI); capping is performed with 16% *N*-methyl imidazole in THF (ABI) and 10% acetic anhydride/10% 2,6-lutidine in THF (ABI); and oxidation solution is 16.9 mM I₂, 49 mM
 5 pyridine, 9% water in THF (PERSEPTIVE™). Burdick & Jackson Synthesis Grade acetonitrile is used directly from the reagent bottle. S-Ethyltetrazole solution (0.25 M in acetonitrile) is made up from the solid obtained from American International Chemical, Inc. Alternately, for the introduction of phosphorothioate linkages, Beaucage reagent (3H-1,2-Benzodithiol-3-one 1,1-dioxide, 0.05 M in acetonitrile) is used.

10 Deprotection of the DNA-based oligonucleotides is performed as follows: the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution of 40% aqueous methylamine (1 mL) at 65 °C for 10 minutes. After cooling to -20 °C, the supernatant is removed from the polymer support. The support is washed three times with 1.0 mL of EtOH:MeCN:H₂O/3:1:1, vortexed and
 15 the supernatant is then added to the first supernatant. The combined supernatants, containing the oligoribonucleotide, are dried to a white powder.

The method of synthesis used for RNA including certain siNA molecules of the invention follows the procedure as described in Usman *et al.*, 1987, *J. Am. Chem. Soc.*, 109, 7845; Scaringe *et al.*, 1990, *Nucleic Acids Res.*, 18, 5433; and Wincott *et al.*, 1995,
 20 *Nucleic Acids Res.* 23, 2677-2684 Wincott *et al.*, 1997, *Methods Mol. Bio.*, 74, 59, and makes use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. In a non-limiting example, small scale syntheses are conducted on a 394 Applied Biosystems, Inc. synthesizer using a 0.2 μmol scale protocol with a 7.5 min coupling step for alkylsilyl
 25 protected nucleotides and a 2.5 min coupling step for 2'-O-methylated nucleotides. **Table V** outlines the amounts and the contact times of the reagents used in the synthesis cycle. Alternatively, syntheses at the 0.2 μmol scale can be done on a 96-well plate synthesizer, such as the instrument produced by Protogene (Palo Alto, CA) with minimal modification to the cycle. A 33-fold excess (60 μL of 0.11 M = 6.6 μmol) of 2'-O-methyl
 30 phosphoramidite and a 75-fold excess of S-ethyl tetrazole (60 μL of 0.25 M = 15 μmol) can be used in each coupling cycle of 2'-O-methyl residues relative to polymer-bound 5'-hydroxyl. A 66-fold excess (120 μL of 0.11 M = 13.2 μmol) of alkylsilyl (ribo) protected

phosphoramidite and a 150-fold excess of S-ethyl tetrazole (120 μ L of 0.25 M = 30 μ mol) can be used in each coupling cycle of ribo residues relative to polymer-bound 5'-hydroxyl. Average coupling yields on the 394 Applied Biosystems, Inc. synthesizer, determined by colorimetric quantitation of the trityl fractions, are typically 97.5-99%.

5 Other oligonucleotide synthesis reagents for the 394 Applied Biosystems, Inc. synthesizer include the following: detritylation solution is 3% TCA in methylene chloride (ABI); capping is performed with 16% *N*-methyl imidazole in THF (ABI) and 10% acetic anhydride/10% 2,6-lutidine in THF (ABI); oxidation solution is 16.9 mM I₂, 49 mM pyridine, 9% water in THF (PERSEPTIVE™). Burdick & Jackson Synthesis Grade
10 acetonitrile is used directly from the reagent bottle. S-Ethyltetrazole solution (0.25 M in acetonitrile) is made up from the solid obtained from American International Chemical, Inc. Alternately, for the introduction of phosphorothioate linkages, Beaucage reagent (3H-1,2-Benzodithiol-3-one 1,1-dioxide 0.05 M in acetonitrile) is used.

Deprotection of the RNA is performed using either a two-pot or one-pot protocol.
15 For the two-pot protocol, the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution of 40% aq. methylamine (1 mL) at 65 °C for 10 minutes. After cooling to -20 °C, the supernatant is removed from the polymer support. The support is washed three times with 1.0 mL of EtOH:MeCN:H₂O/3:1:1, vortexed and the supernatant is then added to the first
20 supernatant. The combined supernatants, containing the oligoribonucleotide, are dried to a white powder. The base deprotected oligoribonucleotide is resuspended in anhydrous TEA/HF/NMP solution (300 μ L of a solution of 1.5 mL *N*-methylpyrrolidinone, 750 μ L TEA and 1 mL TEA•3HF to provide a 1.4 M HF concentration) and heated to 65 °C. After 1.5 h, the oligomer is quenched with 1.5 M NH₄HCO₃.

25 Alternatively, for the one-pot protocol, the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution of 33% ethanolic methylamine/DMSO: 1/1 (0.8 mL) at 65 °C for 15 minutes. The vial is brought to room temperature TEA•3HF (0.1 mL) is added and the vial is heated at 65 °C for 15 minutes. The sample is cooled at -20 °C and then quenched with
30 1.5 M NH₄HCO₃.

For purification of the trityl-on oligomers, the quenched NH_4HCO_3 solution is loaded onto a C-18 containing cartridge that had been prewashed with acetonitrile followed by 50 mM TEAA. After washing the loaded cartridge with water, the RNA is detritylated with 0.5% TFA for 13 minutes. The cartridge is then washed again with
5 water, salt exchanged with 1 M NaCl and washed with water again. The oligonucleotide is then eluted with 30% acetonitrile.

The average stepwise coupling yields are typically >98% (Wincott *et al.*, 1995 *Nucleic Acids Res.* 23, 2677-2684). Those of ordinary skill in the art will recognize that the scale of synthesis can be adapted to be larger or smaller than the example described
10 above including but not limited to 96-well format.

Alternatively, the nucleic acid molecules of the present invention can be synthesized separately and joined together post-synthetically, for example, by ligation (Moore *et al.*, 1992, *Science* 256, 9923; Draper *et al.*, International PCT publication No. WO 93/23569; Shabarova *et al.*, 1991, *Nucleic Acids Research* 19, 4247; Bellon *et al.*,
15 1997, *Nucleosides & Nucleotides*, 16, 951; Bellon *et al.*, 1997, *Bioconjugate Chem.* 8, 204), or by hybridization following synthesis and/or deprotection.

The siNA molecules of the invention can also be synthesized via a tandem synthesis methodology as described in Example 1 herein, wherein both siNA strands are synthesized as a single contiguous oligonucleotide fragment or strand separated by a
20 cleavable linker which is subsequently cleaved to provide separate siNA fragments or strands that hybridize and permit purification of the siNA duplex. The linker can be a polynucleotide linker or a non-nucleotide linker. The tandem synthesis of siNA as described herein can be readily adapted to both multiwell/multiplate synthesis platforms such as 96 well or similarly larger multi-well platforms. The tandem synthesis of siNA as
25 described herein can also be readily adapted to large scale synthesis platforms employing batch reactors, synthesis columns and the like.

A siNA molecule can also be assembled from two distinct nucleic acid strands or fragments wherein one fragment includes the sense region and the second fragment includes the antisense region of the RNA molecule.

The nucleic acid molecules of the present invention can be modified extensively to enhance stability by modification with nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-H (for a review see Usman and Cedergren, 1992, *TIBS* 17, 34; Usman *et al.*, 1994, *Nucleic Acids Symp. Ser.* 31, 163). siNA constructs can
 5 be purified by gel electrophoresis using general methods or can be purified by high pressure liquid chromatography (HPLC; see Wincott *et al.*, *supra*, the totality of which is hereby incorporated herein by reference) and re-suspended in water.

In another aspect of the invention, siNA molecules of the invention are expressed from transcription units inserted into DNA or RNA vectors. The recombinant vectors can
 10 be DNA plasmids or viral vectors. siNA expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus, retrovirus, adenovirus, or alphavirus. The recombinant vectors capable of expressing the siNA molecules can be delivered as described herein, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of siNA molecules.

15 Optimizing Activity of the nucleic acid molecule of the invention.

Chemically synthesizing nucleic acid molecules with modifications (base, sugar and/or phosphate) can prevent their degradation by serum ribonucleases, which can increase their potency (see *e.g.*, Eckstein *et al.*, International Publication No. WO 92/07065; Perrault *et al.*, 1990 *Nature* 344, 565; Pieken *et al.*, 1991, *Science* 253, 314;
 20 Usman and Cedergren, 1992, *Trends in Biochem. Sci.* 17, 334; Usman *et al.*, International Publication No. WO 93/15187; and Rossi *et al.*, International Publication No. WO 91/03162; Sproat, U.S. Pat. No. 5,334,711; Gold *et al.*, U.S. Pat. No. 6,300,074; and Burgin *et al.*, *supra*; all of which are incorporated by reference herein). All of the above references describe various chemical modifications that can be made to the base,
 25 phosphate and/or sugar moieties of the nucleic acid molecules described herein. Modifications that enhance their efficacy in cells, and removal of bases from nucleic acid molecules to shorten oligonucleotide synthesis times and reduce chemical requirements are desired.

There are several examples in the art describing sugar, base and phosphate
 30 modifications that can be introduced into nucleic acid molecules with significant enhancement in their nuclease stability and efficacy. For example, oligonucleotides are

modified to enhance stability and/or enhance biological activity by modification with nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-O-allyl, 2'-H, nucleotide base modifications (for a review see Usman and Cedergren, 1992, *TIBS*, 17, 34; Usman *et al.*, 1994, *Nucleic Acids Symp. Ser.* 31, 163; Burgin *et al.*, 1996, *Biochemistry*, 35, 14090). Sugar modification of nucleic acid molecules have been extensively described in the art (see Eckstein *et al.*, *International Publication* PCT No. WO 92/07065; Perrault *et al.* *Nature*, 1990, 344, 565-568; Pieken *et al.* *Science*, 1991, 253, 314-317; Usman and Cedergren, *Trends in Biochem. Sci.*, 1992, 17, 334-339; Usman *et al.* *International Publication* PCT No. WO 93/15187; Sproat, *U.S. Pat.* No. 5,334,711 and Beigelman *et al.*, 1995, *J. Biol. Chem.*, 270, 25702; Beigelman *et al.*, *International PCT publication* No. WO 97/26270; Beigelman *et al.*, *U.S. Pat.* No. 5,716,824; Usman *et al.*, *U.S. Pat.* No. 5,627,053; Woolf *et al.*, *International PCT Publication* No. WO 98/13526; Thompson *et al.*, *USSN* 60/082,404 which was filed on April 20, 1998; Karpeisky *et al.*, 1998, *Tetrahedron Lett.*, 39, 1131; Earnshaw and Gait, 1998, *Biopolymers (Nucleic Acid Sciences)*, 48, 39-55; Verma and Eckstein, 1998, *Annu. Rev. Biochem.*, 67, 99-134; and Burlina *et al.*, 1997, *Bioorg. Med. Chem.*, 5, 1999-2010; all of the references are hereby incorporated in their totality by reference herein). Such publications describe general methods and strategies to determine the location of incorporation of sugar, base and/or phosphate modifications and the like into nucleic acid molecules without modulating catalysis, and are incorporated by reference herein. In view of such teachings, similar modifications can be used as described herein to modify the siNA nucleic acid molecules of the instant invention so long as the ability of siNA to promote RNAi in cells is not significantly inhibited.

While chemical modification of oligonucleotide internucleotide linkages with phosphorothioate, phosphorodithioate, and/or 5'-methylphosphonate linkages improves stability, excessive modifications can cause some toxicity or decreased activity. Therefore, when designing nucleic acid molecules, the amount of these internucleotide linkages should be minimized. The reduction in the concentration of these linkages should lower toxicity, resulting in increased efficacy and higher specificity of these molecules.

Short interfering nucleic acid (siNA) molecules having chemical modifications that maintain or enhance activity are provided. Such a nucleic acid is also generally more


resistant to nucleases than an unmodified nucleic acid. Accordingly, the *in vitro* and/or *in vivo* activity should not be significantly lowered. In cases in which modulation is the goal, therapeutic nucleic acid molecules delivered exogenously should optimally be stable within cells until translation of the target RNA has been modulated long enough to reduce the levels of the undesirable protein. This period of time varies between hours to days depending upon the disease state. Improvements in the chemical synthesis of RNA and DNA (Wincott *et al.*, 1995, *Nucleic Acids Res.* 23, 2677; Caruthers *et al.*, 1992, *Methods in Enzymology* 211,3-19 (incorporated by reference herein)) have expanded the ability to modify nucleic acid molecules by introducing nucleotide modifications to enhance their nuclease stability, as described above.

In one embodiment, nucleic acid molecules of the invention include one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) G-clamp nucleotides. A G-clamp nucleotide is a modified cytosine analog wherein the modifications confer the ability to hydrogen bond both Watson-Crick and Hoogsteen faces of a complementary guanine within a duplex, see for example Lin and Matteucci, 1998, *J. Am. Chem. Soc.*, 120, 8531-8532. A single G-clamp analog substitution within an oligonucleotide can result in substantially enhanced helical thermal stability and mismatch discrimination when hybridized to complementary oligonucleotides. The inclusion of such nucleotides in nucleic acid molecules of the invention results in both enhanced affinity and specificity to nucleic acid targets, complementary sequences, or template strands. In another embodiment, nucleic acid molecules of the invention include one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) LNA "locked nucleic acid" nucleotides such as a 2', 4'-C methylene bicyclo nucleotide (see for example Wengel *et al.*, International PCT Publication No. WO 00/66604 and WO 99/14226).

In another embodiment, the invention features conjugates and/or complexes of siNA molecules of the invention. Such conjugates and/or complexes can be used to facilitate delivery of siNA molecules into a biological system, such as a cell. The conjugates and complexes provided by the instant invention can impart therapeutic activity by transferring therapeutic compounds across cellular membranes, altering the pharmacokinetics, and/or modulating the localization of nucleic acid molecules of the invention. The present invention encompasses the design and synthesis of novel conjugates and complexes for the delivery of molecules, including, but not limited to,


small molecules, lipids, cholesterol, phospholipids, nucleosides, nucleotides, nucleic acids, antibodies, toxins, negatively charged polymers and other polymers, for example, proteins, peptides, hormones, carbohydrates, polyethylene glycols, or polyamines, across cellular membranes. In general, the transporters described are designed to be used either
 5 individually or as part of a multi-component system, with or without degradable linkers. These compounds are expected to improve delivery and/or localization of nucleic acid molecules of the invention into a number of cell types originating from different tissues, in the presence or absence of serum (see Sullenger and Cech, U.S. Pat. No. 5,854,038). Conjugates of the molecules described herein can be attached to biologically active
 10 molecules via linkers that are biodegradable, such as biodegradable nucleic acid linker molecules.

In one embodiment, the invention features a compound having Formula 1:



15 wherein each R₁, R₃, R₄, R₅, R₆, R₇ and R₈ is independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, or a protecting group, each “n” is independently an integer from 0 to about 200, R₁₂ is a straight or branched chain alkyl, substituted alkyl, aryl, or substituted aryl, and R₂ is a siNA molecule or a portion thereof.


In one embodiment, the invention features a compound having Formula 2:



20 wherein each R₃, R₄, R₅, R₆ and R₇ is independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, or a protecting group, each “n” is independently an integer

from 0 to about 200, R_{12} is a straight or branched chain alkyl, substituted alkyl, aryl, or substituted aryl, and R_2 is a siNA molecule or a portion thereof.

In one embodiment, the invention features a compound having Formula 3:




5

3

wherein each R_1 , R_3 , R_4 , R_5 , R_6 and R_7 is independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, or a protecting group, each “n” is independently an integer from 0 to about 200, R_{12} is a straight or branched chain alkyl, substituted alkyl, aryl, or substituted aryl, and R_2 is a siNA molecule or a portion thereof.

10


In one embodiment, the invention features a compound having Formula 4:



4

wherein each R_3 , R_4 , R_5 , R_6 and R_7 is independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, or a protecting group, each “n” is independently an integer from 0 to about 200, R_2 is a siNA molecule or a portion thereof, and R_{13} is an amino acid side chain.


In one embodiment, the invention features a compound having Formula 5:



5

wherein each R_1 and R_4 is independently a protecting group or hydrogen, each R_3 , R_5 , R_6 , R_7 and R_8 is independently hydrogen, alkyl or nitrogen protecting group, each
 5 “n” is independently an integer from 0 to about 200, R_{12} is a straight or branched chain alkyl, substituted alkyl, aryl, or substituted aryl, and each R_9 and R_{10} is independently a nitrogen containing group, cyanoalkoxy, alkoxy, aryloxy, or alkyl group.


In one embodiment, the invention features a compound having Formula 6:



6

wherein each R_4 , R_5 , R_6 and R_7 is independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, or a protecting group, R_2 is a siNA molecule or a portion thereof, each “n” is independently an integer from 0 to about 200, and L is a degradable linker.


15 In one embodiment, the invention features a compound having Formula 7:



7

wherein each R_1 , R_3 , R_4 , R_5 , R_6 and R_7 is independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, or a protecting group, each “n” is independently an integer from 0 to about 200, R_{12} is a straight or branched chain alkyl, substituted alkyl, aryl, or substituted aryl, and R_2 is a siNA molecule or a portion thereof.

5 In one embodiment, the invention features a compound having Formula 8:



8

wherein each R_1 and R_4 is independently a protecting group or hydrogen, each R_3 , R_5 , R_6 and R_7 is independently hydrogen, alkyl or nitrogen protecting group, each “n” is independently an integer from 0 to about 200, R_{12} is a straight or branched chain alkyl, substituted alkyl, aryl, or substituted aryl, and each R_9 and R_{10} is independently a nitrogen containing group, cyanoalkoxy, alkoxy, aryloxy, or alkyl group.

In one embodiment, R_{13} of a compound of the invention comprises an alkylamino or an alkoxy group, for example, $-\text{CH}_2\text{O}-$ or $-\text{CH}(\text{CH}_2)\text{CH}_2\text{O}-$.

15 In another embodiment, R_{12} of a compound of the invention is an alkylhydroxyl, for example, $-(\text{CH}_2)_n\text{OH}$, where n comprises an integer from about 1 to about 10.

In another embodiment, L of Formula 6 of the invention comprises serine, threonine, or a photolabile linkage.

20 In one embodiment, R_9 of a compound of the invention comprises a phosphorus protecting group, for example $-\text{OCH}_2\text{CH}_2\text{CN}$ (oxyethylcyano).

In one embodiment, R_{10} of a compound of the invention comprises a nitrogen containing group, for example, $-\text{N}(\text{R}_{14})$ wherein R_{14} is a straight or branched chain alkyl having from about 1 to about 10 carbons.

In another embodiment, R₁₀ of a compound of the invention comprises a heterocycloalkyl or heterocycloalkenyl ring containing from about 4 to about 7 atoms, and having from about 1 to about 3 heteroatoms comprising oxygen, nitrogen, or sulfur.

In another embodiment, R₁ of a compound of the invention comprises an acid labile protecting group, such as a trityl or substituted trityl group, for example, a dimethoxytrityl or mono-methoxytrityl group.

In another embodiment, R₄ of a compound of the invention comprises a *tert*-butyl, Fm (fluorenyl-methoxy), or allyl group.


In one embodiment, R₆ of a compound of the invention comprises a TFA (trifluoroacetyl) group.

In another embodiment, R₃, R₅, R₇ and R₈ of a compound of the invention are independently hydrogen.

In one embodiment, R₇ of a compound of the invention is independently isobutryl, dimethylformamide, or hydrogen.


In another embodiment, R₁₂ of a compound of the invention comprises a methyl group or ethyl group.

In one embodiment, the invention features a compound having Formula 27:




27

wherein “n” is an integer from about 0 to about 20, R₄ is H or a cationic salt, X is a siNA molecule or a portion thereof, and R₂₄ is a sulfur containing leaving group, for example a group comprising:




In one embodiment, the invention features a compound having Formula 39:



39

5 wherein "n" is an integer from about 0 to about 20, X is a siNA molecule or a portion thereof, and P is a phosphorus containing group.

In another embodiment, a thiol containing linker of the invention is a compound having Formula 41:




41

10

wherein "n" is an integer from about 0 to about 20, P is a phosphorus containing group, for example a phosphine, phosphite, or phosphate, and R₂₄ is any alkyl, substituted alkyl, alkoxy, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, or substituted alkynyl group with or without additional protecting groups.

15


In one embodiment, the invention features a compound having Formula 43:



43


wherein X comprises a siNA molecule or portion thereof; W comprises a degradable nucleic acid linker; Y comprises a linker molecule or amino acid that can be present or absent; Z comprises H, OH, O-alkyl, SH, S-alkyl, alkyl, substituted alkyl, aryl, substituted aryl, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, or label; n is an integer from about 1 to about 100; and N' is an integer from about 1 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 44:



44


wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent; n is an integer from about 1 to about 50, and PEG represents a compound having Formula 45:



45


wherein Z comprises H, OH, O-alkyl, SH, S-alkyl, alkyl, substituted alkyl, aryl, substituted aryl, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, or label; and n is an integer from about 1 to about 100. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 46:



46


wherein X comprises a siNA molecule or portion thereof; each W independently comprises linker molecule or chemical linkage that can be present or absent, Y comprises
 5 a linker molecule or chemical linkage that can be present or absent; and PEG represents a compound having Formula 45:



45

wherein Z comprises H, OH, O-alkyl, SH, S-alkyl, alkyl, substituted alkyl, aryl,
 10 substituted aryl, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, or label; and n is an integer from about 1 to about 100. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

15 In one embodiment, the invention features a compound having Formula 47:




47

wherein X comprises a siNA molecule or portion thereof; each W independently comprises a linker molecule or chemical linkage that can be the same or different and can
 20 be present or absent, Y comprises a linker molecule that can be present or absent; each Q

independently comprises a hydrophobic group or phospholipid; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and n is an integer from about 1 to about 10. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 48:



48

wherein X comprises a siNA molecule or portion thereof; each W independently comprises a linker molecule or chemical linkage that can be present or absent, Y comprises a linker molecule that can be present or absent; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and B represents a lipophilic group, for example a saturated or unsaturated linear, branched, or cyclic alkyl group, cholesterol, or a derivative thereof. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 49:




49

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, Y comprises a linker molecule that can be present or absent; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and B represents a lipophilic group, for example a saturated or unsaturated linear, branched, or cyclic alkyl group, cholesterol, or a derivative thereof. In another

embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 50:




5

50

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent; Y comprises a linker molecule or chemical linkage that can be present or absent; and each Q independently comprises a hydrophobic group or phospholipid. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In one embodiment, the invention features a compound having Formula 51:




51

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent; Y comprises a linker molecule or amino acid that can be present or absent; Z comprises H, OH, O-alkyl, SH, S-alkyl, alkyl, substituted alkyl, aryl, substituted aryl, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, or label; SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers, and n is an integer from about 1 to about 20. In another

embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 52:



5


52

wherein X comprises a siNA molecule or portion thereof; Y comprises a linker molecule or chemical linkage that can be present or absent; each R1, R2, R3, R4, and R5 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N; Z comprises H, OH, O-alkyl, SH, S-alkyl, alkyl, substituted alkyl, aryl, substituted aryl, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, or label; SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers, n is an integer from about 1 to about 20; and N' is an integer from about 1 to about 20. In another embodiment, X comprises a siNA molecule or a portion thereof. In another embodiment, Y is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

10

15

In another embodiment, the invention features a compound having Formula 53:




20

53

wherein B comprises H, a nucleoside base, or a non-nucleosidic base with or without protecting groups; each R1 independently comprises O, N, S, alkyl, or substituted N; each R2 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylhalo, S, N, substituted N, or a phosphorus containing group; each R3 independently comprises N or O-N, each R4 independently comprises O, CH₂, S, sulfone, or sulfoxy; X comprises H,
 5 a removable protecting group, a siNA molecule or a portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent; SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers,, each n is
 10 independently an integer from about 1 to about 50; and N' is an integer from about 1 to about 10. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 54:




15

54

wherein B comprises H, a nucleoside base, or a non-nucleosidic base with or without protecting groups; each R1 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylhalo, S, N, substituted N, or a phosphorus containing group; X comprises H, a removable protecting group, a siNA molecule or a portion thereof; W comprises a
 20 linker molecule or chemical linkage that can be present or absent; and SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


25 In one embodiment, the invention features a compound having Formula 55:



55

wherein each R1 independently comprises O, N, S, alkyl, or substituted N; each R2 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylhalo, S, N, substituted N, or a phosphorus containing group; each R3 independently comprises H, OH, alkyl, substituted alkyl, or halo; X comprises H, a removable protecting group, a siNA molecule or a portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent; SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers, each n is independently an integer from about 1 to about 50; and N' is an integer from about 1 to about 100. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 56:




56

wherein R1 comprises H, alkyl, alkylhalo, N, substituted N, or a phosphorus containing group; R2 comprises H, O, OH, alkyl, alkylhalo, halo, S, N, substituted N, or a phosphorus containing group; X comprises H, a removable protecting group, a siNA molecule or a portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent; SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L,


alpha or beta isomers, and each n is independently an integer from about 0 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 57:




57

wherein R1 can include the groups:




and wherein R2 can include the groups:



and wherein Tr is a removable protecting group, for example a trityl, monomethoxytrityl, or dimethoxytrityl; SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers, and n is an integer from about 1 to about 20.

In one embodiment, compounds having Formula 52, 53, 54, 55, 56, and 57 are featured wherein each nitrogen adjacent to a carbonyl can independently be substituted for a carbonyl adjacent to a nitrogen or each carbonyl adjacent to a nitrogen can be substituted for a nitrogen adjacent to a carbonyl.


In another embodiment, the invention features a compound having Formula 58:



58

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent; Y comprises a linker molecule or amino acid that can be present or absent; V comprises a signal protein or peptide, for example Human serum albumin protein, Antennapedia peptide, Kaposi fibroblast growth factor peptide, Caiman crocodylus Ig(5) light chain peptide, HIV envelope glycoprotein gp41 peptide, HIV-1 Tat peptide, Influenza hemagglutinin envelope glycoprotein peptide, or transportan A peptide; each n is independently an integer from about 1 to about 50; and N' is an integer from about 1 to about 100. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 59:



59

wherein each R1 independently comprises O, S, N, substituted N, or a phosphorus containing group; each R2 independently comprises O, S, or N; X comprises H, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, or other biologically active molecule; n is an integer from about 1 to about 50, Q comprises H or a removable protecting group which can be optionally absent, each W independently comprises a linker molecule or chemical linkage that can be present or absent, and V comprises a signal protein or peptide, for example Human serum albumin protein, Antennapedia peptide, Kaposi fibroblast growth factor peptide, Caiman crocodylus Ig(5) light chain peptide, HIV envelope glycoprotein gp41 peptide, HIV-1 Tat peptide,


Influenza hemagglutinin envelope glycoprotein peptide, or transportan A peptide, or a compound having Formula 45



45


- 5 wherein Z comprises H, OH, O-alkyl, SH, S-alkyl, alkyl, substituted alkyl, aryl, substituted aryl, amino, substituted amino, a removable protecting group, a siNA molecule or a portion thereof; and n is an integer from about 1 to about 100. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

- 10 In another embodiment, the invention features a compound having Formula 60:




60

wherein R1 can include the groups:




- 15 and wherein R2 can include the groups:



and wherein Tr is a removable protecting group, for example a trityl, monomethoxytrityl, or dimethoxytrityl; n is an integer from about 1 to about 50; and R8 is a nitrogen protecting group, for example a phthaloyl, trifluoroacetyl, Fmoc, or monomethoxytrityl group.


5 In another embodiment, the invention features a compound having Formula 61:



61

wherein X comprises a siNA molecule or portion thereof; each W independently comprises a linker molecule or chemical linkage that can be the same or different and can be present or absent, Y comprises a linker molecule that can be present or absent; each 10 independently comprises a signal protein or peptide, for example Human serum albumin protein, Antennapedia peptide, Kaposi fibroblast growth factor peptide, Caiman crocodylus Ig(5) light chain peptide, HIV envelope glycoprotein gp41 peptide, HIV-1 Tat peptide, Influenza hemagglutinin envelope glycoprotein peptide, or transportan A peptide;; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, 15 O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and n is an integer from about 1 to about 10. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

20 In another embodiment, the invention features a compound having Formula 62:




62

wherein X comprises a siNA molecule or portion thereof; each 5 independently comprises a signal protein or peptide, for example Human serum albumin protein, Antennapedia peptide, Kaposi fibroblast growth factor peptide, Caiman crocodylus Ig(5) 25

light chain peptide, HIV envelope glycoprotein gp41 peptide, HIV-1 Tat peptide, Influenza hemagglutinin envelope glycoprotein peptide, or transportan A peptide; W comprises a linker molecule or chemical linkage that can be present or absent; each R1, R2, and R3 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and each n is independently an integer from about 1 to about 10. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 63:



10


63

wherein X comprises a siNA molecule or portion thereof; V comprises a signal protein or peptide, for example Human serum albumin protein, Antennapedia peptide, Kaposi fibroblast growth factor peptide, Caiman crocodylus Ig(5) light chain peptide, HIV envelope glycoprotein gp41 peptide, HIV-1 Tat peptide, Influenza hemagglutinin envelope glycoprotein peptide, or transportan A peptide; W comprises a linker molecule or chemical linkage that can be present or absent; each R1, R2, R3 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, R4 represents an ester, amide, or protecting group, and each n is independently an integer from about 1 to about 10. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

15

20


In another embodiment, the invention features a compound having Formula 64:



64

wherein X comprises a siNA molecule or portion thereof; each W independently comprises a linker molecule or chemical linkage that can be present or absent, Y comprises a linker molecule that can be present or absent; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, A comprises a nitrogen containing group, and B comprises a lipophilic group. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 65:



65

wherein X comprises a siNA molecule or portion thereof; each W independently comprises a linker molecule or chemical linkage that can be present or absent, Y comprises a linker molecule that can be present or absent; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, RV comprises the lipid or phospholipid component of any of Formulae 47-50, and R6 comprises a nitrogen containing group. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In another embodiment, the invention features a compound having Formula 92:




92

wherein B comprises H, a nucleoside base, or a non-nucleosidic base with or without protecting groups; each R1 independently comprises O, OH, H, alkyl, alkylhalo,

O-alkyl, O-alkylhalo, S, N, substituted N, or a phosphorus containing group; X comprises H, a removable protecting group, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, enzymatic nucleic acid, amino acid, peptide, protein, lipid, phospholipid, biologically active molecule or label; W comprises a linker molecule or chemical linkage that can be present or absent; R2 comprises O, NH, S, CO, COO, ON=C, or alkyl; R3 comprises alkyl, akloxy, or an aminoacyl side chain; and SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 86:



86

wherein R1 comprises H, alkyl, alkylhalo, N, substituted N, or a phosphorus containing group; R2 comprises H, O, OH, alkyl, alkylhalo, halo, S, N, substituted N, or a phosphorus containing group; X comprises H, a removable protecting group, a siNA molecule or a portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent; R3 comprises O, NH, S, CO, COO, ON=C, or alkyl; R4 comprises alkyl, akloxy, or an aminoacyl side chain; and SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers, and each n is independently an integer from about 0 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 87:



87

wherein X comprises a protein, peptide, antibody, lipid, phospholipid, oligosaccharide, label, biologically active molecule, for example a vitamin such as folate, vitamin A, E, B6, B12, coenzyme, antibiotic, antiviral, nucleic acid, nucleotide, nucleoside, or oligonucleotide such as an enzymatic nucleic acid, allozyme, antisense nucleic acid, siNA, 2,5-A chimera, decoy, aptamer or triplex forming oligonucleotide, or polymers such as polyethylene glycol; W comprises a linker molecule or chemical linkage that can be present or absent; and Y comprises siNA or a portion thereof; R1 comprises H, alkyl, or substituted alkyl. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 88:



88

wherein X comprises a protein, peptide, antibody, lipid, phospholipid, oligosaccharide, label, biologically active molecule, for example a vitamin such as folate, vitamin A, E, B6, B12, coenzyme, antibiotic, antiviral, nucleic acid, nucleotide, nucleoside, or oligonucleotide such as an enzymatic nucleic acid, allozyme, antisense nucleic acid, siNA, 2,5-A chimera, decoy, aptamer or triplex forming oligonucleotide, or polymers such as polyethylene glycol; W comprises a linker molecule or chemical linkage that can be present or absent, and Y comprises a siNA or a portion thereof. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 99:



99

wherein X comprises a siNA molecule or portion thereof; each W independently comprises a linker molecule or chemical linkage that can be present or absent, Y comprises a linker molecule that can be present or absent; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine or branched derivative thereof, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 100:



100

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, Y comprises a linker molecule that can be present or absent; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and SG comprises a sugar, for example galactose, galactosamine, N-acetyl-galactosamine or branched derivative thereof, glucose, mannose, fructose, or fucose and the respective D or L, alpha or beta isomers. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In one embodiment, the SG component of any compound having Formulae 99 or 100 comprises a compound having Formula 101:



101

wherein Y comprises a linker molecule or chemical linkage that can be present or absent and each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group.


In one embodiment, the W-SG component of a compound having Formulae 99
5 comprises a compound having Formula 102:



102

wherein R2 comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylhalo, S, N, substituted N, a protecting group, or another compound having Formula 102; R1
10 independently H, OH, alkyl, substituted alkyl, or halo and each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group, and R3 comprises O or R3 in Formula 99, and n is an integer from about 1 to about 20.


In one embodiment, the W-SG component of a compound having Formulae 99
comprises a compound having Formula 103:



103

wherein R1 comprises H, alkyl, alkylhalo, O-alkyl, O-alkylhalo, S, N, substituted N, a protecting group, or another compound having Formula 103; each R7 independently
15 comprises an acyl group that can be present or absent, for example a acetyl group, and R3 comprises H or R3 in Formula 99, and each n is independently an integer from about 1 to about 20.


In one embodiment, the invention features a compound having Formula 104:




104

wherein R3 comprises H, OH, amino, substituted amino, nucleotide, nucleoside,
 5 nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a
 portion thereof, or OR5 where R5 a removable protecting group, R4 comprises O, alkyl,
 alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, each R7
 independently comprises an acyl group that can be present or absent, for example a acetyl
 group, and each n is independently an integer from about 1 to about 20, and


10 wherein R1 can include the groups:



and wherein R2 can include the groups:




15 In one embodiment, the invention features a compound having Formula 105:



105

wherein X comprises a siNA molecule or a portion thereof, R2 comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylhalo, S, N, substituted N, a protecting group, or a nucleotide, polynucleotide, or oligonucleotide or a portion thereof; R1 independently H, OH, alkyl, substituted alkyl, or halo and each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group, and n is an integer from about 1 to about 20.


In one embodiment, the invention features a compound having Formula 106:



106

wherein X comprises a siNA molecule or a portion thereof, R1 comprises H, OH, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a portion thereof, or OR5 where R5 a removable protecting group, each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group, and each n is independently an integer from about 1 to about 20


In another embodiment, the invention features a compound having Formula 107:



107

wherein X comprises a siNA molecule or portion thereof; each W independently comprises a linker molecule or chemical linkage that can be present or absent, Y comprises a linker molecule that can be present or absent; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and Cholesterol comprises cholesterol or an analog, derivative, or metabolite thereof. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In another embodiment, the invention features a compound having Formula 108:



108

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, Y comprises a linker molecule that can be present or absent; each R1, R2, R3, and R4 independently comprises O, OH, H, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, and Cholesterol comprises cholesterol or an analog, derivative, or metabolite thereof. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In one embodiment, the W-Cholesterol component of a compound having Formula 107 comprises a compound having Formula 109:



109

wherein R3 comprises R3 as described in Formula 107, and n is independently an integer from about 1 to about 20.


5 In one embodiment, the invention features a compound having Formula 110:




110

10 wherein R4 comprises O, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, each n is independently an integer from about 1 to about 20, and


wherein R1 can include the groups:



and wherein R2 can include the groups:



In one embodiment, the invention features a compound having Formula 111:




5

111

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, and n is an integer from about 1 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

10

In one embodiment, the invention features a compound having Formula 112:




112

wherein n is an integer from about 1 to about 20. In another embodiment, a compound having Formula 112 is used to generate a compound having Formula 111 via NHS ester mediated coupling with a biologically active molecule, such as a siNA molecule or a portion thereof. In a non-limiting example, the NHS ester coupling can be effectuated via attachment to a free amine present in the siNA molecule, such as an amino linker molecule present on a nucleic acid sugar (e.g. 2'-amino linker) or base (e.g., C5 alkyl amine linker) component of the siNA molecule.


15

In one embodiment, the invention features a compound having Formula 113:




wherein R3 comprises H, OH, amino, substituted amino, nucleotide, nucleoside,
 5 nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a
 portion thereof, or OR5 where R5 a removable protecting group, R4 comprises O, alkyl,
 alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, each n is
 independently an integer from about 1 to about 20, and

wherein R1 can include the groups:




and wherein R2 can include the groups:



In another embodiment, a compound having Formula 113 is used to generate a
 15 compound having Formula 111 via phosphoramidite mediated coupling with a
 biologically active molecule, such as a siNA molecule or a portion thereof.


In one embodiment, the invention features a compound having Formula 114:



114

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, and n is an integer from about 1 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In one embodiment, the invention features a compound having Formula 115:



115

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, R3 comprises H, OH, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a portion thereof, or OR5 where R5 a removable protecting group, and each n is independently an integer from about 1 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In one embodiment, the invention features a compound having Formula 116:




141


116

wherein R3 comprises H, OH, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a portion thereof, or OR5 where R5 a removable protecting group, R4 comprises O, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, each n is
 5 independently an integer from about 1 to about 20, and

wherein R1 can include the groups:




and wherein R2 can include the groups:



10

In another embodiment, a compound having Formula 116 is used to generate a compound having Formula 114 or 115 via phosphoramidite mediated coupling with a biologically active molecule, such as a siNA molecule or a portion thereof.

15 In one embodiment, the invention features a compound having Formula 117:




117


wherein R3 comprises H, OH, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a

portion thereof, or OR5 where R5 a removable protecting group, R4 comprises O, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group, each n is independently an integer from about 1 to about 20, and

5 wherein R1 can include the groups:




and wherein R2 can include the groups:



10 In another embodiment, a compound having Formula 117 is used to generate a compound having Formula 105 via phosphoramidite mediated coupling with a biologically active molecule, such as a siNA molecule or a portion thereof.

In one embodiment, the invention features a compound having Formula 118:




15

118

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, R3 comprises H, OH, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a portion thereof, or OR5 where R5 a removable protecting group, each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group, and each n is independently an integer from about 1 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.


In one embodiment, the invention features a compound having Formula 119:



119

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group, and each n is independently an integer from about 1 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In one embodiment, the invention features a compound having Formula 120:




144


120

wherein R3 comprises H, OH, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a portion thereof, or OR5 where R5 a removable protecting group, R4 comprises O, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group, each n is independently an integer from about 1 to about 20, and

wherein R1 can include the groups:




and wherein R2 can include the groups:



In another embodiment, a compound having Formula 120 is used to generate a compound having Formula 118 or 119 via phosphoramidite mediated coupling with a biologically active molecule, such as a siNA molecule or a portion thereof.

In one embodiment, the invention features a compound having Formula 121:




145

121

wherein X comprises a siNA molecule or portion thereof; W comprises a linker molecule or chemical linkage that can be present or absent, each R7 independently comprises an acyl group that can be present or absent, for example a acetyl group, and
 5 each n is independently an integer from about 1 to about 20. In another embodiment, W is selected from the group consisting of amide, phosphate, phosphate ester, phosphoramidate, or thiophosphate ester linkage.

In one embodiment, the invention features a compound having Formula 122:




10

122


wherein R3 comprises H, OH, amino, substituted amino, nucleotide, nucleoside, nucleic acid, oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, label, or a portion thereof, or OR5 where R5 a removable protecting group, R4 comprises O, alkyl, alkylhalo, O-alkyl, O-alkylcyano, S, S-alkyl, S-alkylcyano, N or substituted N, each R7
 15 independently comprises an acyl group that can be present or absent, for example a acetyl group, each n is independently an integer from about 1 to about 20, and

wherein R1 can include the groups:



and wherein R2 can include the groups:

146



In another embodiment, a compound having Formula 122 is used to generate a compound having Formula 121 via phosphoramidite mediated coupling with a biologically active molecule, such as a siNA molecule or a portion thereof.

5 In one embodiment, the invention features a compound having Formula 94,



94

wherein X comprises a siNA molecule or a portion thereof; each Y independently comprises a linker or chemical linkage that can be present or absent, W comprises a
10 biodegradable nucleic acid linker molecule, and Z comprises a biologically active molecule, for example an enzymatic nucleic acid, allozyme, antisense nucleic acid, siNA, 2,5-A chimera, decoy, aptamer or triplex forming oligonucleotide, peptide, protein, or antibody.

In another embodiment, W of a compound having Formula 94 of the invention
15 comprises 5'-cytidine-deoxythymidine-3', 5'-deoxythymidine-cytidine-3', 5'-cytidine-deoxyuridine-3', 5'-deoxyuridine-cytidine-3', 5'-uridine-deoxythymidine-3', or 5'-deoxythymidine-uridine-3'.

In yet another embodiment, W of a compound having Formula 94 of the invention
20 comprises 5'-adenosine-deoxythymidine-3', 5'-deoxythymidine-adenosine-3', 5'-adenosine-deoxyuridine-3', or 5'-deoxyuridine-adenosine-3'.


In another embodiment, Y of a compound having Formula 94 of the invention comprises a phosphorus containing linkage, phosphoramidate linkage, phosphodiester linkage, phosphorothioate linkage, amide linkage, ester linkage, carbamate linkage, disulfide linkage, oxime linkage, or morpholino linkage.

25 In another embodiment, compounds having Formula 89 and 91 of the invention are synthesized by periodate oxidation of an N-terminal Serine or Threonine residue of a peptide or protein.

In one embodiment, X of compounds having Formulae 43, 44, 46-52, 58, 61-65, 85-88, 92, 94, 95, 99, 100, 105-108, 111, 114, 115, 118, 119, or 121 of the invention comprises a siNA molecule or a portion thereof. In one embodiment, the siNA molecule can be conjugated at the 5' end, 3'-end, or both 5' and 3' ends of the sense strand or region of the siNA. In one embodiment, the siNA molecule can be conjugated at the 3'-end of the antisense strand or region of the siNA with a compound of the invention. In one embodiment, both the sense strand and antisense strands or regions of the siNA molecule are conjugated with a compound of the invention. In one embodiment, only the sense strand or region of the siNA is conjugated with a compound of the invention. In one embodiment, only the antisense strand or region of the siNA is conjugated with a compound of the invention.

In one embodiment, W and/or Y of compounds having Formulae 43, 44, 46-52, 58, 61-65, 85-88, 92, 94, 95, 99, 100, 101, 107, 108, 111, 114, 115, 118, 119, or 121 of the invention comprises a degradable or cleavable linker, for example a nucleic acid sequence comprising ribonucleotides and/or deoxynucleotides, such as a dimer, trimer, or tetramer. A non limiting example of a nucleic acid cleavable linker is an adenosine-deoxythymidine (A-dT) dimer or a cytidine-deoxythymidine (C-dT) dimer. In yet another embodiment, W and/or V of compounds having Formulae 43, 44, 48-51, 58, 63-65, 96, 99, 100, 107, 108, 111, 114, 115, 118, 119, or 121 of the invention comprises a N-hydroxy succinimide (NHS) ester linkage, oxime linkage, disulfide linkage, phosphoramidate, phosphorothioate, phosphorodithioate, phosphodiester linkage, or NHC(O), CH₃NC(O), CONH, C(O)NCH₃, S, SO, SO₂, O, NH, NCH₃ group. In another embodiment, the degradable linker, W and/or Y, of compounds having Formulae 43, 44, 46-52, 58, 61-65, 85-88, 92, 94, 95, 99, 100, 101, 107, 108, 111, 114, 115, 118, 119, or 121 of the invention comprises a linker that is susceptible to cleavage by carboxypeptidase activity.

In another embodiment, W and/or Y of Formulae 43, 44, 46-52, 58, 61-65, 85-88, 92, 94, 95, 99, 100, 101, 107, 108, 111, 114, 115, 118, 119, or 121 comprises a polyethylene glycol linker having Formula 45:



45

wherein Z comprises H, OH, O-alkyl, SH, S-alkyl, alkyl, substituted alkyl, aryl, substituted aryl, amino, substituted amino, nucleotide, nucleoside, nucleic acid,

oligonucleotide, amino acid, peptide, protein, lipid, phospholipid, or label; and n is an integer from about 1 to about 100.

In one embodiment, the nucleic acid conjugates of the instant invention are assembled by solid phase synthesis, for example on an automated peptide synthesizer, for example a Miligen 9050 synthesizer and/or an automated oligonucleotide synthesizer
5 such as an ABI 394, 390Z, or Pharmacia OligoProcess, OligoPilot, OligoMax, or AKTA synthesizer. In another embodiment, the nucleic acid conjugates of the invention are assembled post synthetically, for example, following solid phase oligonucleotide synthesis (see for example **Figures 45, 50, 53, and 73**).

10 In another embodiment, V of compounds having Formula 58-63 and 96 comprise peptides having SEQ ID NOS: **507-516 (Table V)**.

In one embodiment, the nucleic acid conjugates of the instant invention are assembled post synthetically, for example, following solid phase oligonucleotide synthesis.

15 The present invention provides compositions and conjugates comprising nucleosidic and non-nucleosidic derivatives. The present invention also provides nucleic acid, polynucleotide and oligonucleotide derivatives including RNA, DNA, and PNA based conjugates. The attachment of compounds of the invention to nucleosides, nucleotides, non-nucleosides, and nucleic acid molecules is provided at any position
20 within the molecule, for example, at internucleotide linkages, nucleosidic sugar hydroxyl groups such as 5', 3', and 2'-hydroxyls, and/or at nucleobase positions such as amino and carbonyl groups.

The exemplary conjugates of the invention are described as compounds of the formulae herein, however, other peptide, protein, phospholipid, and poly-alkyl glycol
25 derivatives are provided by the invention, including various analogs of the compounds of formulae 1-122, including but not limited to different isomers of the compounds described herein.

The exemplary folate conjugates of the invention are described as compounds shown by formulae herein, however, other folate and antifolate derivatives are provided

by the invention, including various folate analogs of the formulae of the invention, including dihydrofolates, tetrahydrofolates, tetrahydropterins, folinic acid, pteropolyglutamic acid, 1-deza, 3-deaza, 5-deaza, 8-deaza, 10-deaza, 1,5-deaza, 5,10 dideaza, 8,10-dideaza, and 5,8-dideaza folates, antifolates, and pteronic acids. As used
5 herein, the term "folate" is meant to refer to folate and folate derivatives, including pteronic acid derivatives and analogs.

The present invention features compositions and conjugates to facilitate delivery of molecules into a biological system such as cells. The conjugates provided by the instant invention can impart therapeutic activity by transferring therapeutic compounds across
10 cellular membranes. The present invention encompasses the design and synthesis of novel agents for the delivery of molecules, including but not limited to siNA molecules. In general, the transporters described are designed to be used either individually or as part of a multi-component system. The compounds of the invention generally shown in Formulae herein are expected to improve delivery of molecules into a number of cell
15 types originating from different tissues, in the presence or absence of serum.

In another embodiment, the compounds of the invention are provided as a surface component of a lipid aggregate, such as a liposome encapsulated with the predetermined molecule to be delivered. Liposomes, which can be unilamellar or multilamellar, can introduce encapsulated material into a cell by different mechanisms. For example, the
20 liposome can directly introduce its encapsulated material into the cell cytoplasm by fusing with the cell membrane. Alternatively, the liposome can be compartmentalized into an acidic vacuole (i.e., an endosome) and its contents released from the liposome and out of the acidic vacuole into the cellular cytoplasm.

In one embodiment the invention features a lipid aggregate formulation of the compounds described herein, including phosphatidylcholine (of varying chain length; e.g., egg yolk phosphatidylcholine), cholesterol, a cationic lipid, and 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-polythyleneglycol-2000 (DSPE-PEG2000). The cationic lipid component of this lipid aggregate can be any cationic lipid known in the art such as dioleoyl 1,2-diacyl-3-trimethylammonium-propane (DOTAP). In another
25 30 embodiment this cationic lipid aggregate comprises a covalently bound compound described in any of the Formulae herein.

In another embodiment, polyethylene glycol (PEG) is covalently attached to the compounds of the present invention. The attached PEG can be any molecular weight but is preferably between 2000-50,000 daltons.

5 The compounds and methods of the present invention are useful for introducing nucleotides, nucleosides, nucleic acid molecules, lipids, peptides, proteins, and/or non-nucleosidic small molecules into a cell. For example, the invention can be used for nucleotide, nucleoside, nucleic acid, lipids, peptides, proteins, and/or non-nucleosidic small molecule delivery where the corresponding target site of action exists intracellularly.

10 In one embodiment, the compounds of the instant invention provide conjugates of molecules that can interact with cellular receptors, such as high affinity folate receptors and ASGPr receptors, and provide a number of features that allow the efficient delivery and subsequent release of conjugated compounds across biological membranes. The compounds utilize chemical linkages between the receptor ligand and the compound to be
15 delivered of length that can interact preferentially with cellular receptors. Furthermore, the chemical linkages between the ligand and the compound to be delivered can be designed as degradable linkages, for example by utilizing a phosphate linkage that is proximal to a nucleophile, such as a hydroxyl group. Deprotonation of the hydroxyl group or an equivalent group, as a result of pH or interaction with a nuclease, can result in
20 nucleophilic attack of the phosphate resulting in a cyclic phosphate intermediate that can be hydrolyzed. This cleavage mechanism is analogous RNA cleavage in the presence of a base or RNA nuclease. Alternately, other degradable linkages can be selected that respond to various factors such as UV irradiation, cellular nucleases, pH, temperature etc. The use of degradable linkages allows the delivered compound to be released in a
25 predetermined system, for example in the cytoplasm of a cell, or in a particular cellular organelle.

The present invention also provides ligand derived phosphoramidites that are readily conjugated to compounds and molecules of interest. Phosphoramidite compounds of the invention permit the direct attachment of conjugates to molecules of interest
30 without the need for using nucleic acid phosphoramidite species as scaffolds. As such, the use of phosphoramidite chemistry can be used directly in coupling the compounds of

the invention to a compound of interest, without the need for other condensation reactions, such as condensation of the ligand to an amino group on the nucleic acid, for example at the N6 position of adenosine or a 2'-deoxy-2'-amino function. Additionally, compounds of the invention can be used to introduce non-nucleic acid based conjugated linkages into oligonucleotides that can provide more efficient coupling during oligonucleotide synthesis than the use of nucleic acid-based phosphoramidites. This improved coupling can take into account improved steric considerations of abasic or non-nucleosidic scaffolds bearing pendant alkyl linkages.

Compounds of the invention utilizing triphosphate groups can be utilized in the enzymatic incorporation of conjugate molecules into oligonucleotides. Such enzymatic incorporation is useful when conjugates are used in post-synthetic enzymatic conjugation or selection reactions, (see for example Matulic-Adamic *et al.*, 2000, *Bioorg. Med. Chem. Lett.*, 10, 1299-1302; Lee *et al.*, 2001, *NAR.*, 29, 1565-1573; Joyce, 1989, *Gene*, 82, 83-87; Beaudry *et al.*, 1992, *Science* 257, 635-641; Joyce, 1992, *Scientific American* 267, 90-97; Breaker *et al.*, 1994, *TIBTECH* 12, 268; Bartel *et al.*, 1993, *Science* 261:1411-1418; Szostak, 1993, *TIBS* 17, 89-93; Kumar *et al.*, 1995, *FASEB J.*, 9, 1183; Breaker, 1996, *Curr. Op. Biotech.*, 7, 442; Santoro *et al.*, 1997, *Proc. Natl. Acad. Sci.*, 94, 4262; Tang *et al.*, 1997, *RNA* 3, 914; Nakamaye & Eckstein, 1994, *supra*; Long & Uhlenbeck, 1994, *supra*; Ishizaka *et al.*, 1995, *supra*; Vaish *et al.*, 1997, *Biochemistry* 36, 6495; Kuwabara *et al.*, 2000, *Curr. Opin. Chem. Biol.*, 4, 669).

The term "biodegradable linker" as used herein, refers to a nucleic acid or non-nucleic acid linker molecule that is designed as a biodegradable linker to connect one molecule to another molecule, for example, a biologically active molecule to a siNA molecule of the invention or the sense and antisense strands of a siNA molecule of the invention. The biodegradable linker is designed such that its stability can be modulated for a particular purpose, such as delivery to a particular tissue or cell type. The stability of a nucleic acid-based biodegradable linker molecule can be modulated by using various chemistries, for example combinations of ribonucleotides, deoxyribonucleotides, and chemically-modified nucleotides, such as 2'-O-methyl, 2'-fluoro, 2'-amino, 2'-O-amino, 2'-C-allyl, 2'-O-allyl, and other 2'-modified or base modified nucleotides. The biodegradable nucleic acid linker molecule can be a dimer, trimer, tetramer or longer nucleic acid molecule, for example, an oligonucleotide of about 2, 3, 4, 5, 6, 7, 8, 9, 10,

11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides in length, or can comprise a single nucleotide with a phosphorus-based linkage, for example, a phosphoramidate or phosphodiester linkage. The biodegradable nucleic acid linker molecule can also comprise nucleic acid backbone, nucleic acid sugar, or nucleic acid base modifications.

5 The term "biodegradable" as used herein, refers to degradation in a biological system, for example enzymatic degradation or chemical degradation.

 The term "biologically active molecule" as used herein, refers to compounds or molecules that are capable of eliciting or modifying a biological response in a system. Non-limiting examples of biologically active siNA molecules either alone or in
10 combination with other molecules contemplated by the instant invention include therapeutically active molecules such as antibodies, cholesterol, hormones, antivirals, peptides, proteins, chemotherapeutics, small molecules, vitamins, co-factors, nucleosides, nucleotides, oligonucleotides, enzymatic nucleic acids, antisense nucleic acids, triplex forming oligonucleotides, 2,5-A chimeras, siNA, dsRNA, allozymes, aptamers, decoys
15 and analogs thereof. Biologically active molecules of the invention also include molecules capable of modulating the pharmacokinetics and/or pharmacodynamics of other biologically active molecules, for example, lipids and polymers such as polyamines, polyamides, polyethylene glycol and other polyethers.

 The term "phospholipid" as used herein, refers to a hydrophobic molecule
20 comprising at least one phosphorus group. For example, a phospholipid can comprise a phosphorus-containing group and saturated or unsaturated alkyl group, optionally substituted with OH, COOH, oxo, amine, or substituted or unsubstituted aryl groups.

 The term "alkyl" as used herein refers to a saturated aliphatic hydrocarbon, including straight-chain, branched-chain "isoalkyl", and cyclic alkyl groups. The term
25 "alkyl" also comprises alkoxy, alkyl-thio, alkyl-thio-alkyl, alkoxyalkyl, alkylamino, alkenyl, alkynyl, alkoxy, cycloalkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, C1-C6 hydrocarbyl, aryl or substituted aryl groups. Preferably, the alkyl group has 1 to 12 carbons. More preferably it is a lower alkyl of from about 1 to about 7 carbons, more preferably about 1 to about 4 carbons. The alkyl group can be substituted
30 or unsubstituted. When substituted the substituted group(s) preferably comprise hydroxy, oxy, thio, amino, nitro, cyano, alkoxy, alkyl-thio, alkyl-thio-alkyl, alkoxyalkyl,

alkylamino, silyl, alkenyl, alkynyl, alkoxy, cycloalkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, C1-C6 hydrocarbyl, aryl or substituted aryl groups. The term "alkyl" also includes alkenyl groups containing at least one carbon-carbon double bond, including straight-chain, branched-chain, and cyclic groups. Preferably, the alkenyl group has about 2 to about 12 carbons. More preferably it is a lower alkenyl of from about 2 to about 7 carbons, more preferably about 2 to about 4 carbons. The alkenyl group can be substituted or unsubstituted. When substituted the substituted group(s) preferably comprise hydroxy, oxy, thio, amino, nitro, cyano, alkoxy, alkyl-thio, alkyl-thio-alkyl, alkoxyalkyl, alkylamino, silyl, alkenyl, alkynyl, alkoxy, cycloalkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, C1-C6 hydrocarbyl, aryl or substituted aryl groups. The term "alkyl" also includes alkynyl groups containing at least one carbon-carbon triple bond, including straight-chain, branched-chain, and cyclic groups. Preferably, the alkynyl group has about 2 to about 12 carbons. More preferably it is a lower alkynyl of from about 2 to about 7 carbons, more preferably about 2 to about 4 carbons. The alkynyl group can be substituted or unsubstituted. When substituted the substituted group(s) preferably comprise hydroxy, oxy, thio, amino, nitro, cyano, alkoxy, alkyl-thio, alkyl-thio-alkyl, alkoxyalkyl, alkylamino, silyl, alkenyl, alkynyl, alkoxy, cycloalkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, C1-C6 hydrocarbyl, aryl or substituted aryl groups. Alkyl groups or moieties of the invention can also include aryl, alkylaryl, carbocyclic aryl, heterocyclic aryl, amide and ester groups. The preferred substituent(s) of aryl groups are halogen, trihalomethyl, hydroxyl, SH, OH, cyano, alkoxy, alkyl, alkenyl, alkynyl, and amino groups. An "alkylaryl" group refers to an alkyl group (as described above) covalently joined to an aryl group (as described above). Carbocyclic aryl groups are groups wherein the ring atoms on the aromatic ring are all carbon atoms. The carbon atoms are optionally substituted. Heterocyclic aryl groups are groups having from about 1 to about 3 heteroatoms as ring atoms in the aromatic ring and the remainder of the ring atoms are carbon atoms. Suitable heteroatoms include oxygen, sulfur, and nitrogen, and include furanyl, thienyl, pyridyl, pyrrolyl, N-lower alkyl pyrrolo, pyrimidyl, pyrazinyl, imidazolyl and the like, all optionally substituted. An "amide" refers to an -C(O)-NH-R, where R is either alkyl, aryl, alkylaryl or hydrogen. An "ester" refers to an -C(O)-OR', where R is either alkyl, aryl, alkylaryl or hydrogen.

The term “alkoxyalkyl” as used herein refers to an alkyl-O-alkyl ether, for example, methoxyethyl or ethoxymethyl.

The term “alkyl-thio-alkyl” as used herein refers to an alkyl-S-alkyl thioether, for example, methylthiomethyl or methylthioethyl.

5 The term “amino” as used herein refers to a nitrogen containing group as is known in the art derived from ammonia by the replacement of one or more hydrogen radicals by organic radicals. For example, the terms “aminoacyl” and “aminoalkyl” refer to specific N-substituted organic radicals with acyl and alkyl substituent groups respectively.

10 The term “amination” as used herein refers to a process in which an amino group or substituted amine is introduced into an organic molecule.

The term “exocyclic amine protecting moiety” as used herein refers to a nucleobase amino protecting group compatible with oligonucleotide synthesis, for example, an acyl or amide group.

15 The term “alkenyl” as used herein refers to a straight or branched hydrocarbon of a designed number of carbon atoms containing at least one carbon-carbon double bond. Examples of “alkenyl” include vinyl, allyl, and 2-methyl-3-heptene.

20 The term “alkoxy” as used herein refers to an alkyl group of indicated number of carbon atoms attached to the parent molecular moiety through an oxygen bridge. Examples of alkoxy groups include, for example, methoxy, ethoxy, propoxy and isopropoxy.

The term “alkynyl” as used herein refers to a straight or branched hydrocarbon of a designed number of carbon atoms containing at least one carbon-carbon triple bond. Examples of “alkynyl” include propargyl, propyne, and 3-hexyne.

25 The term “aryl” as used herein refers to an aromatic hydrocarbon ring system containing at least one aromatic ring. The aromatic ring can optionally be fused or otherwise attached to other aromatic hydrocarbon rings or non-aromatic hydrocarbon rings. Examples of aryl groups include, for example, phenyl, naphthyl, 1,2,3,4-

tetrahydronaphthalene and biphenyl. Preferred examples of aryl groups include phenyl and naphthyl.

The term "cycloalkenyl" as used herein refers to a C3-C8 cyclic hydrocarbon containing at least one carbon-carbon double bond. Examples of cycloalkenyl include
5 cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclopentadiene, cyclohexenyl, 1,3-cyclohexadiene, cycloheptenyl, cycloheptatrienyl, and cyclooctenyl.

The term "cycloalkyl" as used herein refers to a C3-C8 cyclic hydrocarbon. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

10 The term "cycloalkylalkyl," as used herein, refers to a C3-C7 cycloalkyl group attached to the parent molecular moiety through an alkyl group, as defined above. Examples of cycloalkylalkyl groups include cyclopropylmethyl and cyclopentylethyl.

The terms "halogen" or "halo" as used herein refers to indicate fluorine, chlorine, bromine, and iodine.

15 The term "heterocycloalkyl," as used herein refers to a non-aromatic ring system containing at least one heteroatom selected from nitrogen, oxygen, and sulfur. The heterocycloalkyl ring can be optionally fused to or otherwise attached to other heterocycloalkyl rings and/or non-aromatic hydrocarbon rings. Preferred heterocycloalkyl groups have from 3 to 7 members. Examples of heterocycloalkyl groups
20 include, for example, piperazine, morpholine, piperidine, tetrahydrofuran, pyrrolidine, and pyrazole. Preferred heterocycloalkyl groups include piperidinyl, piperazinyl, morpholinyl, and pyrrolidinyl.

The term "heteroaryl" as used herein refers to an aromatic ring system containing at least one heteroatom selected from nitrogen, oxygen, and sulfur. The heteroaryl ring can
25 be fused or otherwise attached to one or more heteroaryl rings, aromatic or non-aromatic hydrocarbon rings or heterocycloalkyl rings. Examples of heteroaryl groups include, for example, pyridine, furan, thiophene, 5,6,7,8-tetrahydroisoquinoline and pyrimidine. Preferred examples of heteroaryl groups include thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, pyrimidyl, imidazolyl, benzimidazolyl, furanyl, benzofuranyl, thiazolyl,

benzothiazolyl, isoxazolyl, oxadiazolyl, isothiazolyl, benzisothiazolyl, triazolyl, tetrazolyl, pyrrolyl, indolyl, pyrazolyl, and benzopyrazolyl.

The term "C1-C6 hydrocarbyl" as used herein refers to straight, branched, or cyclic alkyl groups having 1-6 carbon atoms, optionally containing one or more carbon-carbon
5 double or triple bonds. Examples of hydrocarbyl groups include, for example, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, 3-methylpentyl, vinyl, 2-pentene, cyclopropylmethyl, cyclopropyl, cyclohexylmethyl, cyclohexyl and propargyl. When reference is made
10 herein to C1-C6 hydrocarbyl containing one or two double or triple bonds it is understood that at least two carbons are present in the alkyl for one double or triple bond, and at least four carbons for two double or triple bonds.

The term "protecting group" as used herein, refers to groups known in the art that are readily introduced and removed from an atom, for example O, N, P, or S. Protecting groups are used to prevent undesirable reactions from taking place that can compete with
15 the formation of a specific compound or intermediate of interest. See also "Protective Groups in Organic Synthesis", 3rd Ed., 1999, Greene, T. W. and related publications.


The term "nitrogen protecting group," as used herein, refers to groups known in the art that are readily introduced on to and removed from a nitrogen. Examples of nitrogen protecting groups include Boc, Cbz, benzoyl, and benzyl. See also "Protective Groups in
20 Organic Synthesis", 3rd Ed., 1999, Greene, T. W. and related publications.

The term "hydroxy protecting group," or "hydroxy protection" as used herein, refers to groups known in the art that are readily introduced on to and removed from an oxygen, specifically an -OH group. Examples of hydroxy protecting groups include trityl or substituted trityl groups, such as monomethoxytrityl and dimethoxytrityl, or substituted
25 silyl groups, such as tert-butyldimethyl, trimethylsilyl, or tert-butyldiphenyl silyl groups. See also "Protective Groups in Organic Synthesis", 3rd Ed., 1999, Greene, T. W. and related publications.


The term "acyl" as used herein refers to -C(O)R groups, wherein R is an alkyl or aryl.

The term “phosphorus containing group” as used herein, refers to a chemical group containing a phosphorus atom. The phosphorus atom can be trivalent or pentavalent, and can be substituted with O, H, N, S, C or halogen atoms. Examples of phosphorus containing groups of the instant invention include but are not limited to phosphorus atoms substituted with O, H, N, S, C or halogen atoms, comprising phosphonate, alkylphosphonate, phosphate, diphosphate, triphosphate, pyrophosphate, phosphorothioate, phosphorodithioate, phosphoramidate, phosphoramidite groups, nucleotides and nucleic acid molecules.


The term “phosphine” or “phosphite” as used herein refers to a trivalent phosphorus species, for example compounds having Formula 97:




wherein R can include the groups:




and wherein S and T independently include the groups:




The term “phosphate” as used herein refers to a pentavalent phosphorus species, for example a compound having Formula 98:



wherein R includes the groups:



and wherein S and T each independently can be a sulfur or oxygen atom or a group which can include:



5

and wherein M comprises a sulfur or oxygen atom. The phosphate of the invention can comprise a nucleotide phosphate, wherein any R, S, or T in Formula 98 comprises a linkage to a nucleic acid or nucleoside.

10 The term “cationic salt” as used herein refers to any organic or inorganic salt having a net positive charge, for example a triethylammonium (TEA) salt.

The term “degradable linker” as used herein, refers to linker moieties that are capable of cleavage under various conditions. Conditions suitable for cleavage can include but are not limited to pH, UV irradiation, enzymatic activity, temperature, hydrolysis, elimination, and substitution reactions, and thermodynamic properties of the linkage.

15

The term “photolabile linker” as used herein, refers to linker moieties as are known in the art, that are selectively cleaved under particular UV wavelengths. Compounds of the invention containing photolabile linkers can be used to deliver compounds to a target cell or tissue of interest, and can be subsequently released in the presence of a UV source.

20 The term “nucleic acid conjugates” as used herein, refers to nucleoside, nucleotide and oligonucleotide conjugates.

The term “lipid” as used herein, refers to any lipophilic compound. Non-limiting examples of lipid compounds include fatty acids and their derivatives, including straight chain, branched chain, saturated and unsaturated fatty acids, carotenoids, terpenes, bile acids, and steroids, including cholesterol and derivatives or analogs thereof.

5 The term “folate” as used herein, refers to analogs and derivatives of folic acid, for example antifolates, dihydrofolates, tetrahydrofolates, tetrahydropterins, folinic acid, pteropolyglutamic acid, 1-deza, 3-deaza, 5-deaza, 8-deaza, 10-deaza, 1,5-deaza, 5,10 dideaza, 8,10-dideaza, and 5,8-dideaza folates, antifolates, and pteric acid derivatives.

10 The term “compounds with neutral charge” as used herein, refers to compositions which are neutral or uncharged at neutral or physiological pH. Examples of such compounds are cholesterol and other steroids, cholesteryl hemisuccinate (CHEMS), dioleoyl phosphatidyl choline, distearoylphosphatidyl choline (DSPC), fatty acids such as oleic acid, phosphatidic acid and its derivatives, phosphatidyl serine, polyethylene glycol -conjugated phosphatidylamine, phosphatidylcholine, phosphatidylethanolamine and
15 related variants, prenylated compounds including farnesol, polyprenols, tocopherol, and their modified forms, diacylsuccinyl glycerols, fusogenic or pore forming peptides, dioleoylphosphatidylethanolamine (DOPE), ceramide and the like.

20 The term “lipid aggregate” as used herein refers to a lipid-containing composition wherein the lipid is in the form of a liposome, micelle (non-lamellar phase) or other aggregates with one or more lipids.

25 The term “nitrogen containing group” as used herein refers to any chemical group or moiety comprising a nitrogen or substituted nitrogen. Non-limiting examples of nitrogen containing groups include amines, substituted amines, amides, alkylamines, amino acids such as arginine or lysine, polyamines such as spermine or spermidine, cyclic amines such as pyridines, pyrimidines including uracil, thymine, and cytosine,
morpholines, phthalimides, and heterocyclic amines such as purines, including guanine and adenine.

30 Therapeutic nucleic acid molecules (e.g., siNA molecules) delivered exogenously optimally are stable within cells until reverse transcription of the RNA has been modulated long enough to reduce the levels of the RNA transcript. The nucleic acid

molecules are resistant to nucleases in order to function as effective intracellular therapeutic agents. Improvements in the chemical synthesis of nucleic acid molecules described in the instant invention and in the art have expanded the ability to modify nucleic acid molecules by introducing nucleotide modifications to enhance their nuclease stability as described above.

In yet another embodiment, siNA molecules having chemical modifications that maintain or enhance enzymatic activity of proteins involved in RNAi are provided. Such nucleic acids are also generally more resistant to nucleases than unmodified nucleic acids. Thus, *in vitro* and/or *in vivo* the activity should not be significantly lowered.

Use of the nucleic acid-based molecules of the invention will lead to better treatment of the disease progression by affording the possibility of combination therapies (e.g., multiple siNA molecules targeted to different genes; nucleic acid molecules coupled with known small molecule modulators; or intermittent treatment with combinations of molecules, including different motifs and/or other chemical or biological molecules). The treatment of subjects with siNA molecules can also include combinations of different types of nucleic acid molecules, such as enzymatic nucleic acid molecules (ribozymes), allozymes, antisense, 2,5-A oligoadenylate, decoys, and aptamers.

In another aspect a siNA molecule of the invention comprises one or more 5' and/or a 3'- cap structure, for example on only the sense siNA strand, the antisense siNA strand, or both siNA strands.

By "cap structure" is meant chemical modifications, which have been incorporated at either terminus of the oligonucleotide (see, for example, Adamic *et al.*, U.S. Pat. No. 5,998,203, incorporated by reference herein). These terminal modifications protect the nucleic acid molecule from exonuclease degradation, and can help in delivery and/or localization within a cell. The cap can be present at the 5'-terminus (5'-cap) or at the 3'-terminal (3'-cap) or can be present on both termini. Non-limiting examples of the 5'-cap include, but are not limited to, glyceryl, inverted deoxy abasic residue (moiety); 4',5'-methylene nucleotide; 1-(beta-D-erythrofuransyl) nucleotide, 4'-thio nucleotide; carbocyclic nucleotide; 1,5-anhydrohexitol nucleotide; L-nucleotides; alpha-nucleotides; modified base nucleotide; phosphorodithioate linkage; *threo*-pentofuransyl nucleotide; acyclic 3',4'-seco nucleotide; acyclic 3,4-dihydroxybutyl nucleotide; acyclic 3,5-

dihydroxypentyl nucleotide, 3'-3'-inverted nucleotide moiety; 3'-3'-inverted abasic moiety; 3'-2'-inverted nucleotide moiety; 3'-2'-inverted abasic moiety; 1,4-butanediol phosphate; 3'-phosphoramidate; hexylphosphate; aminohexyl phosphate; 3'-phosphate; 3'-phosphorothioate; phosphorodithioate; or bridging or non-bridging methylphosphonate moiety.

Non-limiting examples of the 3'-cap include, but are not limited to, glyceryl, inverted deoxy abasic residue (moiety), 4', 5'-methylene nucleotide; 1-(beta-D-erythrofuransyl) nucleotide; 4'-thio nucleotide, carbocyclic nucleotide; 5'-amino-alkyl phosphate; 1,3-diamino-2-propyl phosphate; 3-aminopropyl phosphate; 6-aminohexyl phosphate; 1,2-aminododecyl phosphate; hydroxypropyl phosphate; 1,5-anhydrohexitol nucleotide; L-nucleotide; alpha-nucleotide; modified base nucleotide; phosphorodithioate; *threo*-pentofuransyl nucleotide; acyclic 3',4'-seco nucleotide; 3,4-dihydroxybutyl nucleotide; 3,5-dihydroxypentyl nucleotide, 5'-5'-inverted nucleotide moiety; 5'-5'-inverted abasic moiety; 5'-phosphoramidate; 5'-phosphorothioate; 1,4-butanediol phosphate; 5'-amino; bridging and/or non-bridging 5'-phosphoramidate, phosphorothioate and/or phosphorodithioate, bridging or non bridging methylphosphonate and 5'-mercapto moieties (for more details see Beaucage and Iyer, 1993, *Tetrahedron* 49, 1925; incorporated by reference herein).

By the term "non-nucleotide" is meant any group or compound which can be incorporated into a nucleic acid chain in the place of one or more nucleotide units, including either sugar and/or phosphate substitutions, and allows the remaining bases to exhibit their enzymatic activity. The group or compound is abasic in that it does not contain a commonly recognized nucleotide base, such as adenosine, guanine, cytosine, uracil or thymine and therefore lacks a base at the 1'-position.

By "nucleotide" as used herein is as recognized in the art to include natural bases (standard), and modified bases well known in the art. Such bases are generally located at the 1' position of a nucleotide sugar moiety. Nucleotides generally comprise a base, sugar and a phosphate group. The nucleotides can be unmodified or modified at the sugar, phosphate and/or base moiety, (also referred to interchangeably as nucleotide analogs, modified nucleotides, non-natural nucleotides, non-standard nucleotides and other; see, for example, Usman and McSwiggen, *supra*; Eckstein *et al.*, International PCT

Publication No. WO 92/07065; Usman *et al.*, International PCT Publication No. WO 93/15187; Uhlman & Peyman, *supra*, all are hereby incorporated by reference herein). There are several examples of modified nucleic acid bases known in the art as summarized by Limbach *et al.*, 1994, *Nucleic Acids Res.* 22, 2183. Some of the non-limiting examples of base modifications that can be introduced into nucleic acid molecules include, inosine, purine, pyridin-4-one, pyridin-2-one, phenyl, pseudouracil, 2, 4, 6-trimethoxy benzene, 3-methyl uracil, dihydrouridine, naphthyl, aminophenyl, 5-alkylcytidines (*e.g.*, 5-methylcytidine), 5-alkyluridines (*e.g.*, ribothymidine), 5-halouridine (*e.g.*, 5-bromouridine) or 6-azapyrimidines or 6-alkylpyrimidines (*e.g.* 6-methyluridine), propyne, and others (Burgin *et al.*, 1996, *Biochemistry*, 35, 14090; Uhlman & Peyman, *supra*). By "modified bases" in this aspect is meant nucleotide bases other than adenine, guanine, cytosine and uracil at 1' position or their equivalents.

In one embodiment, the invention features modified siNA molecules, with phosphate backbone modifications comprising one or more phosphorothioate, phosphonoacetate, and/or thiophosphonoacetate, phosphorodithioate, methylphosphonate, phosphotriester, morpholino, amidate carbamate, carboxymethyl, acetamidate, polyamide, sulfonate, sulfonamide, sulfamate, formacetal, thioformacetal, and/or alkylsilyl, substitutions. For a review of oligonucleotide backbone modifications, see Hunziker and Leumann, 1995, *Nucleic Acid Analogues: Synthesis and Properties*, in *Modern Synthetic Methods*, VCH, 331-417, and Mesmaeker *et al.*, 1994, *Novel Backbone Replacements for Oligonucleotides*, in *Carbohydrate Modifications in Antisense Research*, ACS, 24-39.

By "abasic" is meant sugar moieties lacking a base or having other chemical groups in place of a base at the 1' position, see for example Adamic *et al.*, U.S. Pat. No. 5,998,203.

By "unmodified nucleoside" is meant one of the bases adenine, cytosine, guanine, thymine, or uracil joined to the 1' carbon of β -D-ribo-furanose.

By "modified nucleoside" is meant any nucleotide base which contains a modification in the chemical structure of an unmodified nucleotide base, sugar and/or phosphate. Non-limiting examples of modified nucleotides are shown by Formulae I-VII and/or other modifications described herein.

In connection with 2'-modified nucleotides as described for the present invention, by "amino" is meant 2'-NH₂ or 2'-O- NH₂, which can be modified or unmodified. Such modified groups are described, for example, in Eckstein *et al.*, U.S. Pat. No. 5,672,695 and Matulic-Adamic *et al.*, U.S. Pat. No. 6,248,878, which are both incorporated by
 5 reference in their entireties.

Various modifications to nucleic acid siNA structure can be made to enhance the utility of these molecules. Such modifications will enhance shelf-life, half-life *in vitro*, stability, and ease of introduction of such oligonucleotides to the target site, *e.g.*, to enhance penetration of cellular membranes, and confer the ability to recognize and bind
 10 to targeted cells.

Administration of Nucleic Acid Molecules

A siNA molecule of the invention can be adapted for use to treat any disease, infection or condition associated with gene expression, and other indications that can respond to the level of gene product in a cell or tissue, alone or in combination with other
 15 therapies. For example, a siNA molecule can comprise a delivery vehicle, including liposomes, for administration to a subject, carriers and diluents and their salts, and/or can be present in pharmaceutically acceptable formulations. Methods for the delivery of nucleic acid molecules are described in Akhtar *et al.*, 1992, *Trends Cell Bio.*, 2, 139; *Delivery Strategies for Antisense Oligonucleotide Therapeutics*, ed. Akhtar, 1995, Maurer
 20 *et al.*, 1999, *Mol. Membr. Biol.*, 16, 129-140; Hofland and Huang, 1999, *Handb. Exp. Pharmacol.*, 137, 165-192; and Lee *et al.*, 2000, *ACS Symp. Ser.*, 752, 184-192, all of which are incorporated herein by reference. Beigelman *et al.*, U.S. Pat. No. 6,395,713 and Sullivan *et al.*, PCT WO 94/02595 further describe the general methods for delivery of nucleic acid molecules. These protocols can be utilized for the delivery of virtually
 25 any nucleic acid molecule. Nucleic acid molecules can be administered to cells by a variety of methods known to those of skill in the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as biodegradable polymers, hydrogels, cyclodextrins (see for example Gonzalez *et al.*, 1999, *Bioconjugate Chem.*, 10, 1068-1074; Wang *et al.*, International PCT
 30 publication Nos. WO 03/47518 and WO 03/46185), poly(lactic-co-glycolic)acid (PLGA) and PLCA microspheres (see for example US Patent 6,447,796 and US Patent

Application Publication No. US 2002130430), biodegradable nanocapsules, and bioadhesive microspheres, or by proteinaceous vectors (O'Hare and Normand, International PCT Publication No. WO 00/53722). In one embodiment, nucleic acid molecules or the invention are administered via biodegradable implant materials, such as

5 elastic shape memory polymers (see for example Lendelein and Langer, 2002, *Science*, 296, 1673). Alternatively, the nucleic acid/vehicle combination is locally delivered by direct injection or by use of an infusion pump. Direct injection of the nucleic acid molecules of the invention, whether subcutaneous, intramuscular, or intradermal, can take place using standard needle and syringe methodologies, or by needle-free technologies

10 such as those described in Conry *et al.*, 1999, *Clin. Cancer Res.*, 5, 2330-2337 and Barry *et al.*, International PCT Publication No. WO 99/31262. Many examples in the art describe CNS delivery methods of oligonucleotides by osmotic pump, (see Chun *et al.*, 1998, *Neuroscience Letters*, 257, 135-138, D'Aldin *et al.*, 1998, *Mol. Brain Research*, 55, 151-164, Dryden *et al.*, 1998, *J. Endocrinol.*, 157, 169-175, Ghirmikar *et al.*, 1998,

15 *Neuroscience Letters*, 247, 21-24) or direct infusion (Broaddus *et al.*, 1997, *Neurosurg. Focus*, 3, article 4). Other routes of delivery include, but are not limited to oral (tablet or pill form) and/or intrathecal delivery (Gold, 1997, *Neuroscience*, 76, 1153-1158). More detailed descriptions of nucleic acid delivery and administration are provided in Sullivan *et al.*, *supra*, Draper *et al.*, PCT WO93/23569, Beigelman *et al.*, PCT WO99/05094, and

20 Klimuk *et al.*, PCT WO99/04819 all of which have been incorporated by reference herein. The molecules of the instant invention can be used as pharmaceutical agents. Pharmaceutical agents prevent, modulate the occurrence, or treat (alleviate a symptom to some extent, preferably all of the symptoms) of a disease state in a subject.

In addition, the invention features the use of methods to deliver the nucleic acid

25 molecules of the instant invention to hematopoietic cells, including monocytes and lymphocytes. These methods are described in detail by Hartmann *et al.*, 1998, *J. Pharmacol. Exp. Ther.*, 285(2), 920-928; Kronenwett *et al.*, 1998, *Blood*, 91(3), 852-862; Fillion and Phillips, 1997, *Biochim. Biophys. Acta.*, 1329(2), 345-356; Ma and Wei, 1996, *Leuk. Res.*, 20(11/12), 925-930; and Bongartz *et al.*, 1994, *Nucleic Acids Research*,

30 22(22), 4681-8. Such methods, as described above, include the use of free oligonucleotide, cationic lipid formulations, liposome formulations including pH sensitive liposomes and immunoliposomes, and bioconjugates including oligonucleotides

conjugated to fusogenic peptides, for the transfection of hematopoietic cells with oligonucleotides.

In one embodiment, a compound, molecule, or composition for the treatment of ocular conditions (e.g., macular degeneration, diabetic retinopathy etc.) is administered to
5 a subject intraocularly or by intraocular means. In another embodiment, a compound, molecule, or composition for the treatment of ocular conditions (e.g., macular degeneration, diabetic retinopathy etc.) is administered to a subject periorcularly or by periorcular means (see for example Ahlheim et al., International PCT publication No. WO 03/24420). In one embodiment, a siNA molecule and/or formulation or composition
10 thereof is administered to a subject intraocularly or by intraocular means. In another embodiment, a siNA molecule and/or formulation or composition thereof is administered to a subject periorcularly or by periorcular means. Periorcular administration generally provides a less invasive approach to administering siNA molecules and formulation or composition thereof to a subject (see for example Ahlheim et al., International PCT
15 publication No. WO 03/24420). The use of periorcular administration also minimizes the risk of retinal detachment, allows for more frequent dosing or administration, provides a clinically relevant route of administration for macular degeneration and other optic conditions, and also provides the possibility of using reservoirs (e.g., implants, pumps or other devices) for drug delivery.

20 In one embodiment, a siNA molecule of the invention is complexed with membrane disruptive agents such as those described in U.S. Patent Application Publication No. 20010007666, incorporated by reference herein in its entirety including the drawings. In another embodiment, the membrane disruptive agent or agents and the siNA molecule are also complexed with a cationic lipid or helper lipid molecule, such as those lipids
25 described in U.S. Patent No. 6,235,310, incorporated by reference herein in its entirety including the drawings.

In one embodiment, siNA molecules of the invention are formulated or complexed with polyethylenimine (e.g., linear or branched PEI) and/or polyethylenimine derivatives, including for example grafted PEIs such as galactose PEI, cholesterol PEI, antibody
30 derivatized PEI, and polyethylene glycol PEI (PEG-PEI) derivatives thereof (see for example Ogris *et al.*, 2001, *AAPA PharmSci*, 3, 1-11; Furgeson et al., 2003, Bioconjugate

Chem., 14, 840-847; Kunath et al., 2002, Pharmaceutical Research, 19, 810-817; Choi et al., 2001, Bull. Korean Chem. Soc., 22, 46-52; Bettinger et al., 1999, Bioconjugate Chem., 10, 558-561; Peterson et al., 2002, Bioconjugate Chem., 13, 845-854; Erbacher et al., 1999, Journal of Gene Medicine Preprint, 1, 1-18; Godbey et al., 1999, PNAS USA, 96, 5177-5181; Godbey et al., 1999, Journal of Controlled Release, 60, 149-160; Diebold et al., 1999, Journal of Biological Chemistry, 274, 19087-19094; Thomas and Klivanov, 2002, PNAS USA, 99, 14640-14645; and Sagara, US 6,586,524, incorporated by reference herein.

In one embodiment, a siNA molecule of the invention comprises a bioconjugate, for example a nucleic acid conjugate as described in Vargeese et al., USSN 10/427,160, filed April 30, 2003; US 6,528,631; US 6,335,434; US 6, 235,886; US 6,153,737; US 5,214,136; US 5,138,045, all incorporated by reference herein.

Thus, the invention features a pharmaceutical composition comprising one or more nucleic acid(s) of the invention in an acceptable carrier, such as a stabilizer, buffer, and the like. The polynucleotides of the invention can be administered (*e.g.*, RNA, DNA or protein) and introduced into a subject by any standard means, with or without stabilizers, buffers, and the like, to form a pharmaceutical composition. When it is desired to use a liposome delivery mechanism, standard protocols for formation of liposomes can be followed. The compositions of the present invention can also be formulated and used as tablets, capsules or elixirs for oral administration, suppositories for rectal administration, sterile solutions, suspensions for injectable administration, and the other compositions known in the art.

The present invention also includes pharmaceutically acceptable formulations of the compounds described. These formulations include salts of the above compounds, *e.g.*, acid addition salts, for example, salts of hydrochloric, hydrobromic, acetic acid, and benzene sulfonic acid.

A pharmacological composition or formulation refers to a composition or formulation in a form suitable for administration, *e.g.*, systemic administration, into a cell or subject, including for example a human. Suitable forms, in part, depend upon the use or the route of entry, for example oral, transdermal, or by injection. Such forms should not prevent the composition or formulation from reaching a target cell (*i.e.*, a cell to

which the negatively charged nucleic acid is desirable for delivery). For example, pharmacological compositions injected into the blood stream should be soluble. Other factors are known in the art, and include considerations such as toxicity and forms that prevent the composition or formulation from exerting its effect.

5 By "systemic administration" is meant *in vivo* systemic absorption or accumulation of drugs in the blood stream followed by distribution throughout the entire body. Administration routes that lead to systemic absorption include, without limitation: intravenous, subcutaneous, intraperitoneal, inhalation, oral, intrapulmonary and intramuscular. Each of these administration routes exposes the siNA molecules of the
 10 invention to an accessible diseased tissue. The rate of entry of a drug into the circulation has been shown to be a function of molecular weight or size. The use of a liposome or other drug carrier comprising the compounds of the instant invention can potentially localize the drug, for example, in certain tissue types, such as the tissues of the reticular endothelial system (RES). A liposome formulation that can facilitate the association of
 15 drug with the surface of cells, such as, lymphocytes and macrophages is also useful. This approach can provide enhanced delivery of the drug to target cells by taking advantage of the specificity of macrophage and lymphocyte immune recognition of abnormal cells, such as cancer cells.

By "pharmaceutically acceptable formulation" is meant a composition or
 20 formulation that allows for the effective distribution of the nucleic acid molecules of the instant invention in the physical location most suitable for their desired activity. Non-limiting examples of agents suitable for formulation with the nucleic acid molecules of the instant invention include: P-glycoprotein inhibitors (such as Pluronic P85), which can enhance entry of drugs into the CNS (Joliet-Riant and Tillement, 1999, *Fundam. Clin.*
 25 *Pharmacol.*, 13, 16-26); biodegradable polymers, such as poly (DL-lactide-coglycolide) microspheres for sustained release delivery after intracerebral implantation (Emerich, DF *et al*, 1999, *Cell Transplant*, 8, 47-58) (Alkermes, Inc. Cambridge, MA); and loaded nanoparticles, such as those made of polybutylcyanoacrylate, which can deliver drugs
 across the blood brain barrier and can alter neuronal uptake mechanisms (*Prog*
 30 *Neuropsychopharmacol Biol Psychiatry*, 23, 941-949, 1999). Other non-limiting examples of delivery strategies for the nucleic acid molecules of the instant invention include material described in Boado *et al.*, 1998, *J. Pharm. Sci.*, 87, 1308-1315; Tyler *et*

al., 1999, *FEBS Lett.*, 421, 280-284; Pardridge *et al.*, 1995, *PNAS USA.*, 92, 5592-5596; Boado, 1995, *Adv. Drug Delivery Rev.*, 15, 73-107; Aldrian-Herrada *et al.*, 1998, *Nucleic Acids Res.*, 26, 4910-4916; and Tyler *et al.*, 1999, *PNAS USA.*, 96, 7053-7058.

The invention also features the use of the composition comprising surface-modified
 5 liposomes containing poly (ethylene glycol) lipids (PEG-modified, or long-circulating liposomes or stealth liposomes). These formulations offer a method for increasing the accumulation of drugs in target tissues. This class of drug carriers resists opsonization and elimination by the mononuclear phagocytic system (MPS or RES), thereby enabling longer blood circulation times and enhanced tissue exposure for the encapsulated drug
 10 (Lasic *et al. Chem. Rev.* 1995, 95, 2601-2627; Ishiwata *et al., Chem. Pharm. Bull.* 1995, 43, 1005-1011). Such liposomes have been shown to accumulate selectively in tumors, presumably by extravasation and capture in the neovascularized target tissues (Lasic *et al., Science* 1995, 267, 1275-1276; Oku *et al.*, 1995, *Biochim. Biophys. Acta*, 1238, 86-90). The long-circulating liposomes enhance the pharmacokinetics and
 15 pharmacodynamics of DNA and RNA, particularly compared to conventional cationic liposomes which are known to accumulate in tissues of the MPS (Liu *et al., J. Biol. Chem.* 1995, 42, 24864-24870; Choi *et al.*, International PCT Publication No. WO 96/10391; Ansell *et al.*, International PCT Publication No. WO 96/10390; Holland *et al.*, International PCT Publication No. WO 96/10392). Long-circulating liposomes are also
 20 likely to protect drugs from nuclease degradation to a greater extent compared to cationic liposomes, based on their ability to avoid accumulation in metabolically aggressive MPS tissues such as the liver and spleen.

The present invention also includes compositions prepared for storage or
 administration that include a pharmaceutically effective amount of the desired compounds
 25 in a pharmaceutically acceptable carrier or diluent. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co. (A.R. Gennaro edit. 1985), hereby incorporated by reference herein. For example, preservatives, stabilizers, dyes and flavoring agents can be provided. These include sodium benzoate, sorbic acid
 30 and esters of *p*-hydroxybenzoic acid. In addition, antioxidants and suspending agents can be used.

A pharmaceutically effective dose is that dose required to prevent, inhibit the occurrence, or treat (alleviate a symptom to some extent, preferably all of the symptoms) of a disease state. The pharmaceutically effective dose depends on the type of disease, the composition used, the route of administration, the type of mammal being treated, the physical characteristics of the specific mammal under consideration, concurrent medication, and other factors that those skilled in the medical arts will recognize. Generally, an amount between 0.1 mg/kg and 100 mg/kg body weight/day of active ingredients is administered dependent upon potency of the negatively charged polymer.

The nucleic acid molecules of the invention and formulations thereof can be administered orally, topically, parenterally, by inhalation or spray, or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and/or vehicles. The term parenteral as used herein includes percutaneous, subcutaneous, intravascular (*e.g.*, intravenous), intramuscular, or intrathecal injection or infusion techniques and the like. In addition, there is provided a pharmaceutical formulation comprising a nucleic acid molecule of the invention and a pharmaceutically acceptable carrier. One or more nucleic acid molecules of the invention can be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants, and if desired other active ingredients. The pharmaceutical compositions containing nucleic acid molecules of the invention can be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use can be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions can contain one or more such sweetening agents, flavoring agents, coloring agents or preservative agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients can be, for example, inert diluents; such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets can be uncoated or they can be coated by known

techniques. In some cases such coatings can be prepared by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed.

- 5 Formulations for oral use can also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

- 10 Aqueous suspensions contain the active materials in a mixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents can be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example
15 polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with
20 partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions can also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

- 25 Oily suspensions can be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions can contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents and flavoring agents can be added to provide palatable oral preparations. These compositions can be preserved by the addition of an anti-oxidant such as ascorbic acid

- 30 Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or

wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents or suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, can also be present.

5 Pharmaceutical compositions of the invention can also be in the form of oil-in-water emulsions. The oily phase can be a vegetable oil or a mineral oil or mixtures of these. Suitable emulsifying agents can be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for
10 example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions can also contain sweetening and flavoring agents.

 Syrups and elixirs can be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol, glucose or sucrose. Such formulations can also contain a
15 demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions can be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension can be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation can also be a sterile injectable solution or suspension in a
20 non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil can be employed including synthetic mono-or diglycerides. In addition, fatty
25 acids such as oleic acid find use in the preparation of injectables.

 The nucleic acid molecules of the invention can also be administered in the form of suppositories, *e.g.*, for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the
30 rectum to release the drug. Such materials include cocoa butter and polyethylene glycols.

Nucleic acid molecules of the invention can be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

5 Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per subject per day). The amount of active ingredient that can be combined with the carrier materials to produce a single dosage form varies depending upon the host treated and the particular mode of administration. Dosage unit forms
10 generally contain between from about 1 mg to about 500 mg of an active ingredient.

It is understood that the specific dose level for any particular subject depends upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing
15 therapy.

For administration to non-human animals, the composition can also be added to the animal feed or drinking water. It can be convenient to formulate the animal feed and drinking water compositions so that the animal takes in a therapeutically appropriate quantity of the composition along with its diet. It can also be convenient to present the
20 composition as a premix for addition to the feed or drinking water.

The nucleic acid molecules of the present invention can also be administered to a subject in combination with other therapeutic compounds to increase the overall therapeutic effect. The use of multiple compounds to treat an indication can increase the beneficial effects while reducing the presence of side effects.

25 In one embodiment, the invention comprises compositions suitable for administering nucleic acid molecules of the invention to specific cell types. For example, the asialoglycoprotein receptor (ASGPr) (Wu and Wu, 1987, *J. Biol. Chem.* 262, 4429-4432) is unique to hepatocytes and binds branched galactose-terminal glycoproteins, such as asialoorosomuroid (ASOR). In another example, the folate receptor is overexpressed
30 in many cancer cells. Binding of such glycoproteins, synthetic glycoconjugates, or

folates to the receptor takes place with an affinity that strongly depends on the degree of branching of the oligosaccharide chain, for example, triantennary structures are bound with greater affinity than biantennary or monoantennary chains (Baenziger and Fiete, 1980, *Cell*, 22, 611-620; Connolly *et al.*, 1982, *J. Biol. Chem.*, 257, 939-945). Lee and Lee, 1987, *Glycoconjugate J.*, 4, 317-328, obtained this high specificity through the use of N-acetyl-D-galactosamine as the carbohydrate moiety, which has higher affinity for the receptor, compared to galactose. This "clustering effect" has also been described for the binding and uptake of mannosyl-terminating glycoproteins or glycoconjugates (Ponpipom *et al.*, 1981, *J. Med. Chem.*, 24, 1388-1395). The use of galactose, galactosamine, or folate based conjugates to transport exogenous compounds across cell membranes can provide a targeted delivery approach to, for example, the treatment of liver disease, cancers of the liver, or other cancers. The use of bioconjugates can also provide a reduction in the required dose of therapeutic compounds required for treatment. Furthermore, therapeutic bioavailability, pharmacodynamics, and pharmacokinetic parameters can be modulated through the use of nucleic acid bioconjugates of the invention. Non-limiting examples of such bioconjugates are described in Vargeese *et al.*, USSN 10/201,394, filed August 13, 2001; and Matulic-Adamic *et al.*, USSN 10/151,116, filed May 17, 2002. In one embodiment, nucleic acid molecules of the invention are complexed with or covalently attached to nanoparticles, such as Hepatitis B virus S, M, or L envelope proteins (see for example Yamado *et al.*, 2003, *Nature Biotechnology*, 21, 885). In one embodiment, nucleic acid molecules of the invention are delivered with specificity for human tumor cells, specifically non-apoptotic human tumor cells including for example T-cells, hepatocytes, breast carcinoma cells, ovarian carcinoma cells, melanoma cells, intestinal epithelial cells, prostate cells, testicular cells, non-small cell lung cancers, small cell lung cancers, etc.

Examples:

The following are non-limiting examples showing the selection, isolation, synthesis and activity of nucleic acids of the instant invention.

Example 1: Tandem synthesis of siNA constructs

Exemplary siNA molecules of the invention are synthesized in tandem using a cleavable linker, for example, a succinyl-based linker. Tandem synthesis as described

herein is followed by a one-step purification process that provides RNAi molecules in high yield. This approach is highly amenable to siNA synthesis in support of high throughput RNAi screening, and can be readily adapted to multi-column or multi-well synthesis platforms.

5 After completing a tandem synthesis of a siNA oligo and its complement in which the 5'-terminal dimethoxytrityl (5'-O-DMT) group remains intact (trityl on synthesis), the oligonucleotides are deprotected as described above. Following deprotection, the siNA sequence strands are allowed to spontaneously hybridize. This hybridization yields a duplex in which one strand has retained the 5'-O-DMT group while the complementary
10 strand comprises a terminal 5'-hydroxyl. The newly formed duplex behaves as a single molecule during routine solid-phase extraction purification (Trityl-On purification) even though only one molecule has a dimethoxytrityl group. Because the strands form a stable duplex, this dimethoxytrityl group (or an equivalent group, such as other trityl groups or other hydrophobic moieties) is all that is required to purify the pair of oligos, for example,
15 by using a C18 cartridge.

Standard phosphoramidite synthesis chemistry is used up to the point of introducing a tandem linker, such as an inverted deoxy abasic succinate or glyceryl succinate linker (see **Figure 1**) or an equivalent cleavable linker. A non-limiting example of linker coupling conditions that can be used includes a hindered base such as
20 diisopropylethylamine (DIPA) and/or DMAP in the presence of an activator reagent such as Bromotripyrrolidinophosphoniumhexafluorophosphate (PyBrOP). After the linker is coupled, standard synthesis chemistry is utilized to complete synthesis of the second sequence leaving the terminal the 5'-O-DMT intact. Following synthesis, the resulting oligonucleotide is deprotected according to the procedures described herein and quenched
25 with a suitable buffer, for example with 50mM NaOAc or 1.5M $\text{NH}_4\text{H}_2\text{CO}_3$.

Purification of the siNA duplex can be readily accomplished using solid phase extraction, for example using a Waters C18 SepPak 1g cartridge conditioned with 1 column volume (CV) of acetonitrile, 2 CV H_2O , and 2 CV 50mM NaOAc. The sample is loaded and then washed with 1 CV H_2O or 50mM NaOAc. Failure sequences are eluted
30 with 1 CV 14% ACN (Aqueous with 50mM NaOAc and 50mM NaCl). The column is then washed, for example with 1 CV H_2O followed by on-column detritylation, for

example by passing 1 CV of 1% aqueous trifluoroacetic acid (TFA) over the column, then adding a second CV of 1% aqueous TFA to the column and allowing to stand for approximately 10 minutes. The remaining TFA solution is removed and the column washed with H₂O followed by 1 CV 1M NaCl and additional H₂O. The siNA duplex product is then eluted, for example, using 1 CV 20% aqueous CAN.

Figure 2 provides an example of MALDI-TOF mass spectrometry analysis of a purified siNA construct in which each peak corresponds to the calculated mass of an individual siNA strand of the siNA duplex. The same purified siNA provides three peaks when analyzed by capillary gel electrophoresis (CGE), one peak presumably corresponding to the duplex siNA, and two peaks presumably corresponding to the separate siNA sequence strands. Ion exchange HPLC analysis of the same siNA construct only shows a single peak. Testing of the purified siNA construct using a luciferase reporter assay described below demonstrated the same RNAi activity compared to siNA constructs generated from separately synthesized oligonucleotide sequence strands.

15 Example 2: Serum stability of chemically modified siNA constructs

Chemical modifications were introduced into siNA constructs to determine the stability of these constructs compared to native siNA oligonucleotides (containing two thymidine nucleotide overhangs) in human serum. An investigation of the serum stability of RNA duplexes revealed that siNA constructs consisting of all RNA nucleotides containing two thymidine nucleotide overhangs have a half-life in serum of 15 seconds, whereas chemically modified siNA constructs remained stable in serum for 1 to 3 days depending on the extent of modification (see Figure 3). RNAi stability tests were performed by internally labeling one strand (strand 1) of siNA and duplexing with 1.5 X the concentration of the complementary siNA strand (strand 2) (to insure all labeled material was in duplex form). Duplexed siNA constructs were then tested for stability by incubating at a final concentration of 2μM siNA (strand 2 concentration) in 90% mouse or human serum for time-points of 30sec, 1min, 5min, 30min, 90min, 4hrs 10min, 16hrs 24min, and 49hrs. Time points were run on a 15% denaturing polyacrylamide gels and analyzed on a phosphoimager.

Internal labeling was performed via kinase reactions with polynucleotide kinase (PNK) and ^{32}P - γ -ATP, with addition of radiolabeled phosphate at nucleotide 13 of strand 2, counting in from the 3' side. Ligation of the remaining 8-mer fragments with T4 RNA ligase resulted in the full length, 21-mer, strand 2. Duplexing of RNAi was done by
 5 adding appropriate concentrations of the siNA oligonucleotides and heating to 95° C for 5 minutes followed by slow cooling to room temperature. Reactions were performed by adding 100% serum to the siNA duplexes and incubating at 37° C, then removing aliquots at desired time-points. Results of this study are summarized in **Figure 3**. As shown in the Figure 3, chemically modified siNA molecules (e.g., SEQ ID NOs: 412/413, 412/414,
 10 412/415, 412/416, and 412/418) have significantly increased serum stability compared to an siNA construct having all ribonucleotides except a 3'-terminal dithymidine (TT) modification (e.g., SEQ ID NOs: 419/420).

Example 3: Identification of potential siNA target sites in any RNA sequence

The sequence of an RNA target of interest, such as a viral or human mRNA transcript, is screened for target sites, for example by using a computer folding algorithm.
 15 In a non-limiting example, the sequence of a gene or RNA gene transcript derived from a database, such as Genbank, is used to generate siNA targets having complementarity to the target. Such sequences can be obtained from a database, or can be determined experimentally as known in the art. Target sites that are known, for example, those target
 20 sites determined to be effective target sites based on studies with other nucleic acid molecules, for example ribozymes or antisense, or those targets known to be associated with a disease or condition such as those sites containing mutations or deletions, can be used to design siNA molecules targeting those sites. Various parameters can be used to determine which sites are the most suitable target sites within the target RNA sequence.
 25 These parameters include but are not limited to secondary or tertiary RNA structure, the nucleotide base composition of the target sequence, the degree of homology between various regions of the target sequence, or the relative position of the target sequence within the RNA transcript. Based on these determinations, any number of target sites within the RNA transcript can be chosen to screen siNA molecules for efficacy, for
 30 example by using *in vitro* RNA cleavage assays, cell culture, or animal models. In a non-limiting example, anywhere from 1 to 1000 target sites are chosen within the transcript

based on the size of the siNA construct to be used. High throughput screening assays can be developed for screening siNA molecules using methods known in the art, such as with multi-well or multi-plate assays or combinatorial/siNA library screening assays to determine efficient reduction in target gene expression.

5 Example 4: Selection of siNA molecule target sites in a RNA

The following non-limiting steps can be used to carry out the selection of siNAs targeting a given gene sequence or transcript.

10 The target sequence is parsed in silico into a list of all fragments or subsequences of a particular length, for example 23 nucleotide fragments, contained within the target sequence. This step is typically carried out using a custom Perl script, but commercial sequence analysis programs such as Oligo, MacVector, or the GCG Wisconsin Package can be employed as well.

15 In some instances the siNAs correspond to more than one target sequence; such would be the case for example in targeting different transcripts of the same gene, targeting different transcripts of more than one gene, or for targeting both the human gene and an animal homolog. In this case, a subsequence list of a particular length is generated for each of the targets, and then the lists are compared to find matching sequences in each list. The subsequences are then ranked according to the number of target sequences that contain the given subsequence; the goal is to find subsequences that are present in most or
20 all of the target sequences. Alternately, the ranking can identify subsequences that are unique to a target sequence, such as a mutant target sequence. Such an approach would enable the use of siNA to target specifically the mutant sequence and not effect the expression of the normal sequence.

25 In some instances the siNA subsequences are absent in one or more sequences while present in the desired target sequence; such would be the case if the siNA targets a gene with a paralogous family member that is to remain untargeted. As in case 2 above, a subsequence list of a particular length is generated for each of the targets, and then the lists are compared to find sequences that are present in the target gene but are absent in the untargeted paralog.

The ranked siNA subsequences can be further analyzed and ranked according to GC content. A preference can be given to sites containing 30-70% GC, with a further preference to sites containing 40-60% GC.

5 The ranked siNA subsequences can be further analyzed and ranked according to self-folding and internal hairpins. Weaker internal folds are preferred; strong hairpin structures are to be avoided.

The ranked siNA subsequences can be further analyzed and ranked according to whether they have runs of GGG or CCC in the sequence. GGG (or even more Gs) in either strand can make oligonucleotide synthesis problematic and can potentially interfere
10 with RNAi activity, so it is avoided other appropriately suitable sequences are available. CCC is searched in the target strand because that will place GGG in the antisense strand.

The ranked siNA subsequences can be further analyzed and ranked according to whether they have the dinucleotide UU (uridine dinucleotide) on the 3'-end of the sequence, and/or AA on the 5'-end of the sequence (to yield 3' UU on the antisense
15 sequence). These sequences allow one to design siNA molecules with terminal TT thymidine dinucleotides.

Four or five target sites are chosen from the ranked list of subsequences as described above. For example, in subsequences having 23 nucleotides, the right 21 nucleotides of each chosen 23-mer subsequence are then designed and synthesized for the
20 upper (sense) strand of the siNA duplex, while the reverse complement of the left 21 nucleotides of each chosen 23-mer subsequence are then designed and synthesized for the lower (antisense) strand of the siNA duplex (see Tables I). If terminal TT residues are desired for the sequence (as described in paragraph 7), then the two 3' terminal nucleotides of both the sense and antisense strands are replaced by TT prior to
25 synthesizing the oligos.

The siNA molecules are screened in an in vitro, cell culture or animal model system to identify the most active siNA molecule or the most preferred target site within the target RNA sequence.

In an alternate approach, a pool of siNA constructs specific to a target sequence is used to screen for target sites in cells expressing target RNA, such as human HeLa cells. The general strategy used in this approach is shown in **Figure 21**. A non-limiting example of such a pool is a pool comprising sequences having antisense sequences complementary to the target RNA sequence and sense sequences complementary to the antisense sequences. Cells (e.g., HeLa cells) expressing the target gene are transfected with the pool of siNA constructs and cells that demonstrate a phenotype associated with gene silencing are sorted. The pool of siNA constructs can be chemically modified as described herein and synthesized, for example, in a high throughput manner. The siNA from cells demonstrating a positive phenotypic change (e.g., decreased target mRNA levels or target protein expression), are identified, for example by positional analysis within the assay, and are used to determine the most suitable target site(s) within the target RNA sequence based upon the complementary sequence to the corresponding siNA antisense strand identified in the assay.

15 Example 5: RNAi activity of chemically modified siNA constructs

Short interfering nucleic acid (siNA) is emerging as a powerful tool for gene regulation. All-ribose siNA duplexes activate the RNAi pathway but have limited utility as therapeutic compounds due to their nuclease sensitivity and short half-life in serum, as shown in Example 2 above. To develop nuclease-resistant siNA constructs for *in vivo* applications, siNAs that target luciferase mRNA and contain stabilizing chemical modifications were tested for activity in HeLa cells. The sequences for the siNA oligonucleotide sequences used in this study are shown in **Table I**. Modifications included phosphorothioate linkages (P=S), 2'-O-methyl nucleotides, or 2'-fluoro (F) nucleotides in one or both siNA strands and various 3'-end stabilization chemistries, including 3'-glyceryl, 3'-inverted abasic, 3'-inverted Thymidine, and/or Thymidine. The RNAi activity of chemically stabilized siNA constructs was compared with the RNAi activity of control siNA constructs consisting of all ribonucleotides at every position except the 3'-terminus which comprised two thymidine nucleotide overhangs. Active siNA molecules containing stabilizing modifications such as described herein should prove useful for *in vivo* applications, given their enhanced nuclease-resistance.

A luciferase reporter system was utilized to test RNAi activity of chemically modified siNA constructs compared to siNA constructs consisting of all RNA nucleotides containing two thymidine nucleotide overhangs. Sense and antisense siNA strands (20 uM each) were annealed by incubation in buffer (100 mM potassium acetate, 30 mM HEPES-KOH, pH 7.4, 2 mM magnesium acetate) for 1 min. at 90°C followed by 1 hour at 37°C. Plasmids encoding firefly luciferase (pGL2) and renilla luciferase (pRLSV40) were purchased from Promega Biotech.

HeLa S3 cells were grown at 37°C in DMEM with 5% FBS and seeded at 15,300 cells in 100 ul media per well of a 96-well plate 24 hours prior to transfection. For transfection, 4 ul Lipofectamine 2000 (Life Technologies) was added to 96 ul OPTI-MEM, vortexed and incubated at room temperature for 5 minutes. The 100 ul diluted lipid was then added to a microtiter tube containing 5 ul pGL2 (200ng/ul), 5 ul pRLSV40 (8 ng/ul) 6 ul siNA (25 nM or 10 nM final), and 84 ul OPTI-MEM, vortexed briefly and incubated at room temperature for 20 minutes. The transfection mix was then mixed briefly and 50 ul was added to each of three wells that contained HeLa S3 cells in 100 ul media. Cells were incubated for 20 hours after transfection and analyzed for luciferase expression using the Dual luciferase assay according to the manufacturer's instructions (Promega Biotech). The results of this study are summarized in **Figures 4-16**. The sequences of the siNA strands used in this study are shown in Table I and are referred to by Sirna/RPI # in the figures. Normalized luciferase activity is reported as the ratio of firefly luciferase activity to renilla luciferase activity in the same sample. Error bars represent standard deviation of triplicate transfections. As shown in **Figures 4-16**, the RNAi activity of chemically modified constructs is often comparable to that of unmodified control siNA constructs, which consist of all ribonucleotides at every position except the 3'-terminus which comprises two thymidine nucleotide overhangs. In some instances, the RNAi activity of the chemically modified constructs is greater than the unmodified control siNA construct consisting of all ribonucleotides..

For example, **Figure 4** shows results obtained from a screen using phosphorothioate modified siNA constructs. The Sirna/RPI 27654/27659 construct contains phosphorothioate substitutions for every pyrimidine nucleotide in both sequences, the Sirna/RPI 27657/27662 construct contains 5 terminal 3'-phosphorothioate substitutions in each strand, the Sirna/RPI 27649/27658 construct contains all phosphorothioate

substitutions only in the antisense strand, whereas the Sirna/RPI 27649/27660 and Sirna/RPI 27649/27661 constructs have unmodified sense strands and varying degrees of phosphorothioate substitutions in the antisense strand. All of these constructs show significant RNAi activity when compared to a scrambled siNA control construct (27651/27652).

Figure 5 shows results obtained from a screen using phosphorothioate (Sirna/RPI 28253/28255 and Sirna/RPI 28254/28256) and universal base substitutions (Sirna/RPI 28257/28259 and Sirna/RPI 28258/28260) compared to the same controls described above, these modifications show equivalent or better RNAi activity when compared to the unmodified control siNA construct.

Figure 6 shows results obtained from a screen using 2'-O-methyl modified siNA constructs in which the sense strand contains either 10 (Sirna/RPI 28244/27650) or 5 (Sirna/RPI 28245/27650) 2'-O-methyl substitutions, both with comparable activity to the unmodified control siNA construct.

Figure 7 shows results obtained from a screen using 2'-O-methyl or 2'-deoxy-2'-fluoro modified siNA constructs compared to a control construct consisting of all ribonucleotides at every position except the 3'-terminus which comprises two thymidine nucleotide overhangs.

Figure 8 compares a siNA construct containing six phosphorothioate substitutions in each strand (Sirna/RPI 28460/28461), where 5 phosphorothioates are present at the 3' end and a single phosphorothioate is present at the 5' end of each strand. This motif shows very similar activity to the control siNA construct consisting of all ribonucleotides at every position except the 3'-terminus, which comprises two thymidine nucleotide overhangs.

Figure 9 compares a siNA construct synthesized by the method of the invention described in Example 1, wherein an inverted deoxyabasic succinate linker was used to generate a siNA having a 3'-inverted deoxyabasic cap on the antisense strand of the siNA. This construct shows improved activity compared to the control siNA construct consisting of all ribonucleotides at every position except the 3'-terminus which comprises two thymidine nucleotide overhangs.

Figure 10 shows the results of an RNAi activity screen of chemically modified siNA constructs including 3'-glyceryl modified siNA constructs compared to an all RNA control siNA construct using a luciferase reporter system. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I. As shown in the Figure, the 3'-terminal modified siNA constructs retain significant RNAi activity compared to the unmodified control siNA (siGL2) construct.

Figure 11 shows the results of an RNAi activity screen of chemically modified siNA constructs. The screen compared various combinations of sense strand chemical modifications and antisense strand chemical modifications. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I. As shown in the figure, the chemically modified Sirna/RPI 30063/30430, Sirna/RPI 30433/30430, and Sirna/RPI 30063/30224 constructs retain significant RNAi activity compared to the unmodified control siNA construct. It should be noted that Sirna/RPI 30433/30430 is a siNA construct having no ribonucleotides which retains significant RNAi activity compared to the unmodified control siGL2 construct in vitro, therefore, this construct is expected to have both similar RNAi activity and improved stability in vivo compared to siNA constructs having ribonucleotides.

Figure 12 shows the results of an RNAi activity screen of chemically modified siNA constructs. The screen compared various combinations of sense strand chemical modifications and antisense strand chemical modifications. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM

concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number
 5 (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I. As shown in the figure, the chemically modified Sirna/RPI 30063/30224 and Sirna/RPI 30063/30430 constructs retain significant RNAi activity compared to the control siNA (siGL2) construct. In addition, the antisense strand alone (Sirna/RPI 30430) and an inverted control (Sirna/RPI 30227/30229), having matched
 10 chemistry to Sirna/RPI (30063/30224) were compared to the siNA duplexes described above. The antisense strand (Sirna/RPI 30430) alone provides far less inhibition compared to the siNA duplexes using this sequence.

Figure 13 shows the results of an RNAi activity screen of chemically modified siNA constructs. The screen compared various combinations of sense strand chemical
 15 modifications and antisense strand chemical modifications. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and
 20 antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I. In addition, an inverted control (Sirna/RPI 30226/30229, having matched chemistry to Sirna/RPI 30222/30224) was compared to the siNA duplexes described above. As shown in the figure, the chemically modified Sirna/RPI
 25 28251/30430, Sirna/RPI 28251/30224, and Sirna/RPI 30222/30224 constructs retain significant RNAi activity compared to the control siNA construct, and the chemically modified Sirna/RPI 28251/30430 construct demonstrates improved activity compared to the control siNA (siGL2) construct.

Figure 14 shows the results of an RNAi activity screen of chemically modified
 30 siNA constructs including various 3'-terminal modified siNA constructs compared to an all RNA control siNA construct using a luciferase reporter system. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and

10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by
 5 Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table 1. As shown in the figure, the chemically modified Sirna/RPI 30222/30546, 30222/30224, 30222/30551, 30222/30557 and 30222/30558 constructs retain significant RNAi activity compared to the control siNA construct.

10 **Figure 15** shows the results of an RNAi activity screen of chemically modified siNA constructs. The screen compared various combinations of sense strand chemistries compared to a fixed antisense strand chemistry. These chemically modified siNAs were compared in the luciferase assay described herein at 1 nM and 10nM concentration using an all RNA siNA control (siGL2) having 3'-terminal dithymidine (TT) and its
 15 corresponding inverted control (Inv siGL2). The background level of luciferase expression in the HeLa cells is designated by the "cells" column. Sense and antisense strands of chemically modified siNA constructs are shown by Sirna/RPI number (sense strand/antisense strand). Sequences corresponding to these Sirna/RPI numbers are shown in Table I. As shown in the figure, the chemically modified Sirna/RPI 30063/30430,
 20 30434/30430, and 30435/30430 constructs all demonstrate greater activity compared to the control siNA (siGL2) construct.

Example 6: RNAi activity titration

A titration assay was performed to determine the lower range of siNA concentration required for RNAi activity both in a control siNA construct consisting of all RNA
 25 nucleotides containing two thymidine nucleotide overhangs and a chemically modified siNA construct comprising five phosphorothioate internucleotide linkages in both the sense and antisense strands. The assay was performed as described above, however, the siNA constructs were diluted to final concentrations between 2.5 nM and 0.025 nM. Results are shown in **Figure 16**. As shown in **Figure 16**, the chemically modified siNA
 30 construct shows a very similar concentration dependent RNAi activity profile to the control siNA construct when compared to an inverted siNA sequence control.

Example 7: siNA design

siNA target sites were chosen by analyzing sequences of the target RNA and optionally prioritizing the target sites on the basis of folding (structure of any given sequence analyzed to determine siNA accessibility to the target), by using a library of siNA molecules as described in Example 4, or alternately by using an *in vitro* siNA system as described in Example 9 herein. siNA molecules were designed that could bind each target and are optionally individually analyzed by computer folding to assess whether the siNA molecule can interact with the target sequence. Varying the length of the siNA molecules can be chosen to optimize activity. Generally, a sufficient number of complementary nucleotide bases are chosen to bind to, or otherwise interact with, the target RNA, but the degree of complementarity can be modulated to accommodate siNA duplexes or varying length or base composition. By using such methodologies, siNA molecules can be designed to target sites within any known RNA sequence, for example those RNA sequences corresponding to the any gene transcript.

Chemically modified siNA constructs are designed to provide nuclease stability for systemic administration *in vivo* and/or improved pharmacokinetic, localization, and delivery properties while preserving the ability to mediate RNAi activity. Chemical modifications as described herein are introduced synthetically using synthetic methods described herein and those generally known in the art. The synthetic siNA constructs are then assayed for nuclease stability in serum and/or cellular/tissue extracts (e.g. liver extracts). The synthetic siNA constructs are also tested in parallel for RNAi activity using an appropriate assay, such as a luciferase reporter assay as described herein or another suitable assay that can quantify RNAi activity. Synthetic siNA constructs that possess both nuclease stability and RNAi activity can be further modified and re-evaluated in stability and activity assays. The chemical modifications of the stabilized active siNA constructs can then be applied to any siNA sequence targeting any chosen RNA and used, for example, in target screening assays to pick lead siNA compounds for therapeutic development (see for example **Figure 27**).

Example 8: Chemical Synthesis and Purification of siNA

siNA molecules can be designed to interact with various sites in the RNA message, for example, target sequences within the RNA sequences described herein. The sequence

of one strand of the siNA molecule(s) is complementary to the target site sequences described above. The siNA molecules can be chemically synthesized using methods described herein. Inactive siNA molecules that are used as control sequences can be synthesized by scrambling the sequence of the siNA molecules such that it is not complementary to the target sequence. Generally, siNA constructs can be synthesized using solid phase oligonucleotide synthesis methods as described herein (see for example Usman *et al.*, US Patent Nos. 5,804,683; 5,831,071; 5,998,203; 6,117,657; 6,353,098; 6,362,323; 6,437,117; 6,469,158; Scaringe *et al.*, US Patent Nos. 6,111,086; 6,008,400; 6,111,086 all incorporated by reference herein in their entirety).

In a non-limiting example, RNA oligonucleotides are synthesized in a stepwise fashion using the phosphoramidite chemistry as is known in the art. Standard phosphoramidite chemistry involves the use of nucleosides comprising any of 5'-O-dimethoxytrityl, 2'-O-tert-butyldimethylsilyl, 3'-O-2-Cyanoethyl N,N-diisopropylphosphoroamidite groups, and exocyclic amine protecting groups (e.g. N6-benzoyl adenosine, N4 acetyl cytidine, and N2-isobutyryl guanosine). Alternately, 2'-O-Silyl Ethers can be used in conjunction with acid-labile 2'-O-orthoester protecting groups in the synthesis of RNA as described by Scaringe *supra*. Differing 2' chemistries can require different protecting groups, for example 2'-deoxy-2'-amino nucleosides can utilize N-phthaloyl protection as described by Usman *et al.*, US Patent 5,631,360, incorporated by reference herein in its entirety).

During solid phase synthesis, each nucleotide is added sequentially (3'- to 5'-direction) to the solid support-bound oligonucleotide. The first nucleoside at the 3'-end of the chain is covalently attached to a solid support (e.g., controlled pore glass or polystyrene) using various linkers. The nucleotide precursor, a ribonucleoside phosphoramidite, and activator are combined resulting in the coupling of the second nucleoside phosphoramidite onto the 5'-end of the first nucleoside. The support is then washed and any unreacted 5'-hydroxyl groups are capped with a capping reagent such as acetic anhydride to yield inactive 5'-acetyl moieties. The trivalent phosphorus linkage is then oxidized to a more stable phosphate linkage. At the end of the nucleotide addition cycle, the 5'-O-protecting group is cleaved under suitable conditions (e.g., acidic conditions for trityl-based groups and Fluoride for silyl-based groups). The cycle is repeated for each subsequent nucleotide.

Modification of synthesis conditions can be used to optimize coupling efficiency, for example by using differing coupling times, differing reagent/phosphoramidite concentrations, differing contact times, differing solid supports and solid support linker chemistries depending on the particular chemical composition of the siNA to be synthesized. Deprotection and purification of the siNA can be performed as is generally described in Deprotection and purification of the siNA can be performed as is generally described in Usman *et al.*, US 5,831,071, US 6,353,098, US 6,437,117, and Bellon *et al.*, US 6,054,576, US 6,162,909, US 6,303,773, or Scaringe *supra*, incorporated by reference herein in their entireties. Additionally, deprotection conditions can be modified to provide the best possible yield and purity of siNA constructs. For example, applicant has observed that oligonucleotides comprising 2'-deoxy-2'-fluoro nucleotides can degrade under inappropriate deprotection conditions. Such oligonucleotides are deprotected using aqueous methylamine at about 35°C for 30 minutes. If the 2'-deoxy-2'-fluoro containing oligonucleotide also comprises ribonucleotides, after deprotection with aqueous methylamine at about 35°C for 30 minutes, TEA-HF is added and the reaction maintained at about 65°C for an additional 15 minutes.

Example 9: RNAi *in vitro* assay to assess siNA activity

An *in vitro* assay that recapitulates RNAi in a cell free system is used to evaluate siNA constructs specific to target RNA. The assay comprises the system described by Tuschl *et al.*, 1999, *Genes and Development*, 13, 3191-3197 and Zamore *et al.*, 2000, *Cell*, 101, 25-33 adapted for use with target RNA. A *Drosophila* extract derived from syncytial blastoderm is used to reconstitute RNAi activity *in vitro*. Target RNA is generated via *in vitro* transcription from an appropriate plasmid using T7 RNA polymerase or via chemical synthesis as described herein. Sense and antisense siNA strands (for example 20 uM each) are annealed by incubation in buffer (such as 100 mM potassium acetate, 30 mM HEPES-KOH, pH 7.4, 2 mM magnesium acetate) for 1 minute at 90°C followed by 1 hour at 37°C, then diluted in lysis buffer (for example 100 mM potassium acetate, 30 mM HEPES-KOH at pH 7.4, 2mM magnesium acetate). Annealing can be monitored by gel electrophoresis on an agarose gel in TBE buffer and stained with ethidium bromide. The *Drosophila* lysate is prepared using zero to two-hour-old embryos from Oregon R flies collected on yeasted molasses agar that are dechorionated and lysed. The lysate is centrifuged and the supernatant isolated. The assay comprises a reaction

mixture containing 50% lysate [vol/vol], RNA (10-50 pM final concentration), and 10% [vol/vol] lysis buffer containing siNA (10 nM final concentration). The reaction mixture also contains 10 mM creatine phosphate, 10 ug/ml creatine phosphokinase, 100 μ M GTP, 100 μ M UTP, 100 μ M CTP, 500 μ M ATP, 5 mM DTT, 0.1 U/ μ L RNasin (Promega), and 100 μ M of each amino acid. The final concentration of potassium acetate is adjusted to 100 mM. The reactions are pre-assembled on ice and preincubated at 25° C for 10 minutes before adding RNA, then incubated at 25° C for an additional 60 minutes. Reactions are quenched with 4 volumes of 1.25 x Passive Lysis Buffer (Promega). Target RNA cleavage is assayed by RT-PCR analysis or other methods known in the art and are compared to control reactions in which siNA is omitted from the reaction.

Alternately, internally-labeled target RNA for the assay is prepared by *in vitro* transcription in the presence of [α -³²P] CTP, passed over a G 50 Sephadex column by spin chromatography and used as target RNA without further purification. Optionally, target RNA is 5'-³²P-end labeled using T4 polynucleotide kinase enzyme. Assays are performed as described above and target RNA and the specific RNA cleavage products generated by RNAi are visualized on an autoradiograph of a gel. The percentage of cleavage is determined by Phosphor Imager[®] quantitation of bands representing intact control RNA or RNA from control reactions without siNA and the cleavage products generated by the assay.

In one embodiment, this assay is used to determine target sites the RNA target for siNA mediated RNAi cleavage, wherein a plurality of siNA constructs are screened for RNAi mediated cleavage of the RNA target, for example, by analyzing the assay reaction by electrophoresis of labeled target RNA, or by northern blotting, as well as by other methodology well known in the art.

25 Example 10: Nucleic acid inhibition of target RNA *in vivo*

siNA molecules targeted to the target RNA are designed and synthesized as described above. These nucleic acid molecules can be tested for cleavage activity *in vivo*, for example, using the following procedure.

Two formats are used to test the efficacy of siNAs targeting a particular gene transcript. First, the reagents are tested on target expressing cells (e.g., HeLa), to determine the extent of RNA and protein inhibition. siNA reagents are selected against the RNA target. RNA inhibition is measured after delivery of these reagents by a suitable
 5 transfection agent to cells. Relative amounts of target RNA are measured versus actin using real-time PCR monitoring of amplification (*eg.*, ABI 7700 Taqman®). A comparison is made to a mixture of oligonucleotide sequences made to unrelated targets or to a randomized siNA control with the same overall length and chemistry, but with randomly substituted nucleotides at each position. Primary and secondary lead reagents
 10 are chosen for the target and optimization performed. After an optimal transfection agent concentration is chosen, a RNA time-course of inhibition is performed with the lead siNA molecule. In addition, a cell-plating format can be used to determine RNA inhibition.

Delivery of siNA to Cells

Cells (e.g., HeLa) are seeded, for example, at 1×10^5 cells per well of a six-well dish
 15 in EGM-2 (BioWhittaker) the day before transfection. siNA (final concentration, for example 20nM) and cationic lipid (e.g., final concentration $2 \mu\text{g/ml}$) are complexed in EGM basal media (Biowhittaker) at 37°C for 30 mins in polystyrene tubes. Following vortexing, the complexed siNA is added to each well and incubated for the times indicated. For initial optimization experiments, cells are seeded, for example, at 1×10^3 in
 20 96 well plates and siNA complex added as described. Efficiency of delivery of siNA to cells is determined using a fluorescent siNA complexed with lipid. Cells in 6-well dishes are incubated with siNA for 24 hours, rinsed with PBS and fixed in 2% paraformaldehyde for 15 minutes at room temperature. Uptake of siNA is visualized using a fluorescent microscope.

25 Taqman and Lightcycler quantification of mRNA

Total RNA is prepared from cells following siNA delivery, for example, using Qiagen RNA purification kits for 6-well or Rneasy extraction kits for 96-well assays. For Taqman analysis, dual-labeled probes are synthesized with the reporter dye, FAM or JOE, covalently linked at the 5'-end and the quencher dye TAMRA conjugated to the 3'-end.
 30 One-step RT-PCR amplifications are performed on, for example, an ABI PRISM 7700

Sequence Detector using 50 μ l reactions consisting of 10 μ l total RNA, 100 nM forward primer, 900 nM reverse primer, 100 nM probe, 1X TaqMan PCR reaction buffer (PE-Applied Biosystems), 5.5 mM MgCl₂, 300 μ M each dATP, dCTP, dGTP, and dTTP, 10U RNase Inhibitor (Promega), 1.25U AmpliTaq Gold (PE-Applied Biosystems) and 10U M-
 5 MLV Reverse Transcriptase (Promega). The thermal cycling conditions can consist of 30 min at 48°C, 10 min at 95°C, followed by 40 cycles of 15 sec at 95°C and 1 min at 60°C. Quantitation of mRNA levels is determined relative to standards generated from serially diluted total cellular RNA (300, 100, 33, 11 ng/rxn) and normalizing to β -actin or GAPDH mRNA in parallel TaqMan reactions. For each gene of interest an upper and
 10 lower primer and a fluorescently labeled probe are designed. Real time incorporation of SYBR Green I dye into a specific PCR product can be measured in glass capillary tubes using a lightcycler. A standard curve is generated for each primer pair using control cRNA. Values are represented as relative expression to GAPDH in each sample.

Western blotting

15 Nuclear extracts can be prepared using a standard micro preparation technique (see for example Andrews and Faller, 1991, *Nucleic Acids Research*, 19, 2499). Protein extracts from supernatants are prepared, for example using TCA precipitation. An equal volume of 20% TCA is added to the cell supernatant, incubated on ice for 1 hour and pelleted by centrifugation for 5 minutes. Pellets are washed in acetone, dried and
 20 resuspended in water. Cellular protein extracts are run on a 10% Bis-Tris NuPage (nuclear extracts) or 4-12% Tris-Glycine (supernatant extracts) polyacrylamide gel and transferred onto nitro-cellulose membranes. Non-specific binding can be blocked by incubation, for example, with 5% non-fat milk for 1 hour followed by primary antibody for 16 hour at 4°C. Following washes, the secondary antibody is applied, for example
 25 (1:10,000 dilution) for 1 hour at room temperature and the signal detected with SuperSignal reagent (Pierce).

Example 11: Animal Models

Various animal models can be used to screen siNA constructs *in vivo* as are known in the art, for example those animal models that are used to evaluate other nucleic acid
 30 technologies such as enzymatic nucleic acid molecules (ribozymes) and/or antisense. Such animal models are used to test the efficacy of siNA molecules described herein. In

a non-limiting example, siNA molecules that are designed as anti-angiogenic agents can be screened using animal models. There are several animal models available in which to test the anti-angiogenesis effect of nucleic acids of the present invention, such as siNA, directed against genes associated with angiogenesis and/or metastasis, such as VEGFR (e.g., VEGFR1, VEGFR2, and VEGFR3) genes. Typically a corneal model has been used to study angiogenesis in rat and rabbit, since recruitment of vessels can easily be followed in this normally avascular tissue (Pandey *et al.*, 1995 *Science* 268: 567-569). In these models, a small Teflon or Hydrion disk pretreated with an angiogenesis factor (e.g. bFGF or VEGF) is inserted into a pocket surgically created in the cornea. Angiogenesis is monitored 3 to 5 days later. siNA molecules directed against VEGFR mRNAs would be delivered in the disk as well, or dropwise to the eye over the time course of the experiment. In another eye model, hypoxia has been shown to cause both increased expression of VEGF and neovascularization in the retina (Pierce *et al.*, 1995 *Proc. Natl. Acad. Sci. USA.* 92: 905-909; Shweiki *et al.*, 1992 *J. Clin. Invest.* 91: 2235-2243).

Several animal models exist for screening of anti-angiogenic agents. These include corneal vessel formation following corneal injury (Burger *et al.*, 1985 *Cornea* 4: 35-41; Lepri, *et al.*, 1994 *J. Ocular Pharmacol.* 10: 273-280; Ormerod *et al.*, 1990 *Am. J. Pathol.* 137: 1243-1252) or intracorneal growth factor implant (Grant *et al.*, 1993 *Diabetologia* 36: 282-291; Pandey *et al.* 1995 *supra*; Zieche *et al.*, 1992 *Lab. Invest.* 67: 711-715), vessel growth into Matrigel matrix containing growth factors (Passaniti *et al.*, 1992 *supra*), female reproductive organ neovascularization following hormonal manipulation (Shweiki *et al.*, 1993 *Clin. Invest.* 91: 2235-2243), several models involving inhibition of tumor growth in highly vascularized solid tumors (O'Reilly *et al.*, 1994 *Cell* 79: 315-328; Senger *et al.*, 1993 *Cancer and Metas. Rev.* 12: 303-324; Takahashi *et al.*, 1994 *Cancer Res.* 54: 4233-4237; Kim *et al.*, 1993 *supra*), and transient hypoxia-induced neovascularization in the mouse retina (Pierce *et al.*, 1995 *Proc. Natl. Acad. Sci. USA.* 92: 905-909).gene

The cornea model, described in Pandey *et al. supra*, is the most common and well characterized anti-angiogenic agent efficacy screening model. This model involves an avascular tissue into which vessels are recruited by a stimulating agent (growth factor, thermal or alkalai burn, endotoxin). The corneal model utilizes the intrastromal corneal implantation of a Teflon pellet soaked in a VEGF-Hydrion solution to recruit blood

vessels toward the pellet, which can be quantitated using standard microscopic and image analysis techniques. To evaluate their anti-angiogenic efficacy, siNA molecules are applied topically to the eye or bound within Hydron on the Teflon pellet itself. This avascular cornea as well as the Matrigel model (described below) provide for low background assays. While the corneal model has been performed extensively in the rabbit, studies in the rat have also been conducted.

The mouse model (Passaniti et al., *supra*) is a non-tissue model which utilizes Matrigel, an extract of basement membrane (Kleinman et al., 1986) or Millipore® filter disk, which can be impregnated with growth factors and anti-angiogenic agents in a liquid form prior to injection. Upon subcutaneous administration at body temperature, the Matrigel or Millipore® filter disk forms a solid implant. VEGF embedded in the Matrigel or Millipore® filter disk is used to recruit vessels within the matrix of the Matrigel or Millipore® filter disk which can be processed histologically for endothelial cell specific vWF (factor VIII antigen) immunohistochemistry, Trichrome-Masson stain, or hemoglobin content. Like the cornea, the Matrigel or Millipore® filter disk are avascular; however, it is not tissue. In the Matrigel or Millipore® filter disk model, siNA molecules are administered within the matrix of the Matrigel or Millipore® filter disk to test their anti-angiogenic efficacy. Thus, delivery issues in this model, as with delivery of siNA molecules by Hydron- coated Teflon pellets in the rat cornea model, may be less problematic due to the homogeneous presence of the siNA within the respective matrix.

The Lewis lung carcinoma and B-16 murine melanoma models are well accepted models of primary and metastatic cancer and are used for initial screening of anti-cancer agents. These murine models are not dependent upon the use of immunodeficient mice, are relatively inexpensive, and minimize housing concerns. Both the Lewis lung and B-16 melanoma models involve subcutaneous implantation of approximately 10^6 tumor cells from metastatically aggressive tumor cell lines (Lewis lung lines 3LL or D122, LLC-LN7; B-16-BL6 melanoma) in C57BL/6J mice. Alternatively, the Lewis lung model can be produced by the surgical implantation of tumor spheres (approximately 0.8 mm in diameter). Metastasis also may be modeled by injecting the tumor cells directly intravenously. In the Lewis lung model, microscopic metastases can be observed approximately 14 days following implantation with quantifiable macroscopic metastatic

tumors developing within 21-25 days. The B-16 melanoma exhibits a similar time course with tumor neovascularization beginning 4 days following implantation. Since both primary and metastatic tumors exist in these models after 21-25 days in the same animal, multiple measurements can be taken as indices of efficacy. Primary tumor volume and growth latency as well as the number of micro- and macroscopic metastatic lung foci or number of animals exhibiting metastases can be quantitated. The percent increase in lifespan can also be measured. Thus, these models would provide suitable primary efficacy assays for screening systemically administered siNA molecules and siNA formulations.

In the Lewis lung and B-16 melanoma models, systemic pharmacotherapy with a wide variety of agents usually begins 1-7 days following tumor implantation/inoculation with either continuous or multiple administration regimens. Concurrent pharmacokinetic studies can be performed to determine whether sufficient tissue levels of siNA can be achieved for pharmacodynamic effect to be expected. Furthermore, primary tumors and secondary lung metastases can be removed and subjected to a variety of *in vitro* studies (*i.e.* target RNA reduction).

Ohno-Matsui *et al.*, 2002, *Am. J. Pathology*, 160, 711-719 describe a model of severe proliferative retinopathy and retinal detachment in mice under inducible expression of vascular endothelial growth factor. In this model, expression of a VEGF transgene results in elevated levels of ocular VEGF that is associated with severe proliferative retinopathy and retinal detachment. Furthermore, Mori *et al.*, 2001, *J. Cellular Physiology*, 188, 253-263, describe a model of laser induced choroidal neovascularization that can be used in conjunction with intravitreal or subretinal injection of siNA molecules of the invention to evaluate the efficacy of siNA treatment of severe proliferative retinopathy and retinal detachment.

In utilizing these models to assess siNA activity, VEGFR1, VEGFR2, and/or VEGFR3 protein levels can be measured clinically or experimentally by FACS analysis. VEGFR1, VEGFR2, and/or VEGFR3 encoded mRNA levels can be assessed by Northern analysis, RNase-protection, primer extension analysis and/or quantitative RT-PCR. siNA molecules that block VEGFR1, VEGFR2, and/or VEGFR3 protein encoding mRNAs and therefore result in decreased levels of VEGFR1, VEGFR2, and/or VEGFR3

activity by more than 20% *in vitro* can be identified using the techniques described herein.

Example 12: siNA-mediated inhibition of angiogenesis *in vivo*

The purpose of this study was to assess the anti-angiogenic activity of siNA
5 targeted against VEGFR1, using the rat cornea model of VEGF induced angiogenesis
discussed in Example 11 above). The siNA molecules shown in **Figure 23** have matched
inverted controls which are inactive since they are not able to interact with the RNA
target. The siNA molecules and VEGF were co-delivered using the filter disk method.
Nitrocellulose filter disks (Millipore®) of 0.057 diameter were immersed in appropriate
10 solutions and were surgically implanted in rat cornea as described by Pandey *et al., supra*.

The stimulus for angiogenesis in this study was the treatment of the filter disk with
30 μ M VEGF which is implanted within the cornea's stroma. This dose yields
reproducible neovascularization stemming from the pericorneal vascular plexus growing
toward the disk in a dose-response study 5 days following implant. Filter disks treated
15 only with the vehicle for VEGF show no angiogenic response. The siNA were co-
administered with VEGF on a disk in three different siNA concentrations. One concern
with the simultaneous administration is that the siNA would not be able to inhibit
angiogenesis since VEGF receptors can be stimulated. However, Applicant has observed
that in low VEGF doses, the neovascular response reverts to normal suggesting that the
20 VEGF stimulus is essential for maintaining the angiogenic response. Blocking the
production of VEGF receptors using simultaneous administration of anti-VEGF-R mRNA
siNA could attenuate the normal neovascularization induced by the filter disk treated with
VEGF.

Materials and Methods:

25 *Test Compounds and Controls*

R&D Systems VEGF, carrier free at 75 μ M in 82 mM Tris-Cl, pH 6.9

siNA, 1.67 $\mu\text{G}/\mu\text{L}$, SITE 2340 (SIRNA/RPI 29695/29699) sense/antisense

siNA, 1.67 $\mu\text{G}/\mu\text{L}$, INVERTED CONTROL FOR SITE 2340 (SIRNA/RPI 29983/29984) sense/antisense

siNA 1.67 $\mu\text{g}/\mu\text{L}$, Site 2340 (Sirna/RPI 30196/30416) sense/antisense

5 *Animals*

Harlan Sprague-Dawley Rats, Approximately 225-250g

45 males, 5 animals per group.

Husbandry

10

Animals are housed in groups of two. Feed, water, temperature and humidity are determined according to Pharmacology Testing Facility performance standards (SOP's) which are in accordance with the 1996 Guide for the Care and Use of Laboratory Animals (NRC). Animals are acclimated to the facility for at least 7 days prior to experimentation.

15 During this time, animals are observed for overall health and sentinels are bled for baseline serology.

Experimental Groups

Each solution (VEGF and siNAs) was prepared as a 1X solution for final
20 concentrations shown in the experimental groups described in **Table III**.

siNA Annealing Conditions

25 siNA sense and antisense strands are annealed for 1 minute in H_2O at 1.67mg/mL/strand followed by a 1 hour incubation at 37°C producing 3.34 mg/mL of duplexed siNA. For the 20 $\mu\text{g}/\text{eye}$ treatment, 6 μL s of the 3.34 mg/mL duplex is injected into the eye (see below). The 3.34 mg/mL duplex siNA can then be serially diluted for dose response assays.

Preparation of VEGF Filter Disk

For corneal implantation, 0.57 mm diameter nitrocellulose disks, prepared from 0.45 μm pore diameter nitrocellulose filter membranes (Millipore Corporation), were soaked for 30 min in 1 μL of 75 μM VEGF in 82 mM Tris-HCl (pH 6.9) in covered petri dishes on ice. Filter disks soaked only with the vehicle for VEGF (83 mM Tris-Cl pH 6.9) elicit no angiogenic response.

Corneal surgery

The rat corneal model used in this study was a modified from Koch *et al. Supra* and Pandey *et al., supra*. Briefly, corneas were irrigated with 0.5% povidone iodine solution followed by normal saline and two drops of 2% lidocaine. Under a dissecting microscope (Leica MZ-6), a stromal pocket was created and a presoaked filter disk (see above) was inserted into the pocket such that its edge was 1 mm from the corneal limbus.

Intraconjunctival injection of test solutions

Immediately after disk insertion, the tip of a 40-50 μm OD injector (constructed in our laboratory) was inserted within the conjunctival tissue 1 mm away from the edge of the corneal limbus that was directly adjacent to the VEGF-soaked filter disk. Six hundred nanoliters of test solution (siNA, inverted control or sterile water vehicle) were dispensed at a rate of 1.2 $\mu\text{L}/\text{min}$ using a syringe pump (Kd Scientific). The injector was then removed, serially rinsed in 70% ethanol and sterile water and immersed in sterile water between each injection. Once the test solution was injected, closure of the eyelid was maintained using microaneurism clips until the animal began to recover gross motor activity. Following treatment, animals were warmed on a heating pad at 37°C.

Quantitation of angiogenic response

Five days after disk implantation, animals were euthanized following administration of 0.4 mg/kg atropine and corneas were digitally imaged. The neovascular surface area (NSA, expressed in pixels) was measured *postmortem* from blood-filled corneal vessels using computerized morphometry (Image Pro Plus, Media Cybernetics, v2.0). The individual mean NSA was determined in triplicate from three regions of

identical size in the area of maximal neovascularization between the filter disk and the limbus. The number of pixels corresponding to the blood-filled corneal vessels in these regions was summated to produce an index of NSA. A group mean NSA was then calculated. Data from each treatment group were normalized to VEGF/siNA vehicle-treated control NSA and finally expressed as percent inhibition of VEGF-induced angiogenesis.

Statistics

After determining the normality of treatment group means, group mean percent inhibition of VEGF-induced angiogenesis was subjected to a one-way analysis of variance. This was followed by two post-hoc tests for significance including Dunnett's (comparison to VEGF control) and Tukey-Kramer (all other group mean comparisons) at $\alpha = 0.05$. Statistical analyses were performed using JMP v.3.1.6 (SAS Institute).

Results of the study are graphically represented in **Figures 23 and 76**. As shown in **Figure 23**, VEGFr1 site 4229 active siNA (Sirna/RPI 29695/29699) at three concentrations were effective at inhibiting angiogenesis compared to the inverted siNA control (Sirna/RPI 29983/29984) and the VEGF control. A chemically modified version of the VEGFr1 site 4229 active siNA comprising a sense strand having 2'-deoxy-2'-fluoro pyrimidines and ribo purines with 5' and 3' terminal inverted deoxyabasic residues and an antisense strand having having 2'-deoxy-2'-fluoro pyrimidines and ribo purines with a terminal 3'-phosphorothioate internucleotide linkage (Sirna/RPI 30196/30416), showed similar inhibition. Furthermore, VEGFr1 site 349 active siNA having "Stab 9/10" chemistry (Sirna # 31270/31273) was tested for inhibition of VEGF-induced angiogenesis at three different concentrations (2.0 ug, 1.0 ug, and 0.1 μ g dose response) as compared to a matched chemistry inverted control siNA construct (Sirna # 31276/31279) at each concentration and a VEGF control in which no siNA was administered. As shown in **Figure 76**, the active siNA construct having "Stab 9/10" chemistry (Sirna # 31270/31273) is highly effective in inhibiting VEGF-induced angiogenesis in the rat corneal model compared to the matched chemistry inverted control siNA at concentrations from 0.1 μ g to 2.0 ug. These results demonstrate that siNA molecules having different chemically modified compositions, such as the modifications described herein, are capable of significantly inhibiting angiogenesis *in vivo*.

Example 13: Inhibition of HBV using siNA Molecules of the Invention

Transfection of HepG2 Cells with psHBV-1 and siNA

The human hepatocellular carcinoma cell line Hep G2 was grown in Dulbecco's modified Eagle media supplemented with 10% fetal calf serum, 2 mM glutamine, 0.1 mM nonessential amino acids, 1 mM sodium pyruvate, 25 mM Hepes, 100 units penicillin, and 100 µg/ml streptomycin. To generate a replication competent cDNA, prior to transfection the HBV genomic sequences are excised from the bacterial plasmid sequence contained in the psHBV-1 vector. Other methods known in the art can be used to generate a replication competent cDNA. This was done with an EcoRI and Hind III restriction digest. Following completion of the digest, a ligation was performed under dilute conditions (20 µg/ml) to favor intermolecular ligation. The total ligation mixture was then concentrated using Qiagen spin columns.

siNA Activity Screen and Dose Response Assay

Transfection of the human hepatocellular carcinoma cell line, Hep G2, with replication-competent HBV DNA results in the expression of HBV proteins and the production of virions. To test the efficacy of siNAs targeted against HBV RNA, several siNA duplexes targeting different sites within HBV pregenomic RNA were co-transfected with HBV genomic DNA once at 25 nM with lipid at 12.5 µg/ml into Hep G2 cells, and the subsequent levels of secreted HBV surface antigen (HBsAg) were analyzed by ELISA (see **Figure 24**). Inverted sequence duplexes were used as negative controls. Subsequently, dose response studies were performed in which the siNA duplexes were co-transfected with HBV genomic DNA at 0.5, 5, 10 and 25 nM with lipid at 12.5 µg/ml into Hep G2 cells, and the subsequent levels of secreted HBV surface antigen (HBsAg) were analyzed by ELISA (see **Figure 25**).

Analysis of HBsAg Levels Following siNA Treatment

To determine siNA activity, HbsAg levels were measured following transfection with siNA. Immulon 4 (Dynax) microtiter wells were coated overnight at 4° C with anti-HBsAg Mab (Biostride B88-95-31ad,ay) at 1 µg/ml in Carbonate Buffer (Na₂CO₃ 15 mM, NaHCO₃ 35 mM, pH 9.5). The wells were then washed 4x with PBST (PBS, 0.05%

Tween® 20) and blocked for 1 hr at 37° C with PBST, 1% BSA. Following washing as above, the wells were dried at 37° C for 30 min. Biotinylated goat anti-HBsAg (Accurate YVS1807) was diluted 1:1000 in PBST and incubated in the wells for 1 hr. at 37° C. The wells were washed 4x with PBST. Streptavidin/Alkaline Phosphatase Conjugate (Pierce 21324) was diluted to 250 ng/ml in PBST, and incubated in the wells for 1 hr. at 37° C. After washing as above, p-nitrophenyl phosphate substrate (Pierce 37620) was added to the wells, which were then incubated for 1 hour at 37° C. The optical density at 405 nm was then determined. Results of the HBV screen study are summarized in **Figure 24**, whereas the results of a dose response assay using lead siNA constructs targeting sites 262 and 1580 of the HBV pregenomic RNA are shown in **Figure 25**. As shown in **Figure 25**, the siNA constructs targeting sites 262 and 1580 of HBV RNA provides significant dose response inhibition of viral replication/activity when compared to inverted siNA controls.

Comparison of different chemically stabilized siNA motifs targeting HBV RNA site 1580

Two different siNA stabilization chemistries were compared in a dose response HBsAg assay using inverted matched chemistry controls. The “Stab7/8” (**Table IV**) constructs comprise a sense strand having 2'-deoxy-2'-fluoro pyrimidine nucleotides and 2'-deoxy purine nucleotides with 5' and 3' terminal inverted deoxyabasic residues and an antisense strand having 2'-deoxy-2'-fluoro pyrimidine nucleotides and 2'-O-methyl purine nucleotides with a terminal 3' phosphorothioate linkage. The “Stab7/11” (**Table IV**) constructs comprise a sense strand having 2'-deoxy-2'-fluoro pyrimidine nucleotides and 2'-deoxy purine nucleotides with 5' and 3' terminal inverted deoxyabasic residues and an antisense strand having 2'-deoxy-2'-fluoro pyrimidine nucleotides and 2'-deoxy purine nucleotides with a terminal 3' phosphorothioate linkage (see for example **Table I**). As shown in **Figure 26**, the chemically stabilized siNA constructs both show significant inhibition of HBV antigen in a dose dependent manner compared to matched inverted controls. A separate direct screen of Stab 7/8 constructs targeting HBV RNA in HepG2 cells that identified stabilized siNA constructs with potent activity is shown in **Figure 87**.

Time course evaluation of different chemically stabilized siNA motifs targeting HBV RNA site 1580

Four different siNA constructs having different stabilization chemistries were compared to an unstabilized siRNA construct in a dose response time course HBsAg assay, the results of which are shown in **Figures 28-31**. The different constructs were compared to an unstabilized ribonucleotide control siRNA construct (Sirna/RPI#30287/30298) at different concentrations (5nM, 10 nM, 25 nM, 50 nM, and 100 nM) over the course of nine days. Activity based on HBsAg levels was determined at day 3, day 6, and day 9. The “Stab 4/5” (**Table IV**) constructs comprise a sense strand (Sirna/RPI#30355) having 2'-deoxy-2'-fluoro pyrimidine nucleotides and purine ribonucleotides with 5' and 3' terminal inverted deoxyabasic residues and an antisense strand (Sirna/RPI#30366) having 2'-deoxy-2'-fluoro pyrimidine nucleotides and purine ribonucleotides with a terminal 3' phosphorothioate linkage (data shown in **Figure 28**). The “Stab7/8” (**Table IV**) constructs comprise a sense strand (Sirna/RPI#30612) having 2'-deoxy-2'-fluoro pyrimidine nucleotides and 2'-deoxy purine nucleotides with 5' and 3' terminal inverted deoxyabasic residues and an antisense strand (Sirna/RPI#30620) having 2'-deoxy-2'-fluoro pyrimidine nucleotides and 2'-O-methyl purine nucleotides with a terminal 3' phosphorothioate linkage (data shown in **Figure 29**). The “Stab7/11” (**Table IV**) constructs comprise a sense (Sirna/RPI#30612) strand having 2'-deoxy-2'-fluoro pyrimidine nucleotides and 2'-deoxy purine nucleotides with 5' and 3' terminal inverted deoxyabasic residues and an antisense strand (Sirna/RPI#31175) having 2'-deoxy-2'-fluoro pyrimidine nucleotides and 2'-deoxy purine nucleotides with a terminal 3' phosphorothioate linkage (data shown in **Figure 30**). The “Stab9/10” (**Table IV**) constructs comprise a sense (Sirna/RPI#31335) strand having ribonucleotides with 5' and 3' terminal inverted deoxyabasic residues and an antisense strand (Sirna/RPI#31337) having ribonucleotides with a terminal 3' phosphorothioate linkage (data shown in **Figure 31**). As shown in **Figures 28-31**, the chemically stabilized siNA constructs all show significantly greater inhibition of HBV antigen in a dose dependent manner over the time course experiment compared to the unstabilized siRNA construct.

A second study was performed using the stab 4/5 (Sirna 30355/30366), stab 7/8 (Sirna 30612/30620), and stab 7/11 (Sirna 30612/31175) siNA constructs described above to examine the duration of effect of the modified siNA constructs out to 21 days post transfection compared to an all RNA control siNA (Sirna 30287/30298). A single transfection was performed with siRNAs targeted to HBV site 1580 and the culture media

was subsequently replaced every three days. Secreted HBsAg levels were monitored for at 3, 6, 9, 12, 15, 18 and 21 days post-transfection. **Figure 77** shows activity of siNAs in reduction of HBsAg levels compared to matched inverted controls at **A.** 3 days, **B.** 9 days, and **C.** 21 days post transfection. Also shown is the corresponding percent inhibition as function of time at siNA concentrations of **D.** 100 nM, **E.** 50 nM, and **F.** 25 nM.

Example 14: Inhibition of HCV using siNA Molecules of the Invention

siNA Inhibition of a chimeric HCV/Poliovirus in HeLa Cells

Inhibition of a chimeric HCV/Poliovirus was investigated using 21 nucleotide siNA duplexes in HeLa cells. Seven siNA constructs were designed that target three regions in the highly conserved 5' untranslated region (UTR) of HCV RNA. The siNAs were screened in two cell culture systems dependent upon the 5'-UTR of HCV; one requires translation of an HCV/luciferase gene, while the other involves replication of a chimeric HCV/poliovirus (PV) (see Blatt *et al.*, USSN 09/740,332, filed December 18, 2000, incorporated by reference herein). Two siNAs (29579/29586; 29578/29585) targeting the same region (shifted by one nucleotide) are active in both systems (see **Figure 32**) as compared with inverse control siNA (29593/29600). For example, a >85% reduction in HCVPV replication was observed in siNA-treated cells compared to an inverse siNA control (**Figure 32**) with an IC₅₀ = ~2.5 nM (**Figure 33**). To develop nuclease-resistant siNA for in vivo applications, siNAs can be modified to contain stabilizing chemical modifications. Such modifications include phosphorothioate linkages (P=S), 2'-O-methyl nucleotides, 2'-fluoro (F) nucleotides, 2'-deoxy nucleotides, universal base nucleotides, 5' and/or 3' end modifications and a variety of other nucleotide and non-nucleotide modifications, in one or both siNA strands. Several of these constructs were tested in the HCV/poliovirus chimera system, demonstrating significant reduction in viral replication (**Figures 34-37**). siNA constructs shown in **Figures 34-37** are referred to by Sirna/RPI#s that are cross referenced to **Table III**, which shows the sequence and chemical modifications of the constructs. siNA activity is compared to relevant controls (untreated cells, scrambled/inactive control sequences, or transfection controls). As shown in the Figures, siNA constructs of the invention provide potent inhibition of HCV RNA in the HCV/poliovirus chimera system. As such, siNA constructs, including

chemically modified, nuclease resistant siNA molecules, represent an important class of therapeutic agents for treating chronic HCV infection.

siNA Inhibition of a HCV RNA expression in a HCV replicon system

In addition, a HCV replicon system was used to test the efficacy of siNAs targeting HCV RNA. The reagents are tested in cell culture using Huh7 cells (see for example Randall *et al.*, 2003, *PNAS USA*, 100, 235-240) to determine the extent of RNA and protein inhibition. siNA were selected against the HCV target as described herein. RNA inhibition was measured after delivery of these reagents by a suitable transfection agent to Huh7 cells. Relative amounts of target RNA are measured versus actin using real-time PCR monitoring of amplification (*eg.*, ABI 7700 Taqman®). A comparison is made to a mixture of oligonucleotide sequences designed to target unrelated targets or to a randomized siNA control with the same overall length and chemistry, but with randomly substituted nucleotides at each position. Primary and secondary lead reagents were chosen for the target and optimization performed. After an optimal transfection agent concentration is chosen, a RNA time-course of inhibition is performed with the lead siNA molecule. In addition, a cell-plating format can be used to determine RNA inhibition. A non-limiting example of a multiple target screen to assay siNA mediated inhibition of HCV RNA is shown in **Figure 38**. siNA reagents (Table I) were transfected at 25 nM into Huh7 cells and HCV RNA quantitated compared to untreated cells (“cells” column in the figure) and cells transfected with lipofectamine (“LFA2K” column in the figure). As shown in the Figure, several siNA constructs show significant inhibition of HCV RNA expression in the Huh7 replicon system. Chemically modified siNA constructs were then screened as described above, with a non-limiting example of a Stab 7/8 (see **Table IV**) chemistry siNA construct screen shown in **Figure 40**. A follow up dose response study using chemically modified siNA constructs (Stab 4/5, see **Table IV**) at concentrations of 5nM, 10nM, 25 nM and 100 nM compared to matched chemistry inverted controls is shown in **Figure 39**, whereas a dose response study for Stab 7/8 constructs at concentrations of 5nM, 10nM, 25 nM, 50 nM and 100 nM compared to matched chemistry inverted controls is shown in **Figure 41**. A separate direct screen of Stab 7/8 constructs targeting HCV RNA that identified stabilized siNA constructs with potent activity is shown in **Figure 86**.

Example 15: Target Discovery in Mammalian Cells using siNA molecules

In a non-limiting example, compositions and methods of the invention are used to discover genes involved in a process of interest within mammalian cells, such as cell growth, proliferation, apoptosis, morphology, angiogenesis, differentiation, migration, viral multiplication, drug resistance, signal transduction, cell cycle regulation, or temperature sensitivity or other process. First, a randomized siNA library is generated. These constructs are inserted into a vector capable of expressing a siNA from the library inside mammalian cells. Alternately, a pool of synthetic siNA molecules is generated.

Reporter System

In order to discover genes playing a role in the expression of certain proteins, such as proteins involved in a cellular process described herein, a readily assayable reporter system is constructed in which a reporter molecule is co-expressed when a particular protein of interest is expressed. The reporter system consists of a plasmid construct bearing a gene coding for a reporter gene, such as Green Fluorescent Protein (GFP) or other reporter proteins known and readily available in the art. The promoter region of the GFP gene is replaced by a portion of a promoter for the protein of interest sufficient to direct efficient transcription of the GFP gene. The plasmid can also contain a drug resistance gene, such as neomycin resistance, in order to select cells containing the plasmid.

Host Cell Lines for Target Discovery

A cell line is selected as host for target discovery. The cell line is preferably known to express the protein of interest, such that upstream genes controlling the expression of the protein can be identified when modulated by a siNA construct expressed therein. The cells preferably retain protein expression characteristics in culture. The reporter plasmid is transfected into cells, for example, using a cationic lipid formulation. Following transfection, the cells are subjected to limiting dilution cloning, for example, under selection by 600 µg/mL Geneticin. Cells retaining the plasmid survive the Geneticin treatment and form colonies derived from single surviving cells. The resulting clonal cell lines are screened by flow cytometry for the capacity to upregulate GFP production. Treating the cells with, for example, sterilized M9 bacterial medium in which

Pseudomonas aeruginosa had been cultured (Pseudomonas conditioned medium, PCM) is used to induce the promoter. The PCM is supplemented with phorbol myristate acetate (PMA). A clonal cell line highly responsive to promoter induction is selected as the reporter line for subsequent studies.

5 *siNA Library Construction*

A siNA library was constructed with oligonucleotides containing hairpin siNA constructs having randomized antisense regions and self complementary sense regions. The library is generated synthesizing siNA constructs having randomized sequence. Alternately, the siNA libraries are constructed as described in Usman *et al.*, USSN 10 60/402,996 (incorporated by reference herein) Oligo sequence 5' and 3' of the siNA contains restriction endonuclease cleavage sites for cloning. The 3' trailing sequence forms a stem-loop for priming DNA polymerase extension to form a hairpin structure. The hairpin DNA construct is melted at 90°C allowing DNA polymerase to generate a dsDNA construct. The double-stranded siNA library is cloned into, for example, a 15 U6+27 transcription unit located in the 5' LTR region of a retroviral vector containing the human nerve growth factor receptor (hNGFr) reporter gene. Positioning the U6+27/siNA transcription unit in the 5' LTR results in a duplication of the transcription unit when the vector integrates into the host cell genome. As a result, the siNA is transcribed by RNA polymerase III from U6+27 and by RNA polymerase II activity directed by the 5' LTR. 20 The siNA library is packaged into retroviral particles that are used to infect and transduce clonal cells selected above. Assays of the hNGFr reporter are used to indicate the percentage of cells that incorporated the siNA construct. By randomized region is meant a region of completely random sequence and/or partially random sequence. By completely random sequence is meant a sequence wherein theoretically there is equal 25 representation of A, T, G and C nucleotides or modified derivatives thereof, at each position in the sequence. By partially random sequence is meant a sequence wherein there is an unequal representation of A, T, G and C nucleotides or modified derivatives thereof, at each position in the sequence. A partially random sequence can therefore have one or more positions of complete randomness and one or more positions with defined 30 nucleotides.

Enriching for Non-responders to Induction

Sorting of siNA library-containing cells is performed to enrich for cells that produce less reporter GFP after treatment with the promoter inducers PCM and PMA. Lower GFP production can be due to RNAi activity against genes involved in the activation of the mucin promoter. Alternatively, siNA can directly target the mucin/GFP transcript resulting in reduced GFP expression.

Cells are seeded at a certain density, such as 1×10^6 per 150 cm² style cell culture flasks and grown in the appropriate cell culture medium with fetal bovine serum. After 72 hours, the cell culture medium is replaced with serum-free medium. After 24 hours of serum deprivation, the cells are treated with serum-containing medium supplemented with PCM (to 40%) and PMA (to 50 nM) to induced GFP production. After 20 to 22 hours, cells are monitored for GFP level on, for example, a FACStar Plus cell sorter. Sorting is performed if $\geq 90\%$ of siNA library cells from an unsorted control sample were induced to produce GFP above background levels. Two cell fractions are collected in each round of sorting. Following the appropriate round of sorting, the M1 fraction is selected to generate a database of siNA molecules present in the sorted cells.

Recovery of siNA Sequence from Sorted Cells

Genomic DNA is obtained from sorted siNA library cells by standard methods. Nested polymerase chain reaction (PCR) primers that hybridized to the retroviral vector 5' and 3' of the siNA are used to recover and amplify the siNA sequences from the particular clone of library cell DNA. The PCR product is ligated into a bacterial cloning vector. The recovered siNA library in plasmid form can be used to generate a database of siNA sequences. For example, the library is cloned into *E. coli*. DNA is prepared by plasmid isolation from bacterial colonies or by direct colony PCR and siNA sequence is determined. A second method can use the siNA library to transfect cloned cells. Clonal lines of stably transfected cells are established and induced with, for example, PCM and PMA. Those lines which fail to respond to GFP induction are probed by PCR for single siNA integration events. The unique siNA sequences obtained by both methods are added to a Target Sequence Tag (TST) database.

Bioinformatics

The antisense region sequences of the isolated siNA constructs are compared to public and private gene data banks. Gene matches are compiled according to perfect and imperfect matches. Potential gene targets are categorized by the number of different siNA sequences matching each gene. Genes with more than one perfect siNA match are
 5 selected for Target Validation studies.

Validation of the Target Gene

To validate a target as a regulator of protein expression, siNA reagents are designed to the target gene cDNA sequence from Genbank. The siNA reagents are complexed with a cationic lipid formulation prior to administration to cloned cells at appropriate
 10 concentrations (e.g. 5-50 nM or less). Cells are treated with siNA reagents, for example from 72 to 96 hours. Before the termination of siNA treatment, PCM (to 40 %) and PMA (to 50 nM), for example, are added to induce the promoter. After twenty hours of induction the cells are harvested and assayed for phenotypic and molecular parameters. Reduced GFP expression in siNA treated cells (measured by flow cytometry) is taken as
 15 evidence for validation of the target gene. Knockdown of target RNA in siNA treated cells can correlate with reduced endogenous RNA and reduced GFP RNA to complete validation of the target.

Example 16: Screening siNA constructs for improved pharmacokinetics

In a non-limiting example, siNA constructs are screened in vivo for improved
 20 pharmacokinetic properties compared to all RNA or unmodified siNA constructs. Chemical modifications are introduced into the siNA construct based on educated design parameters (e.g. introducing 2'-modifications, base modifications, backbone modifications, terminal cap modifications, or covalently attached conjugates etc). The modified construct is tested in an appropriate system (e.g human serum for nuclease resistance,
 25 shown, or an animal model for PK/delivery parameters). In parallel, the siNA construct is tested for RNAi activity, for example in a cell culture system such as a luciferase reporter assay). Lead siNA constructs are then identified which possess a particular characteristic while maintaining RNAi activity, and can be further modified and assayed once again. This same approach can be used to identify siNA-conjugate molecules with improved
 30 pharmacokinetic profiles, delivery, localized delivery, cellular uptake, and RNAi activity.

Example 17: Indications

The siNA molecules of the invention can be used to treat a variety of diseases and conditions through modulation of gene expression. Using the methods described herein, chemically modified siNA molecules can be designed to modulate the expression any

5 number of target genes, including but not limited to genes associated with cancer, metabolic diseases, infectious diseases such as viral, bacterial or fungal infections, neurologic diseases, musculoskeletal diseases, diseases of the immune system, diseases associated with signaling pathways and cellular messengers, and diseases associated with transport systems including molecular pumps and channels.

10 Non-limiting examples of various viral genes that can be targeted using siNA molecules of the invention include Hepatitis C Virus (HCV, for example Genbank Accession Nos: D11168, D50483.1, L38318 and S82227), Hepatitis B Virus (HBV, for example GenBank Accession No. AF100308.1), Human Immunodeficiency Virus type 1 (HIV-1, for example GenBank Accession No. U51188), Human Immunodeficiency Virus

15 type 2 (HIV-2, for example GenBank Accession No. X60667), West Nile Virus (WNV for example GenBank accession No. NC_001563), cytomegalovirus (CMV for example GenBank Accession No. NC_001347), respiratory syncytial virus (RSV for example GenBank Accession No. NC_001781), influenza virus (for example example GenBank Accession No. AF037412, rhinovirus (for example, GenBank accession numbers:

20 D00239, X02316, X01087, L24917, M16248, K02121, X01087), papillomavirus (for example GenBank Accession No. NC_001353), Herpes Simplex Virus (HSV for example GenBank Accession No. NC_001345), and other viruses such as HTLV (for example GenBank Accession No. AJ430458). Due to the high sequence variability of many viral genomes, selection of siNA molecules for broad therapeutic applications would likely

25 involve the conserved regions of the viral genome. Nonlimiting examples of conserved regions of the viral genomes include but are not limited to 5'-Non Coding Regions (NCR), 3'- Non Coding Regions (NCR) LTR regions and/or internal ribosome entry sites (IRES). siNA molecules designed against conserved regions of various viral genomes will enable efficient inhibition of viral replication in diverse patient populations and may

30 ensure the effectiveness of the siNA molecules against viral quasi species which evolve due to mutations in the non-conserved regions of the viral genome.

Non-limiting examples of human genes that can be targeted using siNA molecules of the invention using methods described herein include any human RNA sequence, for example those commonly referred to by Genbank Accession Number. These RNA sequences can be used to design siNA molecules that inhibit gene expression and therefore abrogate diseases, conditions, or infections associated with expression of those genes. Such non-limiting examples of human genes that can be targeted using siNA molecules of the invention include VEGF (for example GenBank Accession No. NM_003376.3), VEGFr (VEGFR1 for example GenBank Accession No. XM_067723, VEGFR2 for example GenBank Accession No. AF063658), HER1, HER2, HER3, and HER4 (for example Genbank Accession Nos: NM_005228, NM_004448, NM_001982, and NM_005235 respectively), telomerase (TERT, for example GenBank Accession No. NM_003219), telomerase RNA (for example GenBank Accession No. U86046), NFkappaB, Rel-A (for example GenBank Accession No. NM_005228), NOGO (for example GenBank Accession No. AB020693), NOGO (for example GenBank Accession No. XM_015620), RAS (for example GenBank Accession No. NM_004283), RAF (for example GenBank Accession No. XM_033884), CD20 (for example GenBank Accession No. X07203), METAP2 (for example GenBank Accession No. NM_003219), CLCA1 (for example GenBank Accession No. NM_001285), phospholamban (for example GenBank Accession No. NM_002667), PTP1B (for example GenBank Accession No. M31724), PCNA (for example GenBank Accession No. NM_002592.1), PKC-alpha (for example GenBank Accession No. NM_002737) and others. The genes described herein are provided as non-limiting examples of genes that can be targeted using siNA molecules of the invention. Additional examples of such genes are described by accession number in Beigelman *et al.*, USSN 60/363,124, filed March 11, 2002 and incorporated by reference herein in its entirety.

The siNA molecule of the invention can also be used in a variety of agricultural applications involving modulation of endogenous or exogenous gene expression in plants using siNA, including use as insecticidal, antiviral and anti-fungal agents or modulate plant traits such as oil and starch profiles and stress resistance.

Example 18: Diagnostic uses

The siNA molecules of the invention can be used in a variety of diagnostic applications, such as in the identification of molecular targets (e.g., RNA) in a variety of applications, for example, in clinical, industrial, environmental, agricultural and/or research settings. Such diagnostic use of siNA molecules involves utilizing reconstituted RNAi systems, for example, using cellular lysates or partially purified cellular lysates. siNA molecules of this invention can be used as diagnostic tools to examine genetic drift and mutations within diseased cells or to detect the presence of endogenous or exogenous, for example viral, RNA in a cell. The close relationship between siNA activity and the structure of the target RNA allows the detection of mutations in any region of the molecule, which alters the base-pairing and three-dimensional structure of the target RNA. By using multiple siNA molecules described in this invention, one can map nucleotide changes, which are important to RNA structure and function *in vitro*, as well as in cells and tissues. Cleavage of target RNAs with siNA molecules can be used to inhibit gene expression and define the role of specified gene products in the progression of disease or infection. In this manner, other genetic targets can be defined as important mediators of the disease. These experiments will lead to better treatment of the disease progression by affording the possibility of combination therapies (e.g., multiple siNA molecules targeted to different genes, siNA molecules coupled with known small molecule inhibitors, or intermittent treatment with combinations siNA molecules and/or other chemical or biological molecules). Other *in vitro* uses of siNA molecules of this invention are well known in the art, and include detection of the presence of mRNAs associated with a disease, infection, or related condition. Such RNA is detected by determining the presence of a cleavage product after treatment with a siNA using standard methodologies, for example, fluorescence resonance emission transfer (FRET).

In a specific example, siNA molecules that cleave only wild-type or mutant forms of the target RNA are used for the assay. The first siNA molecules (*i.e.*, those that cleave only wild-type forms of target RNA) are used to identify wild-type RNA present in the sample and the second siNA molecules (*i.e.*, those that cleave only mutant forms of target RNA) are used to identify mutant RNA in the sample. As reaction controls, synthetic substrates of both wild-type and mutant RNA are cleaved by both siNA molecules to demonstrate the relative siNA efficiencies in the reactions and the absence of cleavage of

the "non-targeted" RNA species. The cleavage products from the synthetic substrates also serve to generate size markers for the analysis of wild-type and mutant RNAs in the sample population. Thus, each analysis requires two siNA molecules, two substrates and one unknown sample, which is combined into six reactions. The presence of cleavage products is determined using an RNase protection assay so that full-length and cleavage fragments of each RNA can be analyzed in one lane of a polyacrylamide gel. It is not absolutely required to quantify the results to gain insight into the expression of mutant RNAs and putative risk of the desired phenotypic changes in target cells. The expression of mRNA whose protein product is implicated in the development of the phenotype (*i.e.*, disease related or infection related) is adequate to establish risk. If probes of comparable specific activity are used for both transcripts, then a qualitative comparison of RNA levels is adequate and decreases the cost of the initial diagnosis. Higher mutant form to wild-type ratios are correlated with higher risk whether RNA levels are compared qualitatively or quantitatively.

Example 19: Synthesis of siNA conjugates

The introduction of conjugate moieties to siNA molecules of the invention is accomplished either during solid phase synthesis using phosphoramidite chemistry described above, or post-synthetically using, for example, N-hydroxysuccinimide (NHS) ester coupling to an amino linker present in the siNA. Typically, a conjugate introduced during solid phase synthesis will be added to the 5'-end of a nucleic acid sequence as the final coupling reaction in the synthesis cycle using the phosphoramidite approach. Coupling conditions can be optimized for high yield coupling, for example by modification of coupling times and reagent concentrations to effectuate efficient coupling. As such, the 5'-end of the sense strand of a siNA molecule is readily conjugated with a conjugate moiety having a reactive phosphorus group available for coupling (e.g., a compound having Formulae 1, 5, 8, 55, 56, 57, 60, 86, 92, 104, 110, 113, 115, 116, 117, 118, 120, or 122) using the phosphoramidite approach, providing a 5'-terminal conjugate (see for example **Figure 65**).

Conjugate precursors having a reactive phosphorus group and a protected hydroxyl group can be used to incorporate a conjugate moiety anywhere in the siNA sequence, such as in the loop portion of a single stranded hairpin siNA construct (see for example

Figure 66). For example, using the phosphoramidite approach, a conjugate moiety comprising a phosphoramidite and protected hydroxyl (e.g., a compound having Formulae 86, 92, 104, 113, 115, 116, 117, 118, 120, or 122 herein) is first coupled at the desired position within the siNA sequence using solid phase synthesis phosphoramidite coupling. Second, removal of the protecting group (e.g., dimethoxytrityl) allows coupling of additional nucleotides to the siNA sequence. This approach allows the conjugate moiety to be positioned anywhere within the siNA molecule.


Conjugate derivatives can also be introduced to a siNA molecule post synthetically. Post synthetic conjugation allows a conjugate moiety to be introduced at any position within the siNA molecule where an appropriate functional group is present (e.g., a C5 alkylamine linker present on a nucleotide base or a 2'-alkylamine linker present on a nucleotide sugar can provide a point of attachment for an NHS-conjugate moiety). Generally, a reactive chemical group present in the siNA molecule is unmasked following synthesis, thus allowing post-synthetic coupling of the conjugate to occur. In a non-limiting example, an protected amino linker containing nucleotide (e.g., TFA protected C5 propylamino thymidine) is introduced at a desired position of the siNA during solid phase synthesis. Following cleavage and deprotection of the siNA, the free amine is made available for NHS ester coupling of the conjugate at the desired position within the siNA sequence, such as at the 3'-end of the sense and/or antisense strands, the 3' and/or 5'-end of the sense strand, or within the siNA sequence, such as in the loop portion of a single stranded hairpin siNA sequence.

A conjugate moiety can be introduced at different locations within a siNA molecule using both solid phase synthesis and post-synthetic coupling approaches. For example, solid phase synthesis can be used to introduce a conjugate moiety at the 5'-end of the siNA (e.g. sense strand) and post-synthetic coupling can be used to introduce a conjugate moiety at the 3'-end of the siNA (e.g. sense strand and/or antisense strand). As such, a siNA sense strand having 3' and 5' end conjugates can be synthesized (see for example **Figure 65**). Conjugate moieties can also be introduced in other combinations, such as at the 5'-end, 3'-end and/or loop portions of a siNA molecule (see for example **Figure 66**).

Example 20: Pharmacokinetics of siNA conjugates (Figure 67)


Three nuclease resistant siNA molecule targeting site 1580 of hepatitis B virus (HBV) RNA were designed using Stab 7/8 chemistry (see **Table IV**) and a 5'-terminal conjugate moiety.

One siNA conjugate comprises a branched cholesterol conjugate linked to the sense strand of the siNA. The “cholesterol” siNA conjugate molecule has the structure shown below:




where T stands for thymidine, B stands for inverted deoxyabasic, G stands for 2'-deoxy guanosine, A stands for 2'-deoxy adenosine, G stands for 2'-O-methyl guanosine, A stands for 2'-O-methyl adenosine, u stands for 2'-fluoro uridine, c stands for 2'-fluoro cytidine, a stands for adenosine, and s stands for phosphorothioate linkage.

Another siNA conjugate comprises a branched phospholipid conjugate linked to the sense strand of the siNA. The “phospholipid” siNA conjugate molecule has the structure shown below:



where T stands for thymidine, B stands for inverted deoxyabasic, G stands for 2'-deoxy guanosine, A stands for 2'-deoxy adenosine, G stands for 2'-O-methyl guanosine, A stands for 2'-O-methyl adenosine, u stands for 2'-fluoro uridine, c stands for 2'-fluoro cytidine, a stands for adenosine, and s stands for phosphorothioate linkage.

Another siNA conjugate comprises a polyethylene glycol (PEG) conjugate linked to the sense strand of the siNA. The "PEG" siNA conjugate molecule has the structure shown below:



where T stands for thymidine, B stands for inverted deoxyabasic, G stands for 2'-deoxy guanosine, A stands for 2'-deoxy adenosine, G stands for 2'-O-methyl guanosine, A stands for 2'-O-methyl adenosine, u stands for 2'-fluoro uridine, c stands for 2'-fluoro cytidine, a stands for adenosine, and s stands for phosphorothioate linkage.

5 The Cholesterol, Phospholipid, and PEG conjugates were evaluated for pharmacokinetic properties in mice compared to a non-conjugated siNA construct having matched chemistry and sequence. This study was conducted in female CD-1 mice approximately 26 g (6-7 weeks of age). Animals were housed in groups of 3. Food and water were provided ad libitum. Temperature and humidity were according to
10 Pharmacology Testing Facility performance standards (SOP's) which are in accordance with the 1996 Guide for the Care and Use of Laboratory Animals (NRC). Animals were acclimated to the facility for at least 3 days prior to experimentation.

Absorbance at 260 nm was used to determine the actual concentration of the stock solution of pre-annealed HBV siNA. An appropriate amount of HBV siNA was diluted
15 in sterile veterinary grade normal saline (0.9%) based on the average body weight of the mice. A small amount of the antisense (Stab 7) strand was internally labeled with gamma 32P-ATP. The 32P-labeled stock was combined with excess sense strand (Stab 8) and annealed. Annealing was confirmed prior to combination with unlabeled drug. Each mouse received a subcutaneous bolus of 30 mg/kg (based on duplex) and approximately
20 10 million cpm (specific activity of approximately 15 cpm/ng).

Three animals per timepoint (1, 4, 8, 24, 72, 96 h) were euthanized by CO2 inhalation followed immediately by exsanguination. Blood was sampled from the heart and collected in heparinized tubes. After exsanguination, animals were perfused with 10-15 mL of sterile veterinary grade saline via the heart. Samples of liver were then
25 collected and frozen.

Tissue samples were homogenized in a digestion buffer prior to compound quantitation. Quantitation of intact compound was determined by scintillation counting followed by PAGE and phosphorimage analysis. Results are shown in Figure 43. As shown in the figure, the conjugated siNA constructs shown vastly improved liver PK
30 compared to the unconjugated siNA construct.

Example 21: Cell culture of siNA conjugates (Figure 68)

The Cholesterol conjugates and Phospholipid conjugated siNA constructs described in Example 20 above were evaluated for cell culture efficacy in a HBV cell culture system.

5 *Transfection of HepG2 Cells with psHBV-1 and siNA*

The human hepatocellular carcinoma cell line Hep G2 was grown in Dulbecco's modified Eagle media supplemented with 10% fetal calf serum, 2 mM glutamine, 0.1 mM nonessential amino acids, 1 mM sodium pyruvate, 25 mM Hepes, 100 units penicillin, and 100 µg/ml streptomycin. To generate a replication competent cDNA, prior to
 10 transfection the HBV genomic sequences are excised from the bacterial plasmid sequence contained in the psHBV-1 vector. Other methods known in the art can be used to generate a replication competent cDNA. This was done with an EcoRI and Hind III restriction digest. Following completion of the digest, a ligation was performed under dilute conditions (20 µg/ml) to favor intermolecular ligation. The total ligation mixture
 15 was then concentrated using Qiagen spin columns.

siNA Activity Screen and Dose Response Assay

Transfection of the human hepatocellular carcinoma cell line, Hep G2, with replication-competent HBV DNA results in the expression of HBV proteins and the production of virions. To test the efficacy of siNA conjugates targeted against HBV
 20 RNA, the Cholesterol siNA conjugate and Phospholipid siNA conjugate described in Example 12 were compared to a non-conjugated control siNA (see **Figure 68**). An inverted sequence duplex was used as a negative control for the unconjugated siNA. Dose response studies were performed in which HBV genomic DNA was transfected with HBV genomic DNA with lipid at 12.5 µg/ml into Hep G2 cells. 24 hours after
 25 transfection with HBV DNA, cell culture media was removed and siNA duplexes were added to cells without lipid at 10 µM, 5 µM, 2.5 µM, 1 µM, and 100 nM and the subsequent levels of secreted HBV surface antigen (HBsAg) were analyzed by ELISA 72 hours post treatment (see Figure 44). To determine siNA activity, HbsAg levels were measured following transfection with siNA. Immulon 4 (Dynax) microtiter wells were
 30 coated overnight at 4° C with anti-HBsAg Mab (Biostride B88-95-31ad,ay) at 1 µg/ml in

Carbonate Buffer (Na₂CO₃ 15 mM, NaHCO₃ 35 mM, pH 9.5). The wells were then washed 4x with PBST (PBS, 0.05% Tween® 20) and blocked for 1 hr at 37° C with PBST, 1% BSA. Following washing as above, the wells were dried at 37° C for 30 min. Biotinylated goat anti-HBsAg (Accurate YVS1807) was diluted 1:1000 in PBST and
 5 incubated in the wells for 1 hr. at 37° C. The wells were washed 4x with PBST. Streptavidin/Alkaline Phosphatase Conjugate (Pierce 21324) was diluted to 250 ng/ml in PBST, and incubated in the wells for 1 hr. at 37° C. After washing as above, p-nitrophenyl phosphate substrate (Pierce 37620) was added to the wells, which were then incubated for 1 hour at 37° C. The optical density at 405 nm was then determined. As
 10 shown in **Figure 68**, the phospholipid and cholesterol conjugates demonstrate marked dose dependent inhibition of HBsAg expression compared to the unconjugated siNA construct when delivered to cells without any transfection agent (lipid).

Example 22: Ex vivo stability of siNA constructs

Chemically modified siNA constructs were designed and synthesized in order
 15 to improve resistance to nucleases while maintaining silencing in cell culture systems. Modified strands, designated Stab 4, Stab 5, Stab 7, Stab 8, and Stab 11 (**Table IV**), were tested in three sets of duplexes that demonstrated a range of stability and activity. These duplexes contained differentially modified sense and antisense strands. All modified sense strands contain terminal 5' and 3' inverted abasic caps, while antisense strands
 20 possess a 3' terminal phosphorothioate linkage. The results characterize the impact of chemical modifications on nuclease resistance in *ex vivo* models of the environments sampled by drugs.

Active siNAs were assessed for their resistance to degradation in serum and liver extracts. Stability in blood will be a requirement for a systemically administered siNA,
 25 and an anti-HBV or anti-HCV siNA would require stability and activity in the hepatic intracellular environment. Liver extracts potentially provide an extreme nuclease model where many catabolic enzymes are present. Both mouse and human systems were assessed.

Individual strands of siNA duplexes were internally labeled with ^{32}P and incubated as single strands or as duplex siRNAs in human or mouse serum and liver extracts. Representative data is shown in **Table VI**. Throughout the course of the experiments, constant levels of ribonuclease activity were verified. The extent and pattern of all-RNA siNA degradation (3 minute time point) did not change following preincubation of serum or liver extract at 37°C for up to 24 hours.

The biological activity of siRNAs containing all-ribose residues has been well established. The extreme instability ($t_{1/2} = 0.017$ hours) of these compounds in serum underscores the need for chemical modification for use in systemic therapeutic applications. The Stab 4/5 duplex modifications provide significant stability in human and mouse serum ($t_{1/2}$'s = 10 – 408 hours) and human liver extract ($t_{1/2}$'s = 28 – 43 hours). In human serum the Stab 4 strand chemistry in the context of the Stab 4/5 duplex, possesses greater stability than the Stab 5 strand chemistry ($t_{1/2} = 408$ vs. 39 hours). This result highlights the impact terminal modifications have on stability. A fully-modified Stab 7/11 construct (no ribonucleotides present) was generated from the Stab 4/5 constructs by substituting the ribonucleotides in all purine positions with deoxyribonucleotides. Another fully modified construct, Stab 7/8, was generated by replacing all purine positions in the antisense strand with 2'-O-methyl nucleotides. This proved to be the most stable antisense strand chemistry observed, with $t_{1/2} = 816$ hours in human liver extract.

The dramatic stability of Stab 8 modifications was also observed when non-duplexed single strands were incubated in human serum and liver extract, as shown in **Table VII**. An approximate five-fold increase in serum stability is seen for the double stranded constructs, compared to that observed for the individual strands. In liver extract, the siNA duplex provides even greater stability compared to the single strands. For example, the Stab 5 chemistry is greater than 100-fold more stable in the Stab 4/5 duplex relative to its stability alone.

Terminal modifications have a large impact on stability in human serum, as can be seen from a comparison of sense verses antisense stabilities in duplex form, and the Stab 4 and Stab 5 single-strand stabilities. Therefore, a number of 3' antisense capping moieties on Stab 4/5 chemistry duplexes were assessed for their contribution to stability in human serum. The structures of these modifications are shown in **Figure 22**, and resultant half-lives are shown in **Table VIII**. A wide range of different stabilities were observed, from half-lives as short as one hour to greater than 770 hours. Thus, in the context of 2'-fluoro modified pyrimidines, 3'-exonuclease becomes the primary mode of attack on duplexes in human serum; a number of chemistries minimize this site of attack. These results suggest that susceptibility to 3' exonucleases is a major path to degradation in the serum.

Example 23: Activity of siNA molecules delivered via hydrodynamic injection

An in vivo mouse model that utilizes hydrodynamic tail vein injection of a replication competent HBV vector has been used to assess the activity of chemically stabilized siRNA targeted to HBV RNA. The hydrodynamic delivery of nucleic acids in the mouse has been described by Liu *et al.*, 1999, *Gene Therapy*, 6, 1258-1266, who showed that the vast majority of the nucleic acid is delivered to the liver by this technique. The use of the hydrodynamic technology to develop an HBV mouse model has been described by Yang *et al.*, 2002, *PNAS*, 99, 13825-13830. In the vector-based model, HBV replicates in the liver for approximately 10 days, resulting in detectable levels of HBV RNA and antigens in the liver and HBV DNA and antigens in the serum.

To assess the activity of chemically stabilized siNAs against HBV, co-injection of the siNAs along with the HBV vector was done in mouse strain C57BL/6. The HBV vector used, pWTD, is a head-to-tail dimer of the complete HBV genome (see for example Buckwold *et al.*, 1996, *J. Virology*, 70, 5845-5851). For a 20 gram mouse, a total injection of 1.6 ml containing 10 μ g or 1 μ g of pWTD and 100 μ g of siNA duplex in saline, was injected into the tail vein within 5 seconds. For a larger mouse, the volume is scaled to maintain a level of 140% of the blood volume of the mouse. The injection is done using a 3 cc syringe and 27g1/2 needle. The animals were sacrificed at 72 hrs post-injection. Animals were treated with siNA constructs and matched chemistry inverted controls. Analysis of the HBV DNA (**Figure 80**) and HBsAg (**Figure 81**) levels in serum

was conducted by real-time PCR and ELISA respectively. The levels of HBV RNA in the liver (**Figure 82**) were analyzed by real-time RT-PCR. In a separate experiment, analysis of HBV DNA levels in serum was carried out at 5 days and 7 days (**Figure 83**) after co-injection of siNA and the HBV vector.

5 Example 24: Activity screens using chemically modified siNA

Two formats can be used to identify active chemically modified siNA molecules against target nucleic acid molecules (e.g., RNA). One format involves screening unmodified siNA constructs in an appropriate system (e.g., cell culture or animal models) then applying chemical modifications to the sequence of identified leads and rescreening the modified constructs. Another format involves direct screening of chemically modified constructs to identify chemically modified leads (see for example the Stab 7/8 HCV screen shown in **Figure 86** and the Stab 7/8 HBV screen shown **Figure 87**, as described above). The latter approach can be useful in identifying active constructs that are specific to various combinations of chemical modifications (e.g., Stab 1-18 chemistries shown in **Table V** herein). Additionally, different iterations of such chemical modifications can be assessed using active chemically modified leads and appropriate rules for selective active constructs given a particular chemistry can be established using this approach. Non-limiting examples of such activity screen are described below.

Activity screen of Stab 7/8 constructs targeting luciferase RNA

HeLa cells were co-transfected with pGF3 vector (250ng/well), renilla luciferase vector (10 ng/well) and siNA (0.5 - 25 nM) using 0.5 ul lipofectamine2000 per well. Twenty-four hours post-transfection, the cells were assayed for luciferase activity using the Promega Dual Luciferase Assay Kit per the manufacturer's instruction. siNA constructs having high levels of activity were identified and tested in a dose response assay with concentrations ranging from 0.5 to 25 nM. Results for siNA constructs targeting sites 80, 237, and 1478 are shown in **Figure 84** and sites 1544 and 1607 are shown in **Figure 85**. As shown in the Figures, several active Stab 7/8 constructs were identified that demonstrate potent dose related inhibition of luciferase expression.

Activity screen of combination siNA constructs targeting HBV RNA in HepG2 cells

The HBV HepG2 cell culture system described in Example 13 above was utilized to evaluate the efficacy of various combinations of chemical modifications (**Table V**) in the sense strand and antisense strand of siNA molecules as compared to matched chemistry

5 inverted controls. To determine siNA activity, HbsAg levels were measured following transfection with siNA. Immulon 4 (Dynax) microtiter wells were coated overnight at 4° C with anti-HBsAg Mab (Biostride B88-95-31ad,ay) at 1 µg/ml in Carbonate Buffer (Na₂CO₃ 15 mM, NaHCO₃ 35 mM, pH 9.5). The wells were then washed 4x with PBST (PBS, 0.05% Tween® 20) and blocked for 1 hr at 37° C with PBST, 1% BSA. Following

10 washing as above, the wells were dried at 37° C for 30 min. Biotinylated goat ant-HBsAg (Accurate YVS1807) was diluted 1:1000 in PBST and incubated in the wells for 1 hr. at 37° C. The wells were washed 4x with PBST. Streptavidin/Alkaline Phosphatase Conjugate (Pierce 21324) was diluted to 250 ng/ml in PBST, and incubated in the wells for 1 hr. at 37° C. After washing as above, p-nitrophenyl phosphate substrate (Pierce

15 37620) was added to the wells, which were then incubated for 1 hour at 37° C. The optical density at 450 nm was then determined. Results of the combination HBV siNA screen are shown in **Figures 88-90**. As shown in the Figures, the various combinations of differing sense and antisense chemistries (e.g., sense/antisense constructs having Stab 7/8, 7/10, 7/11, 9/8, 9/10, 6/10, 16/8, 16/10, 18/8, 18/10, 4/8, 4/10, 7/5, 9/5, and 9/11

20 chemistry) result in active siNA constructs.

All patents and publications mentioned in the specification are indicative of the levels of skill of those skilled in the art to which the invention pertains. All references cited in this disclosure are incorporated by reference to the same extent as if each reference had been incorporated by reference in its entirety individually.

25 One skilled in the art would readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The methods and compositions described herein as presently representative of preferred embodiments are exemplary and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to

30 those skilled in the art, which are encompassed within the spirit of the invention, are defined by the scope of the claims.

It will be readily apparent to one skilled in the art that varying substitutions and modifications can be made to the invention disclosed herein without departing from the scope and spirit of the invention. Thus, such additional embodiments are within the scope of the present invention and the following claims. The present invention teaches one
5 skilled in the art to test various combinations and/or substitutions of chemical modifications described herein toward generating nucleic acid constructs with improved activity for mediating RNAi activity. Such improved activity can comprise improved stability, improved bioavailability, and/or improved activation of cellular responses mediating RNAi. Therefore, the specific embodiments described herein are not limiting
10 and one skilled in the art can readily appreciate that specific combinations of the modifications described herein can be tested without undue experimentation toward identifying siNA molecules with improved RNAi activity.

The invention illustratively described herein suitably can be practiced in the absence of any element or elements, limitation or limitations that are not specifically
15 disclosed herein. Thus, for example, in each instance herein any of the terms "comprising", "consisting essentially of", and "consisting of" may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described
20 or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments, optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within
25 the scope of this invention as defined by the description and the appended claims.

In addition, where features or aspects of the invention are described in terms of Markush groups or other grouping of alternatives, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group or other group.

Table I

Sirna/ RPI#	Aliases	Sequence	SEQ ID#
28443	Sirna/RPI GL2 Str1 (sense) 2'-amino U C	cGuAcGcGGA <u>AuA</u> cuucGATT	349
28444	Sirna/RPI GL2 Str2 (antisense) 2'-amino U C	ucGAAG <u>GuA</u> uuccGcGuAcGTT	350
28445	Sirna/RPI GL2 Str1 (sense) 2'-amino U C uT 3'end	cGuAcGcGGAA <u>AuA</u> cuucGAuT	351
28446	Sirna/RPI GL2 Str2 (antisense) 2'-amino U C uT 3'end	ucGAAG <u>GuA</u> uuccGcGuAcGuT	352
30051	HCV-Luc:325U21 siNA 5' P=S + 3' univ. base 2 + 5'/3' invAba (antisense)	BC ₅ C ₅ C ₅ C ₅ G ₅ GGAGGUCUCGUAGAXXB	353
30052	HCV-Luc:325U21 siNA rev 5' P=S + 3' univ. base 2 + 5'/3' invAba (antisense)	BA ₅ G ₅ A ₅ U ₅ G ₅ CUCUGGAGGGCCCCCXXB	354
30053	HCV-Luc:345L21 siNA (325C) (antisense) 5' P=S + 3' univ. base 2 + 3' invAba (sense)	U ₅ C ₅ U ₅ A ₅ C ₅ GAGACCUCCCGGGGXXB	355
30054	HCV-Luc:345L21 siNA (325C) (antisense) rev 5' P=S + 3' univ. base 2 + 3' invAba (sense)	G ₅ G ₅ G ₅ C ₅ CCUCCAGAGCAUCUXXB	356
30055	HCV-Luc:325U21 siNA all Y P=S + 3' univ. base 2 + 5'/3' invAba (antisense)	BC ₅ C ₅ C ₅ C ₅ GGGAGGU ₅ C ₅ U ₅ C ₅ GU ₅ AGAXXB	357
30056	HCV-Luc:325U21 siNA rev all Y P=S + 3' univ. base 2 + 5'/3' invAba (antisense)	BAGAU ₅ GC ₅ U ₅ C ₅ U ₅ GGAGGGC ₅ C ₅ C ₅ C ₅ XXB	358
30057	HCV-Luc:345L21 siNA (325C) (antisense) all Y P=S + 3' univ. base 2 + 3' invAba (sense)	U ₅ C ₅ U ₅ AC ₅ GAGAC ₅ C ₅ U ₅ C ₅ C ₅ GGGGXXB	359
30058	HCV-Luc:345L21 siNA (325C) (antisense) rev all Y P=S + 3' univ. base 2 + 3' invAba (sense)	GGGGC ₅ C ₅ C ₅ U ₅ C ₅ C ₅ AGAGC ₅ AU ₅ C ₅ U ₅ XXB	360
30059	HCV-Luc:325U21 siNA 4/3 P=S ends + Y-2'F + 3' univ. base 2 + 5'/3' invAba (antisense)	Bc ₅ C ₅ C ₅ C ₅ GGGAGGucucGuA ₅ G ₅ A ₅ XXB	361
30060	HCV-Luc:325U21 siNA rev 4/3 P=S ends + all Y-2'F + 3' univ. base 2 + 5'/3' invAba (antisense)	BA ₅ G ₅ A ₅ u ₅ GucucGGAGGGGcc ₅ c ₅ c ₅ XXB	362
30170	HCV-Luc:325U21 siNA all Y-2'F + 3' univ. base 2 + 5'/3' invAba (antisense)	B cccccGGGAGGucucGuAAGAXX B	363
30171	HCV-Luc:325U21 siNA rev all Y-2'F + 3' univ. base 2 + 5'/3' invAba (antisense)	B AGAuGcucGGAGGGGccccXX B	364
30172	HCV-Luc:345L21 siNA (325C) (antisense)	B U ₅ C ₅ U ₅ AC ₅ GAGAC ₅ C ₅ U ₅ C ₅ C ₅ GGGGXX B	365

Simla/ RPI#	Aliases	Sequence	SEQ ID#
	all Y P=S + 3' univ. base 2 + 5'/3' invAba (antisense)		
30173	HCV-Luc:345L21 siNA (325C) (antisense) all Y-2'F	ucuAcGAGAccuuccGGGG	366
30174	HCV-Luc:345L21 siNA (325C) (antisense) rev all Y-2'F	GGGGccuccAGAGcAucu	367
30175	HCV-Luc:345L21 siNA (325C) (antisense) all Y-2'F + 3' univ. base 2	ucuAcGAGAccuuccGGGGXX	368
30176	HCV-Luc:345L21 siNA (325C) (antisense) rev all Y-2'F + 3' univ. base 2	GGGGccuccAGAGcAucuXX	369
30177	HCV-Luc:345L21 siNA (325C) (antisense) all Y-2'F + 3' univ. base 2 + 5'/3' iB	B ucuAcGAGAccuuccGGGGXX B	370
30178	HCV-Luc:325U21 siNA all Y P=S + 3' univ. base 2 + 3' invAba (sense)	C ₅ C ₅ C ₅ C ₅ GGGAGGU ₅ C ₅ U ₅ C ₅ GU ₅ AGAXX B	371
30063	Simla/RPI GL2 Str1 (sense) 2'-F U,C + 3', 5' abasic	BcGuAcGcGGAAuAcuucGATTB	372
30222	Simla/RPI GL2 Str1 (sense) Y 2'-O-Me with 3'-TT & 5'/3' iB	B cGuAcGcGGAAuAcuucGATT B	373
30224	Simla/RPI GL2 Str2 (antisense) Y 2'-F & 3' TsT	ucGAAGuAuuccGcGuAcGT ₅ T	374
30430	Simla/RPI GL2 Str2 (antisense) 2'-F U,C + 5', 3' abasic, A,G= 2'-O-Me	ucgaag uauuucgcuacgT ₅ T	375
30431	Simla/RPI GL2 Str1 (sense) 2'-F U,C + 3', 5' abasic, TT; 2'-O-Me-A,G	BcguacgcggaaucuuucgaTTB	376
30433	Simla/RPI GL2 Str1 (sense) 2'-F U,C + 3', 5' abasic, TT; 2'-deoxy-A,G	BcGuAcGcGGAAuAcuucGATTB	377
30550	Simla/RPI GL2 Str2 (antisense) 2'-F U,C 3'-dT ₅ T	ucGAAGuAuuccGcGuAcGT ₅ t	378
30555	Simla/RPI GL2 Str2 (antisense) 2'-F U,C 3'-glycerol.T	ucGAAGuAuuccGcGuAcGTL	379
30556	Simla/RPI GL2 Str2 (antisense) 2'-F U,C 3'-glycerol,2T	ucGAAGuAuuccGcGuAcGTTL	380
30226	rev Simla/RPI GL2 Str1 (sense) Y 2'-O-Me with 3'-TT & 5'/3' iB	B AGcuucAuAAGGcGcAuGcTT B	381
30227	rev Simla/RPI GL2 Str1 (sense) Y 2'-F with 3'-TT & 5'/3' iB	B AGcuucAuAAGGcGcAuGcTT B	382
30229	rev Simla/RPI GL2 Str2 (antisense) Y 2'-F & 3' TsT	GcAuGcGccuuAuGAAAGcuT ₅ T	383
30434	Simla/RPI GL2 Str1 (sense) 2'-F U,C + 3', 5' Abasic, TT; 2'-O-Me-A,G;ribo core	BcguacgcGAAuAcuucgaTTB	384
30435	Simla/RPI GL2 Str1 (sense) 2'-F U,C + 3', 5' Abasic, TT; 2'-deoxyA,G;ribo core	BcGuAcGcGGAAuAcuucGATTB	385
30546	Simla/RPI GL2 Str2 (antisense) 2'-F U,C 3'-	ucGAAGuAuuccGcGuAcG3T	386

Sirna/ RPI#	Aliases	Sequence	SEQ ID#
	dTT		
30551	Sima/RPI GL2 Str2 (antisense) 2'-F U,C dTddC	ucGAA GuAuuccGcGuAcGTddC	387
30557	Sima/RPI GL2 Str2 (antisense) 2'-F U,C 3'- invertedT,T	ucGAA GuAuuccGcGuAcGT	388
30558	Sima/RPI GL2 Str2 (antisense) 2'-F U,C 3'- invertedT,TT	ucGAA GuAuuccGcGuAcGT T	389
30196	FLT1:2340U21 siRNA sense iB caps w/2'FY's	B cAAccAcAAAAuAcAAcAATT B	419
30416	FLT1:2358L21 siRNA (2340C) (antisense) TsT	uuGuuGuAuuuuGuGGuuGT _s T	420
29548	HBV:394L21 siRNA (414C) (antisense)	GAUGAGGCAUAGCAGCAGGTT	421
29544	HBV:414U21 siRNA pos (sense)	CCUGCUGCUAUGCCUCAUCTT	422
29556	HBV:394L21 siRNA neg (414C) (antisense) inv	GGACGACGAUACGGAGUAGTT	423
29552	HBV:414U21 siRNA pos (sense) inv	CUACUCCGUUUCGUCUCCTT	424
30350	HBV:262U21 siRNA stab04 (sense)	B uGGAcuuucucAAuuuuuA B	425
30361	HBV:280L21 siRNA (262C) (antisense) stab05	GAAAAuuGAGAGAAAGuccAT _s T	426
30372	HBV:262U21 siRNA inv stab04 (sense)	B AuccuuuuAAcucucuuuAcGGu B	427
30383	HBV:280L21 siRNA (262C) (antisense) inv stab05	AccuGAAAGAGAGuuAAAAAGT _s T	428
30352	HBV:380U21 siRNA stab04 (sense)	B uGuGucuGcGGcGuuuuAucA B	429
30363	HBV:398L21 siRNA (380C) (antisense) stab05	AuAAAAcGccGcAGAcAcAT _s T	430
30374	HBV:380U21 siRNA inv stab04 (sense)	B AcuAuuuuGcGGcGucuGuGu B	431
30385	HBV:398L21 siRNA (380C) (antisense) inv stab05	AcAcAGAcGccGcAAAAuAT _s T	432
30353	HBV:413U21 siRNA stab04 (sense)	B uccuGcuGcuAuGccucAucu B	433
30364	HBV:431L21 siRNA (413C) (antisense) stab05	AuGAGGcAuAGcAGcAGGAT _s T	434
30375	HBV:413U21 siRNA inv stab04 (sense)	B ucuAcuccGuAuGucGuccu B	435
30386	HBV:431L21 siRNA (413C) (antisense) inv stab05	AGGAcGAcGAuAcGGAGuAT _s T	436
30354	HBV:462U21 siRNA stab04 (sense)	B uAuGuuGcccGuuuGuuccuu B	437
30365	HBV:480L21 siRNA (462C) (antisense) stab05	AGGAcAAAcGGGcAAAcAuAT _s T	438
30376	HBV:462U21 siRNA inv stab04 (sense)	B ucuccuGuuuGcccGuuGuAu B	439
30387	HBV:480L21 siRNA (462C) (antisense) inv stab05	AuAcAAcGGGcAAAcAGGAT _s T	440

Sirna/ RPI#	Aliases	Sequence	SEQ ID#
30355	HBV:1580U21 siRNA stab04 (sense)	B uGuGcAcuucGcuucAccucu B	441
30366	HBV:1598L21 siRNA (1580C) (antisense) stab05	AGGuGAAAGcGAAAGuGcAcAT _s T	442
30377	HBV:1580U21 siRNA inv stab04 (sense)	B ucuccAcuucGcuucAcGuGu B	443
30388	HBV:1598L21 siRNA (1580C) (antisense) inv stab05	AcAcGuGAAGcGAAGuGGAT _s T	444
30356	HBV:1586U21 siRNA stab04 (sense)	B cuucGcuucAccucuGcAcGu B	445
30367	HBV:1604L21 siRNA (1586C) (antisense) stab05	GuGcAGAGGuGAAGcGAAAGT _s T	446
30378	HBV:1586U21 siRNA inv stab04 (sense)	B uGcAcGucuccAcuucGcuuc B	447
30389	HBV:1604L21 siRNA (1586C) (antisense) inv stab05	GAAcGcGAAAGuGGAGAcGuGT _s T	448
30357	HBV:1780U21 siRNA stab04 (sense)	B AGGcuGuAGGcAuAAuGGu B	449
30368	HBV:1798L21 siRNA (1780C) (antisense) stab05	cAAuuuAuGccuAcAGccuT _s T	450
30379	HBV:1780U21 siRNA inv stab04 (sense)	B uGGuuAAuuAcGGAuGuuGGGA B	451
30390	HBV:1798L21 siRNA (1780C) (antisense) inv stab05	uccGAcAuccGuAuuuuAAcT _s T	452
30612	HBV:1580U21 siRNA stab07 (sense)	B uGuGcAcuucGcuucAccuTT B	453
30620	HBV:1598L21 siRNA (1580C) (antisense) stab08	aggugaagcgaagucacaT _s T	454
30628	HBV:1582U21 siRNA inv stab07 (sense)	B ucuccAcuucGcuucAcGuTT B	455
30636	HBV:1596L21 siRNA (1578C) (antisense) inv stab08	gca caccugaa gcaagugT _s T	456
30612	HBV:1580U21 siRNA stab07 (sense)	B uGuGcAcuucGcuucAccuTT B	457
31175	HBV:1598L21 siRNA (1580C) stab11 (antisense)	AGGuGAAAGcGAAAGuGcAcAT _s T	458
30612	HBV:1580U21 siRNA stab07 (sense)	B uGuGcAcuucGcuucAccuTT B	459
31176	HBV:1596L21 siRNA (1578C) (antisense) inv stab11 (antisense)	GcAcAcGuGAAAGcGAAAGuGT _s T	460
30287	HBV:1580U21 siRNA (sense)	UGUGCACUUCGCUUCACCCUCU	461
30298	HBV:1598L21 siRNA (1580C) (antisense)	AGGUGAAGCGAAGUGGCACACG	462
30355	HBV:1580U21 siRNA stab04 (sense)	B uGuGcAcuucGcuucAccucu B	463
30366	HBV:1598L21 siRNA (1580C) (antisense) stab05	AGGuGAAAGcGAAAGuGcAcAT _s T	464
30612	HBV:1580U21 siRNA stab07 (sense)	B uGuGcAcuucGcuucAccuTT B	465
31175	HBV:1598L21 siRNA (1580C) stab11 (antisense)	AGGuGAAAGcGAAAGuGcAcAT _s T	466
30612	HBV:1580U21 siRNA stab07 (sense)	B uGuGcAcuucGcuucAccuTT B	467
30620	HBV:1598L21 siRNA (1580C) (antisense)	AGGuGAAAGcGAAAGuGcAcAT _s T	468

SiRNA/ RPI#	Aliases	Sequence	SEQ ID#
	stab08		
31335	HBV:1580U21 siRNA stab09 (sense)	B UGUGCACUUCGCUUCACCUUTT B	469
31337	HBV:1598L21 siRNA (1580C) stab10 (antisense)	AGGUGAAGCGAAGUGCACATsT	470
31456	HCVa:291U21 siRNA stab04	B cuuGuGGuAcuGccuGAUATT B	471
31468	HCVa:309L21 siRNA (291C) stab05	uAucAGGcAGuAaccAcAAGTsT	472
31480	HCVa:291U21 siRNA inv stab04	B AuAGuccGucAuGGGuucTT B	473
31492	HCVa:309L21 siRNA (291C) inv stab05	GAAcAccAuGAcGGAcuAuTsT	474
31461	HCVa:300U21 siRNA stab04	B cuGccuGAuAGGGGuGcuuGTT B	475
31473	HCVa:318L21 siRNA (300C) stab05	cAAGcAcccuAucAGGcAGTsT	476
31485	HCVa:300U21 siRNA inv stab04	B GuucGuGGGAuAGuccGucTT B	477
31497	HCVa:318L21 siRNA (300C) inv stab05	GAcGGAcuAuuccAcGAAcTsT	478
31463	HCVa:303U21 siRNA stab04	B ccuGAuAGGGGuGcuuGcGATT B	479
31475	HCVa:321L21 siRNA (303C) stab05	ucGcAAGAcccuAucAGGTsT	480
31487	HCVa:303U21 siRNA inv stab04	B AGcGuucGuGGGAuAGuccTT B	481
31499	HCVa:321L21 siRNA (303C) inv stab05	GGAcuAuuccAcGAAcGcuTsT	482
31344	HCVa:325U21 siRNA stab07	B ccccGGGAGGcucGuAGATT B	483
30562	HCVa:345L21 siRNA (325C) Y-2'F, R-2'OMe + TsT	ucuAcGAGAcucccccGGGTsT	484
31345	HCVa:325U21 siRNA inv stab07	B AGAuGcuuGGAGGGcccccTT B	485
31346	HCVa:343L21 siRNA (325C) inv stab08	GGGcccuccAGAGcAucuTsT	486
31702	HCVa:326U21 siRNA stab07	B cccGGGAGGcucGuAGAcTT B	487
31706	HCVa:344L21 siRNA (326C) stab08	GucuAcGAGAcucccccGGTsT	488
31710	HCVa:326U21 siRNA inv stab07	B cAGAuGcuuGGAGGGcccccTT B	489
31714	HCVa:344L21 siRNA (326C) inv stab08	GGGcccuccAGAGcAucuGTsT	490
31703	HCVa:327U21 siRNA stab07	B ccGGGAGGcucGuAGAcTT B	491
31707	HCVa:345L21 siRNA (327C) stab08	GGucuAcGAGAcucccccGGTsT	492
31711	HCVa:327U21 siRNA inv stab07	B ccAGAuGcuuGGAGGGcccccTT B	493
31715	HCVa:345L21 siRNA (327C) inv stab08	GGcccuccAGAGcAucuGGTsT	494
31704	HCVa:328U21 siRNA stab07	B cGGGAGGcucGuAGAccGTT B	495
31708	HCVa:346L21 siRNA (328C) stab08	cGGucuAcGAGAcucccccGTsT	496
31712	HCVa:328U21 siRNA inv stab07	B GccAGAuGcuuGGAGGGcccccTT B	497
31716	HCVa:346L21 siRNA (328C) inv stab08	GcccuccAGAGcAucuGGcTsT	498
31705	HCVa:329U21 siRNA stab07	B GGGAGGcucGuAGAccGuTT B	499
31709	HCVa:347L21 siRNA (329C) stab08	AcGGucuAcGAGAcucccccTsT	500
31713	HCVa:329U21 siRNA inv stab07	B uGccAGAuGcuuGGAGGGTT B	501
31717	HCVa:347L21 siRNA (329C) inv stab08	cccuccAGAGcAucuGGcATsT	502

Sirna/ RPI#	Aliases	Sequence	SEQ ID#
31703	HCVa:327U21 siRNA stab07	B ccGGGAGGucucGuAGAcTT B	503
31707	HCVa:345L21 siRNA (327C) stab08	GGucAcGAGAcuuccGGTsT	504
31711	HCVa:327U21 siRNA inv stab07	B ccAGAuGcuuGGAGGGccTT B	505
31715	HCVa:345L21 siRNA (327C) inv stab08	GGccuccAGAGcAucuGGTsT	506
29579	HCVa:325U21 siRNA	CCCCGGGAGGUCUCGUAGACCGU	543
	HCVa:327 siRNA 3'-classI 10bp	UCUCGUAGACCUUGGUCUACGAGACCUCCCGGTT	544
	HCVa:327 siRNA 3'-classI 8bp	UCGUAGACCUUGGUCUACGAGACCUCCCGGTT	545
	HCVa:327 siRNA 3'-classI 6bp	GUAGACCUUGGUCUACGAGACCUCCCGGTT	546
	HCVa:327 siRNA 3'-classI 4bp	AGACCUUGGUCUACGAGACCUCCCGGTT	547
	HCVa:327 siRNA 5'-classI 10bp	GGUCUACGAGACCUCCCGGUUCCGGAGGUCU	548
	HCVa:327 siRNA 5'-classI 8bp	GGUCUACGAGACCUCCCGGUUCCGGAGGU	549
	HCVa:327 siRNA 5'-classI 6bp	GGUCUACGAGACCUCCCGGUUCCGGGAG	550
	HCVa:327 siRNA 5'-classI 4bp	GGUCUACGAGACCUCCCGGUUCCGGG	551
	HCVa:327 siRNA 3'-gaaa 10bp	CUCGUAGACCGAAAGGUCUACGAGACCUCCCGGTT	552
	HCVa:327 siRNA 3'-gaaa 8bp	CGUAGACCGAAAGGUCUACGAGACCUCCCGGTT	553
	HCVa:327 siRNA 3'-gaaa 6bp	UAGACCGAAAGGUCUACGAGACCUCCCGGTT	554
	HCVa:327 siRNA 3'-gaaa 4bp	GACCGAAAGGUCUACGAGACCUCCCGGTT	555
	HCVa:327 siRNA 5'-gaaa 10bp	GGUCUACGAGACCUCCCGGUUGAAACCGGAGGUC	556
	HCVa:327 siRNA 5'-gaaa 8bp	GGUCUACGAGACCUCCCGGUUGAAACCGGAGG	557
	HCVa:327 siRNA 5'-gaaa 6bp	GGUCUACGAGACCUCCCGGUUGAAACCGGGA	558
	HCVa:327 siRNA 5'-gaaa 4bp	GGUCUACGAGACCUCCCGGUUGAAACCGG	559
	HCVa:327 siRNA 3'-uuuguguag 10bp	CGUAGACCUUUUUGUGUAGGGUCUACGAGACCUCCCGGTT	560
	HCVa:327 siRNA 3'-uuuguguag 8bp	UAGACCUUUUUGUGUAGGGUCUACGAGACCUCCCGGTT	561
	HCVa:327 siRNA 3'-uuuguguag 6bp	GACCUUUUUGUGUAGGGUCUACGAGACCUCCCGGTT	562
	HCVa:327 siRNA 3'-uuuguguag 4bp	CCUUUUUGUGUAGGGUCUACGAGACCUCCCGGTT	563
	HCVa:327 siRNA 5'-uuuguguag 10bp	GGUCUACGAGACCUCCCGGUUUUUGUGUAGCCGGAGGUC	564
	HCVa:327 siRNA 5'-uuuguguag 8bp	GGUCUACGAGACCUCCCGGUUUUUGUGUAGCCGGAGG	565
	HCVa:327 siRNA 5'-uuuguguag 6bp	GGUCUACGAGACCUCCCGGUUUUUGUGUAGCCGGGA	566
	HCVa:327 siRNA 5'-uuuguguag 4bp	GGUCUACGAGACCUCCCGGUUUUUGUGUAGCCGG	567
	HCVa:327 siRNA 3'-classI 10bp stab08	ucucGuAGAccuuGGucAcGAGAccuuccGGTsT	568
	HCVa:327 siRNA 3'-classI 8bp stab08	ucGuAGAccuuGGucAcGAGAccuuccGGTsT	569
	HCVa:327 siRNA 3'-classI 6bp stab08	GuAGAccuuGGucAcGAGAccuuccGGTsT	570
	HCVa:327 siRNA 3'-classI 4bp stab08	AGAccuuGGucAcGAGAccuuccGGTsT	571
	HCVa:327 siRNA 5'-classI 10bp stab08	GGucAcGAGAccuuccGGuuccGGGAGGucu	572
	HCVa:327 siRNA 5'-classI 8bp stab08	GGucAcGAGAccuuccGGuuccGGGAGGGu	573
	HCVa:327 siRNA 5'-classI 6bp stab08	GGucAcGAGAccuuccGGuuccGGGAGG	574

Sirna/ RPI#	Aliases	Sequence	SEQ ID#
	HCVa:327 siRNA 5'-classI 4bp stab08	GGucuaAcGAGAccucccGGuccGGG	575
	HCVa:327 siRNA 3'-gaaa 10bp stab08	cucGuuAGAccGAAAGGucuaAcGAGAccucccGGTsT	576
	HCVa:327 siRNA 3'-gaaa 8bp stab08	cGuuAGAccGAAAGGucuaAcGAGAccucccGGTsT	577
	HCVa:327 siRNA 3'-gaaa 6bp stab08	uAGAccGAAAGGucuaAcGAGAccucccGGTsT	578
	HCVa:327 siRNA 3'-gaaa 4bp stab08	GAccGAAAGGucuaAcGAGAccucccGGTsT	579
	HCVa:327 siRNA 5'-gaaa 10bp stab08	GGucuaAcGAGAccucccGGuuGAAAccGGGAGGuc	580
	HCVa:327 siRNA 5'-gaaa 8bp stab08	GGucuaAcGAGAccucccGGuuGAAAccGGGAGG	581
	HCVa:327 siRNA 5'-gaaa 6bp stab08	GGucuaAcGAGAccucccGGuuGAAAccGGGA	582
	HCVa:327 siRNA 5'-gaaa 4bp stab08	GGucuaAcGAGAccucccGGuuGAAAccGG	583
	HCVa:327 siRNA 3'-uuuguguag 10bp stab08	cGuuAGAccuuuuuuGuGuAGGGGucuaAcGAGAccucccGGTsT	584
	HCVa:327 siRNA 3'-uuuguguag 8bp stab08	uAGAccuuuuuuGuGuAGGGGucuaAcGAGAccucccGGTsT	585
	HCVa:327 siRNA 3'-uuuguguag 6bp stab08	GAccuuuuuuGuGuuAGGGGucuaAcGAGAccucccGGTsT	586
	HCVa:327 siRNA 3'-uuuguguag 4bp stab08	ccuuuuuuGuGuuAGGGGucuaAcGAGAccucccGGTsT	587
	HCVa:327 siRNA 5'-uuuguguag 10bp stab08	GGucuaAcGAGAccucccGGuuuuuuGuGuuAGccGGGAGGuc	588
	HCVa:327 siRNA 5'-uuuguguag 8bp stab08	GGucuaAcGAGAccucccGGuuuuuuGuGuuAGccGGGAGG	589
	HCVa:327 siRNA 5'-uuuguguag 6bp stab08	GGucuaAcGAGAccucccGGuuuuuuGuGuuAGccGGGA	590
	HCVa:327 siRNA 5'-uuuguguag 4bp stab08	GGucuaAcGAGAccucccGGuuuuuuGuGuuAGccGG	591
	HCVa:347L23 siRNA (327C) stab08	AcGGGucuaAcGAGAccucccGGTsT	592
	HCVa:346L22 siRNA (327C) stab08	cGGGucuaAcGAGAccucccGGTsT	593
	HCVa:345L21 siRNA (327C) stab08	GGucuaAcGAGAccucccGGTsT	594
	HCVa:344L20 siRNA (327C) stab08	GucuaAcGAGAccucccGGTsT	595
	HCVa:343L19 siRNA (327C) stab08	ucuAcGAGAccucccGGTsT	596
	HCVa:342L18 siRNA (327C) stab08	cuAcGAGAccucccGGTsT	597
	HCVa:341L17 siRNA (327C) stab08	uAcGAGAccucccGGTsT	598
	HCVa:340L16 siRNA (327C) stab08	AcGAGAccucccGGTsT	599
	HCVa:339L15 siRNA (327C) stab08	cGAGAccucccGGTsT	600
	HCVa:345L21 siRNA (327C) stab08 GG	GGucuaAcGAGAccucccGGGsG	601
	HCVa:345L20 siRNA (327C) stab08 G	GGucuaAcGAGAccucccGGGsG	602
	HCVa:345L20 siRNA (327C) stab08	GGucuaAcGAGAccucccGGTsT	603
	HCVa:345L19 siRNA (327C) stab08	GGucuaAcGAGAccucccGsG	604
	HCVa:345L18 siRNA (327C) stab08	GGucuaAcGAGAccucccsG	605

Sirna/ RPI#	Aliases	Sequence	SEQ ID#
	HCVa:345L17 siRNA (327C) stab08	GGUcUAcGAGAccucsc	606
	HCVa:345L16 siRNA (327C) stab08	GGUcUAcGAGAccucsc	607
	HCVa:345L15 siRNA (327C) stab08	GGUcUAcGAGAccucsc	608
	HCVa:327U21 siRNA stab07	B ccGGGAGGGuCucGuAGAccTT B	609
	HCVa:327U21 siRNA stab07 GT	B ccGGGAGGGuCucGuAGAccGT B	610
	HCVa:327U21 siRNA stab07	B cGGGAGGGuCucGuAGAccTT B	611
	HCVa:328U20 siRNA stab07	B GGGAGGGuCucGuAGAccTT B	612
	HCVa:329U19 siRNA stab07	B GGAGGGuCucGuAGAccTT B	613
	HCVa:330U18 siRNA stab07	B GAGGGuCucGuAGAccTT B	614
	HCVa:331U17 siRNA stab07	B AGGGuCucGuAGAccTT B	615
	HCVa:332U16 siRNA stab07	B ccGGGAGGGuCucGuAGAccT B	616
	HCVa:327U21 siRNA stab07	B ccGGGAGGGuCucGuAGAcc B	617
	HCVa:327U21 siRNA stab07	B ccGGGAGGGuCucGuAGAc B	618
	HCVa:327U21 siRNA stab07	B ccGGGAGGGuCucGuAGAc B	619
	HCVa:327U21 siRNA stab07	B ccGGGAGGGuCucGuAG B	620
31270	FLT1:349U21 siRNA stab09 sense	B CUGAGUUUAAAAGGCACCCTT B	621
31273	FLT1:367L21 siRNA (349C) stab10 antisense	GGGUGCCUUUUAAACUCAGTst	622
31276	FLT1:349U21 siRNA stab09 inv sense	B CCCACGGAAAAUUUGAGUCTT B	623
	FLT1:367L21 siRNA (349C) stab10 inv antisense	GACUAAAAUUUCCGUGGGTst	624
31679	HBV1598 all RNA sense	AGGUGAAGCGAAGUGCACAuu	625
30287	HBV1598 all RNA antisense	UGUGCACUUCGCUUCACCCUCU	626
31336	HBV:1580U21 siRNA inv stab09 sense	B UCCACUUCGCUUCACCGUGTT B	627
	HBV:1598L21 siRNA (1580C) inv stab10 antisense	ACACGUGAAGCGAAGUGGATst	629
32636	Luc3:80U21 siRNA stab07 sense	B AuAAGGcuAuGAAGAGAUATT B	630
32676	Luc3:98L21 siRNA (80C) stab08 antisense	uAucucuuAuAGccuuAuTst	631
32640	Luc3:237U21 siRNA stab07 sense	B cGuAuGcAGuGAAAAcucuTT B	632
32680	Luc3:255L21 siRNA (237C) stab08 antisense	AGAGuuuuAcuGcAuAcGTst	633
32662	Luc3:1478U21 siRNA stab07 sense	B uGAcGGAAAAAGAGAuGuTT B	634
32702	Luc3:1496L21 siRNA (1478C) stab08 antisense	AcGAucucuuuuuccGucATst	635
32666	Luc3:1544U21 siRNA stab07 sense	B GAGuuGuGuuuGuGGAcGATT B	636
32706	Luc3:1562L21 siRNA (1544C) stab08 antisense	ucGuuccAcAAAcAcAucTst	637
32672	Luc3:1607U21 siRNA stab07 sense	B GAGAGAuuccuAuAAAGGcTT B	638

SiRNA/ RPI#	Aliases	Sequence	SEQ ID#
32712	Luc3:1625L21 siRNA (1607C) stab08 antisense	GccuuuAuGAGGAucucucTsT	639
33139	HCVa:282U21 siRNA stab07 sense	B GcGAAAGGccuuGuGGuAcTT B	640
33179	HCVa:300L21 siRNA (282C) stab08 antisense	GuAccAcAAGGccuuucGcTsT	641
33140	HCVa:283U21 siRNA stab07 sense	B cGAAAGGccuuGuGGuAcuTT B	642
33180	HCVa:301L21 siRNA (283C) stab08 antisense	AGuAccAcAAGGccuuucGTsT	643
33145	HCVa:289U21 siRNA stab07 sense	B GccuuGuGGuAcuGccuGATT B	644
33185	HCVa:307L21 siRNA (289C) stab08 antisense	ucAGGcAGuAccAcAAGGcTsT	645
33149	HCVa:304U21 siRNA stab07 sense	B cuGAuAGGGGuGcuuGcGAGTT B	646
33183	HCVa:304L21 siRNA (286C) stab08 antisense	GGcAGuAccAcAAGGccuuTsT	647
33150	HCVa:305U21 siRNA stab07 sense	B uGAuAGGGGuGcuuGcGAGuTT B	648
33190	HCVa:323L21 siRNA (305C) stab08 antisense	AcucGcAAGcAcccuAucATsT	649
33151	HCVa:307U21 siRNA stab07 sense	B AuAGGGGuGcuuGcGAGuGcTT B	650
33191	HCVa:325L21 siRNA (307C) stab08 antisense	GcAcucGcAAGcAcccuAuTsT	651
33158	HCVa:317U21 siRNA stab07 sense	B uGcGAGuGccccGGGAGGuTT B	652
33187	HCVa:317L21 siRNA (299C) stab08 antisense	AAGcAccuAucAGGcAGuTsT	653
33210	HBV:258U21 siRNA stab07 sense	B GuGGuGGAcuucucucA4uTT B	654
33250	HBV:276L21 siRNA (258C) stab08 antisense	AuuGAGAGAAAGuccAccAcTsT	655
33212	HBV:260U21 siRNA stab07 sense	B GGuGGAcuucucucA4uuuTT B	656
33252	HBV:278L21 siRNA (260C) stab08 antisense	AAAuuGAGAGAAAGuccAccTsT	657
33214	HBV:263U21 siRNA stab07 sense	B GGAcuucucucA4uuuucTT B	658
33254	HBV:281L21 siRNA (263C) stab08 antisense	AGAAAAuuGAGAGAAAGuccTsT	659
32429	HBV:1583U21 siRNA stab07 sense	B GcAcuucGcuucAcccuGTT B	660
32438	HBV:1601L21 siRNA (1583C) stab08 antisense	cAGAGGuGAAGcGAAGuGcTsT	661
33226	HBV:1585U21 siRNA stab07 sense	B AcuucGcuucAcccuGcATT B	662
33266	HBV:1603L21 siRNA (1585C) stab08 antisense	uGcAGAGGuGAAGcGAAGuTsT	663
31651	HBV:1580U21 siRNA stab06 sense	B UGUGcACUUCGCUUCACCCUTT B	664
31652	HBV:1580U21 siRNA inv stab06 sense	B UCCACUUCGCUUCACCGUGUTT B	665

Sirna/ RPI#	Aliases	Sequence	SEQ ID#
31653	HBV:1580U21 siRNA stab16 sense	B UGUGCACUUCGCUUCACCU TT B	666
31654	HBV:1580U21 siRNA inv stab16 sense	B UCCACUUCGCUUCACGUG TT B	667
31657	HBV:1580U21 siRNA stab18 sense	B uGuGcAcuucGcuucAccu TT B	668
31658	HBV:1580U21 siRNA inv stab18 sense	B uccAcuucGcuucAcGuGu TT B	669

UPPER CASE = ribonucleotide

UPPER CASE UNDERLINE = 2'-O-methyl nucleotide

Lowercase = 2'-deoxy-2'-fluoro nucleotide

T = thymidine

T = inverted thymidine

t = 3'-deoxy thymidine

B = inverted deoxyabasic succinate linker

B = inverted deoxyabasic

X = universal base (5-nitroindole)

Z = universal base (3-nitropyrrole)

S = phosphorothioate internucleotide linkage

U = 5-bromodeoxyuridine

A = deoxyadenosine

G = deoxyguanosine

L = glyceryl moiety

ddC = dideoxy Cytidine

p = phosphate

Table II**A. 2.5 μ mol Synthesis Cycle ABI 394 Instrument**

Reagent	Equivalents	Amount	Wait Time* DNA	Wait Time* 2'-O-methyl	Wait Time*RNA
Phosphoramidites	6.5	163 μ L	45 sec	2.5 min	7.5 min
S-Ethyl Tetrazole	23.8	238 μ L	45 sec	2.5 min	7.5 min
Acetic Anhydride	100	233 μ L	5 sec	5 sec	5 sec
N-Methyl Imidazole	186	233 μ L	5 sec	5 sec	5 sec
TCA	176	2.3 mL	21 sec	21 sec	21 sec
Iodine	11.2	1.7 mL	45 sec	45 sec	45 sec
Beaucage	12.9	645 μ L	100 sec	300 sec	300 sec
Acetonitrile	NA	6.67 mL	NA	NA	NA

B. 0.2 μ mol Synthesis Cycle ABI 394 Instrument

Reagent	Equivalents	Amount	Wait Time* DNA	Wait Time* 2'-O-methyl	Wait Time*RNA
Phosphoramidites	15	31 μ L	45 sec	233 sec	465 sec
S-Ethyl Tetrazole	38.7	31 μ L	45 sec	233 min	465 sec
Acetic Anhydride	655	124 μ L	5 sec	5 sec	5 sec
N-Methyl Imidazole	1245	124 μ L	5 sec	5 sec	5 sec
TCA	700	732 μ L	10 sec	10 sec	10 sec
Iodine	20.6	244 μ L	15 sec	15 sec	15 sec
Beaucage	7.7	232 μ L	100 sec	300 sec	300 sec
Acetonitrile	NA	2.64 mL	NA	NA	NA

C. 0.2 μ mol Synthesis Cycle 96 well Instrument

Reagent	Equivalents:DNA/ 2'-O-methyl/Ribo	Amount: DNA/2'-O- methyl/Ribo	Wait Time* DNA	Wait Time* 2'-O- methyl	Wait Time* Ribo
Phosphoramidites	22/33/66	40/60/120 μ L	60 sec	180 sec	360sec
S-Ethyl Tetrazole	70/105/210	40/60/120 μ L	60 sec	180 min	360 sec
Acetic Anhydride	265/265/265	50/50/50 μ L	10 sec	10 sec	10 sec
N-Methyl Imidazole	502/502/502	50/50/50 μ L	10 sec	10 sec	10 sec
TCA	238/475/475	250/500/500 μ L	15 sec	15 sec	15 sec
Iodine	6.8/6.8/6.8	80/80/80 μ L	30 sec	30 sec	30 sec
Beaucage	34/51/51	80/120/120	100 sec	200 sec	200 sec
Acetonitrile	NA	1150/1150/1150 μ L	NA	NA	NA

- Wait time does not include contact time during delivery.
- Tandem synthesis utilizes double coupling of linker molecule

Table III

Group	Solution on Filter (1.0 μ L)	Stock VEGF concentration	Number of Animals	Injectate (6.0 μ L)	Dose	Conc. injectate
1	Tris-Cl pH 6.9	NA	5	water	NA	NA
2	R&D Systems VEGF-carrier free 75 μ M	3.53 μ g/ μ L	5	water	NA	NA
3	R&D Systems VEGF-carrier free 75 μ M	3.53 μ g/ μ L	5	Site 2340 Stab1 siRNA	10 μ g/eye	1.67 μ g/ μ L
4	R&D Systems VEGF-carrier free 75 μ M	3.53 μ g/ μ L	5	Site 2340 Stab1 siRNA	3 μ g/eye	0.5 μ g/ μ L
5	R&D Systems VEGF-carrier free 75 μ M	3.53 μ g/ μ L	5	Site 2340 Stab1 siRNA	1 μ g/eye	0.167 μ g/ μ L
6	R&D Systems VEGF-carrier free 75 μ M	3.53 μ g/ μ L	5	Inactive Site 2340 Stab1 siRNA	10 μ g/eye	1.67 μ g/ μ L
7	R&D Systems VEGF-carrier free 75 μ M	3.53 μ g/ μ L	5	Inactive Site 2340 Stab1 siRNA	3 μ g/eye	0.5 μ g/ μ L
8	R&D Systems VEGF-carrier free 75 μ M	3.53 μ g/ μ L	5	Inactive Site 2340 Stab1 siRNA	1 μ g/eye	0.167 μ g/ μ L

Table IV

Non-limiting examples of Stabilization Chemistries for chemically modified siNA constructs

Chemistry	pyrimidine	Purine	cap	p=S	Strand
“Stab 1”	Ribo	Ribo	-	5 at 5'-end 1 at 3'-end	S/AS
“Stab 2”	Ribo	Ribo	-	All linkages	Usually AS
“Stab 3”	2'-fluoro	Ribo	-	4 at 5'-end 4 at 3'-end	Usually S
“Stab 4”	2'-fluoro	Ribo	5' and 3'-ends	-	Usually S
“Stab 5”	2'-fluoro	Ribo	-	1 at 3'-end	Usually AS
“Stab 6”	2'-O-Methyl	Ribo	5' and 3'-ends	-	Usually S
“Stab 7”	2'-fluoro	2'-deoxy	5' and 3'-ends	-	Usually S
“Stab 8”	2'-fluoro	2'-O-Methyl	-	1 at 3'-end	S or AS
“Stab 9”	Ribo	Ribo	5' and 3'-ends	-	Usually S
“Stab 10”	Ribo	Ribo	-	1 at 3'-end	Usually AS
“Stab 11”	2'-fluoro	2'-deoxy	-	1 at 3'-end	Usually AS
Stab 12	2'-fluoro	LNA	5' and 3'-ends		Usually S
“Stab 13”	2'-fluoro	LNA		1 at 3'-end	Usually AS
“Stab 14”	2'-fluoro	2'-deoxy		2 at 5'-end 1 at 3'-end	Usually AS
“Stab 15”	2'-deoxy	2'-deoxy		2 at 5'-end 1 at 3'-end	Usually AS
“Stab 16”	Ribo	2'-O-Methyl	5' and 3'-ends		Usually S
“Stab 17”	2'-O-Methyl	2'-O-Methyl	5' and 3'-ends		Usually S
“Stab 18”	2'-fluoro	2'-O-Methyl		1 at 3'-end	Usually AS

5 CAP = any terminal cap, see for example **Figure 22**.

All Stab 1-18 chemistries can comprise 3'-terminal thymidine (TT) residues

All Stab 1-18 chemistries typically comprise 21 nucleotides, but can vary as described herein.

S = sense strand

10 AS = antisense strand

Table V: Peptides for Conjugation

Peptide	Sequence	SEQ ID NO
ANTENNAPEDIA	RQI KIW FQN RRM KWK K amide	507
Kaposi fibroblast growth factor	AAV ALL PAV LLA LLA P + VQR KRQ KLMP	508
<i>caiman crocodylus</i> Ig(5) light chain	MGL GLH LLV LAA ALQ GA	509
HIV envelope glycoprotein gp41	GAL FLG FLG AAG STM GA + PKS KRK 5 (NLS of the SV40)	510
HIV-1 Tat	RKK RRQ RRR	511
Influenza hemagglutinin envelop glycoprotein	GLFEAIAGFIENGWEGMIDGGGYC	512
RGD peptide	X-RGD-X where X is any amino acid or peptide	513
transportan A	GWT LNS AGY LLG KIN LKA LAA LAK KIL	514
Somatostatin (tyr-3-octreotate)	(S) <i>FC YWK</i> TCT	515
Pre-S-peptide	(S)DH QLN PAF	516

(S) optional Serine for coupling

5 *Italic* = optional D isomer for stability

Table VI: Duplex half-lives in human and mouse serum and liver extracts

Stability S/AS Sirna #	All RNA 47715/47933	4*/5 30355/30366	4/5* 30355/30366	7/11* 30612/31175	7*/8 30612/30620	7/8* 30612/30620
Human Serum t ½ hours	0.017	408 (0.96) [†]	39 (0.65)	54 (0.76)	130 (0.88)	94 (0.86)
Human Liver t ½ hours	2.5	28.6 (0.40)	43.5 (0.66)	0.78 / 2.9 [†] (0.45)	9 (0.39)	816 (0.99)
Mouse Serum t ½ hours	1.17	16.7 (0.9)	10 (0.81)	2.3	16.6 (0.46)	35.7 (0.69)
Mouse Liver t ½ hours	6	1.08	0.80	0.20	0.22	120 (0.89)

* The asterisk designates the strand carrying the radiolabel in the duplex.

[†] For longer half-lives the fraction full-length at the 18 hours is presented as the parenthetic lower number in each cell.

[‡] A biphasic curve was observed, half-lives for both phases are shown.

Table VII: Single strand half-lives in human serum

Stability Sirna #	4 30355	5 30366	7 30612	11 31175	8 30620
Human serum $t_{1/2}$ hours	22	16	13	19	28
Human liver $t_{1/2}$ hours	0.92	0.40	0.43	0.27	192

5

Table VIII. Human serum half-lives for Stab 4/5 duplex chemistry with terminus chemistries of figure 22

Cap Chemistry	2 (R=O) (B=T)	7 (R=O) (B=T)	9 (R=O) (B=T)	2 (R=S) (B=T)	8 (R=O) (B=T)	1 (R=O) (B=T)	3 (R=O) (B=T)	6 (R=O) (B=T)
Human Serum $t_{1/2}$ hours	1	1.2	2.3	39	96 (0.69) [†]	460 (0.95)	770 (0.94)	770 (0.95)

5

The capping structures were in the following position of the 4:5 chemistry formatted sequence:

antisense strand – 5'-uuGuuGuAuuuuGuGGuuG– CAP – 3' where CAP is 1, 2, 3, 6, 7, 8, or 9 from Figure 22. (SEQ ID NO: 670)

10 sense strand 5'-CAP- cAAccAcAAAAuAcAAcAATT– CAP – 3' where CAP is 1 from Figure 22. (SEQ ID NO: 671)

[†] For half-lives that extend beyond the time course sampled the fraction full-length is presented in parentheses.

CLAIMS

What we claim is:


1. A double stranded short interfering nucleic acid (siNA) molecule that comprises a first nucleotide sequence complementary to a target RNA sequence or a portion thereof, and a second sequence having complementarity to said first sequence, wherein said second sequence is chemically modified in a manner that said second sequence can no longer act as a guide sequence for mediating RNA interference.
2. The siNA molecule of claim 1, wherein said second sequence does not have a terminal 5'-hydroxyl (5'-OH) or 5'-phosphate group.
3. The siNA molecule of claim 1, wherein said second sequence comprises a terminal cap moiety at the 5'-end of said second sequence.
4. The siNA molecule of claim 1, wherein said second sequence comprises a terminal cap moiety at the 5'-end and 3'-end of said second sequence.
5. The siNA molecule of claim 3, wherein said terminal cap moiety is an inverted deoxyabasic moiety.
6. The siNA molecule of claim 4, wherein said terminal cap moiety is an inverted deoxyabasic moiety.
7. The siNA molecule of claim 1, wherein said siNA molecule comprises no ribonucleotides.
8. The siNA molecule of claim 1, wherein said siNA molecule comprises ribonucleotides.
9. The siNA molecule of claim 1, wherein said first sequence and said second sequence each comprise about 19 to about 23 nucleotides, and wherein each said sequence comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand.
10. The siNA molecule of claim 1, wherein any pyrimidine nucleotides in said second sequence are 2'-O-methyl pyrimidine nucleotides.
11. The siNA molecule of claim 1, wherein any purine nucleotides in said second sequence are 2'-deoxy purine nucleotides.

12. The siNA molecule of claim 1, wherein any pyrimidine nucleotides in said second sequence are 2'-deoxy-2'-fluoro pyrimidine nucleotides.
13. The siNA molecule of claim 1, wherein any pyrimidine nucleotides in said first sequence are 2'-deoxy-2'-fluoro pyrimidine nucleotides
14. The siNA molecule of claim 1, wherein any purine nucleotides in said first sequence are 2'-O-methyl purine nucleotides.
15. The siNA molecule of claim 1, wherein any purine nucleotides in said first sequence comprise 2'-deoxy- purine nucleotides.
16. The siNA molecule of claim 1, wherein said first sequence comprises a phosphorothioate internucleotide linkage at the 3' end of said first sequence.
17. A pharmaceutical composition comprising the siNA molecule of claim 1 in an acceptable carrier or diluent.

ABSTRACT OF THE DISCLOSURE

The present invention concerns methods and reagents useful in modulating gene expression in a variety of applications, including use in therapeutic, diagnostic, target validation, and genomic discovery applications. Specifically, the invention relates to synthetic chemically modified small nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules capable of mediating RNA interference (RNAi) against target nucleic acid sequences. The small nucleic acid molecules are useful in the treatment of any disease or condition that responds to modulation of gene expression or activity in a cell, tissue, or organism.


Figure 1




= SOLID SUPPORT

R = TERMINAL PROTECTING GROUP
 FOR EXAMPLE:
 DIMETHOXYTRITYL (DMT)

(1) = CLEAVABLE LINKER
 (FOR EXAMPLE: NUCLEOTIDE SUCCINATE OR
 INVERTED DEOXYABASIC SUCCINATE)
 (2) = CLEAVABLE LINKER
 (FOR EXAMPLE: NUCLEOTIDE SUCCINATE OR
 INVERTED DEOXYABASIC SUCCINATE)



INVERTED DEOXYABASIC SUCCINATE
 LINKAGE



GLYCERYL SUCCINATE LINKAGE

Figure 2




Figure 3

5'-CGUACGCGGAUACUUCGATT (SEQ ID NO: 394) 3'-TTGCAUGCGCCUUAUGAAGCU (SEQ ID NO: 395)	$T_{1/2} = 15$ seconds (control)
5'-B cAAccAcAAAAuAcAAcAATT B (SEQ ID NO: 396) 3'-TXGuuGGuGuuuuAuGuuGuu (SEQ ID NO: 397)	$T_{1/2} = 138$ min
5'-B cAAccAcAAAAuAcAAcAATT B (SEQ ID NO: 396) 3'-TDGuuGGuGuuuuAuGuuGuu (SEQ ID NO: 398)	$T_{1/2} = 3.7$ days
5'-B cAAccAcAAAAuAcAAcAATT B (SEQ ID NO: 396) 3'-XTGuuGGuGuuuuAuGuuGuu (SEQ ID NO: 399)	$T_{1/2} = 72$ minutes
5'-B cAAccAcAAAAuAcAAcAATT B (SEQ ID NO: 396) 3'-LTGuuGGuGuuuuAuGuuGuu (SEQ ID NO: 400)	$T_{1/2} = 40$ days
5'-B cAAccAcAAAAuAcAAcAATT B (SEQ ID NO: 396) 3'-tTGuuGGuGuuuuAuGuuGuu (SEQ ID NO: 401)	$T_{1/2} = 32$ days

G, A, U, C = Guanosine, Adenosine, Uridine, Cytidine

T = Thymidine

Lower Case = 2'-deoxy-2'-fluoro

S = phosphorothioate

B = inverted deoxyabasic

D = inverted Thymidine

X = 3'-deoxy Thymidine

t = L-thymidine

L = Glyceryl moiety




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17






Figure 18



POSITIONS (NN) CAN COMPRISE ANY NUCLEOTIDE, SUCH AS DEOXYNUCLEOTIDES (eg. THYMIDINE) OR UNIVERSAL BASES
B = ABASIC, INVERTED ABASIC, INVERTED NUCLEOTIDE OR OTHER TERMINAL CAP THAT IS OPTIONALLY PRESENT
L = GLYCERYL MOIETY THAT IS OPTIONALLY PRESENT
S = PHOSPHOROTHIOATE OR PHOSPHORODITHIOATE

Figure 19



lower case = 2'-O-Methyl or 2'-deoxy-2'-fluoro
 italic lower case = 2'-deoxy-2'-fluoro
underline = 2'-O-methyl

ITALIC UPPER CASE = DEOXY
 B = INVERTED DEOXYABASIC
 L = GLYCERYL MOIETY OPTIONALLY PRESENT
 S = PHOSPHOROTHIOATE OR
 PHOSPHORODITHIOATE

Figure 21: Target site Selection using siRNA






Figure 22




R = O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, or aralkyl
 B = Independently any nucleotide base, either naturally occurring or chemically modified, or optionally H (abasic).

**Figure 23: Inhibition of VEGF-Induced Angiogenesis
by siRNAs**



**Figure 24: Stab4/5 siNA Targeted to HBV:
 HBsAg Levels in Hep G2 Cells**



**Figure 25: Dose Response with Stab4/5 siRNAs Targeted to
 HBV Sites 262 & 1580**




Figure 26: Comparison of Stab7/8 and Stab 7/11 siRNAs Targeted to HBV RNA Site 1580




Figure 27: Modification Strategy




Figure 28: Duration of siRNA Effect
All-Ribo vs. Stab4/5 HBV Site 1580: HBsAg Levels




Figure 29: Duration of siRNA Effect
 All-Ribo vs. Stab7/8 HBV Site 1580: HBsAg Levels




Figure 30: Duration of siRNA Effect
 All-Ribo vs. Stab7/11 HBV Site 1580: HBsAg Levels




Figure 31: Duration of siRNA Effect
 All-Ribo vs. Stab9/10 HBV Site 1580: HBsAg Levels




Figure 32 : siRNAs targeting HCV chimera







Figure 33: HCV siRNA dose response




**Figure 34: Chemically Modified siRNA targeting
HCV chimera**



**Figure 35: Chemically Modified siRNA targeting
HCV chimera**




**Figure 36: Chemically Modified siRNA
 targeting HCV chimera**



**Figure 37: Chemically Modified siRNA
 targeting HCV chimera**

HCV/PV#280-siRNA to HCV-Luc site 325/345



**Figure 38: HCV/Replicon Cells transfected
 with 0.5µl/well LFA 2K-72 hours**




Figure 39: Dose Response with Stab4/5 siNA Leads in HCV Subgenomic Replicon





Figure 40: Activity of Stab 7/8 siNA Leads in HCV Subgenomic Replicon



**Figure 41: Dose Response with Fully Modified
HCV Site 327 siNA**




Figure 42: Solid Phase Post-synthetic conjugation of pterioic acid






Figure 43



NA = siNA or a portion thereof
 p = phosphorous moiety

Figure 44



NA = siNA or a portion thereof
 p = phosphorous moiety



Figure 45: Solid Phase Post-synthetic conjugation of pterioic acid

Figure 46: Synthesis of N-acetyl-D-galactosamine-2'-aminouridine



Reagents and Conditions: (i) diethylamine, DMF, (ii) **8**, diisopropylethylamine, DMF, (iii) 2-cyanoethyl N,N-diisopropylchlorophosphoramidite, 1-methylimidazole, DIPEA, CH_2Cl_2 , (iv) Ac_2O , TEA, CH_3CN , (v) HCl , Ac_2O , (vi) $\text{Hg}(\text{CN})_2$, MS 4A, CH_3NO_2 -toluene 1:1, (vii) H_2 , 5% Pd-C, ethanol, (viii) N-hydroxysuccinimide, DCC, THF.

D-threoninol

i

11

ii


MMTro

iii

12 R = H
 13 R = $\text{NiPr}_2\text{POCH}_2\text{CH}_2\text{CN}$

Reagents and Conditions: (i) 7, DCC, *N*-hydroxysuccinimide, (ii) MMTr-Cl, pyridine, (iii) 2-cyanoethyl *N,N*-diisopropylchlorophosphoramidite, 1-methylimidazole, DIPEA, CH₂Cl₂.

**Figure 48: Conjugation of targeting ligands to the 5'-end
of a siNA molecule**



N-acetyl-D-galactosamine conjugate

Figure 49: Synthesis of dodecanoic acid linker




Figure 50: Oxime linked siNA/Peptide Conjugate





Figure 51: siNA/Phospholipid Conjugates



PEG=polyethylene glycol

CL=cleavable linker (e.g. A-dT, C-dT)

siNA= short interfering nucleic acid molecule or a portion thereof

Figure 52: siNA Phospholipid Conjugate






Figure 53: siNA-NAcGalactosamine post-synthetic coupling



FOR EXAMPLE: OLIGO-LINKER =



Where n is an integer from 1 to 20

3'-siNA-CL-5' (SENSE OR ANTISENSE STRAND/REGION) + 3'-siNA-CL-5'-O-P(=O)(O⁻)-O-CH₂-ODMT → COUPLING → 3'-siNA-CL-5'-O-P(=O)(O⁻)-O-CH₂-OH + DETRIT

3'-siNA-CL-5'-O-P(=O)(O⁻)-O-CH₂-OH + (Pri)₂N-P(=O)(O⁻)-O-CH₂-CH₂-N≡C → COUPLING → 3'-siNA-CL-5'-O-P(=O)(O⁻)-O-CH₂-O-CH₂-ODMT + (Pri)₂N-P(=O)(O⁻)-O-CH₂-CH₂-N≡C

3'-siNA-CL-5'-O-P(=O)(O⁻)-O-CH₂-O-CH₂-ODMT → CLEAVAGE/DEPROTECTION/PURIFICATION → siNA-CONJUGATE

CL = CLEAVABLE LINKER, E.G. ADENOSINE-THYMIDINE DIMER THAT IS OPTIONALLY PRESENT

Chemical reaction scheme showing the deprotection of a 5'-siRNA-CL-O-protected siRNA. The starting material is a 5'-siRNA-CL-O-protected siRNA with a 5'-(5'-siRNA-CL-O)-3'-(5'-siRNA-CL-O) structure. The reaction is labeled "DEPROTECTION". The product is a 5'-siRNA-CL-O-protected siRNA with a 5'-(5'-siRNA-CL-O)-3'-(5'-siRNA-CL-O) structure, where the 5'-siRNA-CL-O group is deprotected to a 5'-siRNA-CL-O group. The reaction is shown with a chemical structure of the starting material and a chemical structure of the product, with a chemical structure of the 5'-siRNA-CL-O group shown in brackets.

CL = CLEAVABLE LINKER, E.G. ADENOSINE-THYMIDINE DIMER THAT IS OPTIONALLY PRESENT

Figure 56: siNA 3'-Cholesterol Conjugate





Figure 57: Nucleic Acid Cholesterol Conjugates



PEG=polyethylene glycol


CL=cleavable linker (e.g. A-dT, C-dT)

siNA= short interfering nucleic acid molecule or a portion thereof

CHOL=cholesterol or an analog or metabolite thereof


W= linker molecule (see for example Formulae 109 or 112)

Figure 58: siNA Cholesterol Conjugates




CL=cleavable linker (e.g. A-dT, C-dT) that is optionally present
 CHOL=cholesterol or an analog or metabolite thereof
 W= linker molecule (see for example Formulae 107, 108, 109 or 115)
 n = integer, e.g. 1, 2, or 3

Figure 59: siNA Cholesterol Conjugates




CL=cleavable linker (e.g. A-dT, C-dT) that is optionally present
 CHOL=cholesterol or an analog or metabolite thereof
 W= linker molecule (see for example Formulae 107, 108, 109 or 115)
 n = integer, e.g. 1, 2, or 3

Figure 60: siNA Cholesterol Conjugates




CL=cleavable linker (e.g. A-dT, C-dT) that is optionally present
 CHOL=cholesterol or an analog or metabolite thereof
 W= linker molecule (see for example Formulae 107, 108, 109 or 112)
 n = integer, e.g. 1, 2, or 3
 N=integer, e.g. 1, 2, 3, or 4

Figure 61: siNA Lipid Conjugates




CL=cleavable linker (e.g. A-dT, C-dT) that is optionally present
 Lipid=Straight chain or branched alkyl or fatty acid, e.g. C₁₈H₃₇
 W= linker molecule (see for example Formulae 48, 49, 64, or 65)
 n = integer, e.g. 1, 2, or 3

Figure 62: siNA Lipid Conjugates



CL=cleavable linker (e.g. A-dT, C-dT) that is optionally present
 Lipid=Straight chain or branched alkyl or fatty acid, e.g. C₁₈H₃₇
 W= linker molecule (see for example Formulae 48, 49, 64, or 65)
 n = integer, e.g. 1, 2, or 3
 N=integer, e.g. 1, 2, 3, or 4

Figure 63: siNA Galactosamine Conjugates




CL=cleavable linker (e.g. A-dT, C-dT) that is optionally present

GAL=GALACTOSAMINE; e.g. compounds having Formulae 51-56, 86, 92, 99, 100, 103, 105, 106

W= linker molecule (see for example Formulae 102 or 103)

n = integer, e.g. 1, 2, or 3

Figure 64: siNA Galactosamine Conjugates



CL=cleavable linker (e.g. A-dT, C-dT) that is optionally present
 GAL=GALACTOSAMINE; e.g. compounds having Formulae 51-56, 86, 92, 99, 100, 103, 105, 106
 W= linker molecule (see for example Formulae 102 or 103)
 n = integer, e.g. 1, 2, or 3
 N=integer, e.g. 1, 2, 3, or 4

5'-CONJ-W-CL $\frac{\text{SENSE}}{\left(\frac{\text{|||||||||||||||||||||}{\text{ANTISENSE}} \right)_n}$

3' $\left(\frac{\text{|||||||||||||||||||||}{\text{ANTISENSE}} \right)_n$ CL-W-CONJ

5'-CONJ-W-CL $\frac{\text{SENSE}}{\left(\frac{\text{|||||||||||||||||||||}{\text{ANTISENSE}} \right)_n}$ CL-W-CONJ

3' $\left(\frac{\text{|||||||||||||||||||||}{\text{ANTISENSE}} \right)_n$

3'-CONJ-W-CL- $\left(\frac{\text{SENSE}}{\left(\frac{\text{|||||||||||||||||||||}{\text{ANTISENSE}} \right)_n} \right)_n$


3'-CONJ-W-CL- $\left(\frac{\text{SENSE}}{\left(\frac{\text{|||||||||||||||||||||}{\text{ANTISENSE}} \right)_n} \right)_n$ CL-W-CONJ

5'-CONJ-W-CL $\frac{\text{SENSE}}{\left(\frac{\text{|||||||||||||||||||||}{\text{ANTISENSE}} \right)_n}$ CL-W-CONJ

3'-CONJ-W-CL- $\left(\frac{\text{|||||||||||||||||||||}{\text{ANTISENSE}} \right)_n$

n = integer, e.g. 1, 2, or 3

Figure 66: Generalized siNA Conjugate design



CONJ=any biologically active molecule or conjugate as described herein

CL=cleavable linker (e.g. A-dT, C-dT) that is optionally present

W= linker molecule

n = integer, e.g. 1, 2, or 3

N=integer, e.g. 1, 2, 3, or 4

Figure 67: Distribution of Intact siNA in Liver After SC Administration of Conjugated or Unconjugated Chemistries




Figure 68: Lipid Free Delivery of HBV siNA Conjugates in Cell Culture




Figure 69: Scale-up of “mono” Galactosamine phosphoramidite




Figure 70: Synthesis of "tri" Galactosamine phosphoramidite




Figure 71: Synthesis of another Tri-Galactosamine Conjugate




Figure 72: Alternate Synthesis of Tri-Galactosamine Conjugate




Figure 73: Synthesis of NHS Cholesterol Conjugate






Figure 74: Phosphorylated siNA constructs



Phosphates can be modified
as described herein



Asymmetric hairpin



Asymmetric duplex
siNA



(n) = number of base
pairs (e.g. 3-18 bp)

[illegible]

Figure 76: siNA Targeting VEGFR-1 Inhibits VEGF-Induced Rat Corneal Angiogenesis




Figure 77: Duration of Effect of Modified siNA Constructs




Figure 77: Duration of Effect of Modified siNA Constructs

HBV siRNA Duration: Day 9

B




Figure 77: Duration of Effect of Modified siNA Constructs

HBV siRNA Duration: Day 21

C




Figure 77: Duration of Effect of Modified siNA Constructs




Figure 77: Duration of Effect of Modified siNA Constructs

E




Figure 77: Duration of Effect of Modified siNA Constructs




Figure 78: Phosphorylated siNA constructs



Phosphates can be modified
 as described herein



Asymmetric hairpin



Asymmetric duplex
 siNA



(n) = number of base
 pairs (e.g. 3-18 bp)

Figure 79: 5'-phosphate modifications






Figure 80: Serum HBV DNA in Mice Treated with siNA Via HDI



**Figure 81: Serum HBsAg in Mice Treated
with siNA Via HDI**



**Figure 82: Liver HBV RNA in Mice Treated
 with siNA Via HDI**






Figure 83: Serum HBV DNA in Mice Treated with siNA Via HDI 5 and 7 days post treatment



**Figure 84: Luciferase Dose Response
 of select active siNA constructs**



**Figure 85: Luciferase Dose Response
 of select active siNA constructs**




Figure 86: Activity of Stab 7/8 Stabilized siNAs in HCV Replicon




Figure 87: Activity of Stabilized 7/8 siNAs Against HBV in HepG2 Cells




Figure 88: HBV/siNA to site 1580 Combination Constructs




Figure 89: HBV/siNA to site 1580 Combination Constructs





Figure 90: HBV/siNA to site 1580 Combination Constructs



Application Data Sheet

Application Information

Application number::	To be Assigned
Filing Date::	November 24, 2003
Application Type::	Regular
Subject Matter::	Utility
Suggested Classification::	
Suggested Group Art Unit::	
CD-Rom or CR-R?::	None
Number of CD disks::	None
Number of copies of CDs::	None
Sequence submission?::	None
Computer Readable Form (CRF)?::	No
Number of copies of CRF::	No
Title::	RNA Interference Mediated Inhibition of Gene Expression Using Chemically Modified Short Interfering Nucleic Acid (siNA)
Attorney Docket Number::	03-465-B (400/138)
Request for Early Publication?::	No
Request for Non-Publication?::	No
Suggested Drawing Figure::	95
Total Drawing Sheets::	95
Small Entity?::	Yes
Petition Included?::	No
Petition Type::	
Licensed US Govt. Agency::	

Contract or Grant Numbers::
Secrecy Order in Parent Appl.?::

Applicant Information

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: James
Middle Name::
Family Name:: McSwiggen
Name Suffix::
City of Residence:: Boulder
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 4866 Franklin Drive
City of mailing address:: Boulder
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80301

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: Bharat
Middle Name:: M
Family Name:: Chowrira
Name Suffix::
City of Residence:: Louisville

State or Province of Residence::	Colorado
Country of Residence::	United States of America
Street of mailing address::	576 Manorwood Lane
City of mailing address::	Louisville
State or Providence of mailing address::	Colorado
Postal or Zip Code of mailing address::	80027
Applicant Authority type::	Inventor
Primary Citizenship Country::	US
Status::	Full Capacity
Given Name::	Leonid
Middle Name::	
Family Name::	Beigleman
Name Suffix::	
City of Residence::	Longmont
State or Province of Residence::	Colorado
Country of Residence::	United States of America
Street of mailing address::	5530 Colt Drive
City of mailing address::	Longmont
State or Providence of mailing address::	Colorado
Postal or Zip Code of mailing address::	80503
Applicant Authority type::	Inventor
Primary Citizenship Country::	US
Status::	Full Capacity
Given Name::	Dennis
Middle Name::	
Family Name::	Macejak

Name Suffix::
City of Residence:: Arvada
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 6595 Union Street
City of mailing address:: Arvada
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80004

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: Shawn

Middle Name::

Family Name:: Zinnen
Name Suffix::

City of Residence:: Denver
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 2378 Birch Street
City of mailing address:: Denver
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80207

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity

Given Name:: Pamela
Middle Name::
Family Name:: Pavco
Name Suffix::
City of Residence:: Lafayette
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 705 Bayberry Circle
City of mailing address:: Lafayette
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80026

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: Peter
Middle Name::
Family Name:: Haeberli
Name Suffix::
City of Residence:: Berthoud
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 705 7th Street
City of mailing address:: Berthoud
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80513

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: David
Middle Name::
Family Name:: Morrisey
Name Suffix::
City of Residence:: Boulder
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 4769 Tanglewood Trail
City of mailing address:: Boulder
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80301

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: Fosnaugh
Middle Name::
Family Name:: Kathy
Name Suffix::
City of Residence:: Boulder
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 1030 Edinboro Drive
City of mailing address:: Boulder

State or Providence of mailing address:: Colorado

Postal or Zip Code of mailing address:: 80301

Applicant Authority type:: Inventor

Primary Citizenship Country:: US

Status:: Full Capacity

Given Name:: Jamison

Middle Name::

Family Name:: Sharon

Name Suffix::

City of Residence:: Boulder

State or Province of Residence:: Colorado

Country of Residence:: United States of America

Street of mailing address:: 4985 Twin Lakes Road #89

City of mailing address:: Boulder

State or Providence of mailing address:: Colorado

Postal or Zip Code of mailing address:: 80301

Applicant Authority type:: Inventor

Primary Citizenship Country:: US

Status:: Full Capacity

Given Name:: Usman

Middle Name::

Family Name:: Nassim

Name Suffix::

City of Residence:: Lafayette

State or Province of Residence:: Colorado

Country of Residence:: United States of America
Street of mailing address:: 2129 Night Sky Line
City of mailing address:: Lafayette
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80026

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: Thompson
Middle Name::
Family Name:: James
Name Suffix::

City of Residence:: Lafayette
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 705 Barberry Circle
City of mailing address:: Lafayette
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80026

Applicant Authority type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: Vargeese
Middle Name::
Family Name:: Chandra

Name Suffix::
City of Residence:: Broomfield
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 14010 Westhampton Place
City of mailing address:: Broomfield
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80020

Applicant Authority type:: Inventor
Primary Citizenship Country:: China
Status:: Full Capacity
Given Name:: Wang
Middle Name::
Family Name:: Weimen
Name Suffix::
City of Residence:: Superior
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 3122 Cimarron Place
City of mailing address:: Superior
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80027

Applicant Authority type:: Inventor
Primary Citizenship Country:: China
Status:: Full Capacity
Given Name:: Tongqian

Middle Name::
Family Name:: Chen
Name Suffix::
City of Residence:: Longmont
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 2450 Airport Road, Apt. 1284
City of mailing address:: Longmont
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80503

Applicant Authority type:: Inventor
Primary Citizenship Country:: India
Status:: Full Capacity
Given Name:: Narendra
Middle Name::
Family Name:: Vaish
Name Suffix::
City of Residence:: Denver
State or Province of Residence:: Colorado
Country of Residence:: United States of America
Street of mailing address:: 1313 Williams Street #503
City of mailing address:: Denver
State or Providence of mailing address:: Colorado
Postal or Zip Code of mailing address:: 80218

Correspondence Information

Correspondence Customer Number:: 020306
Phone Number:: (312) 913-0001
Fax Number:: (312) 913-0002
E-Mail Address:: docketing@mbhb.com

Representative Information

Representative Customer Number::	020306
----------------------------------	--------

Domestic Priority Information

Application::	Continuity Type::	Parent Application::	Parent Filing Date::
This application	Continuation-in-part	10/693,059	10/23/03
10/693,059	Continuation-in-part	10/444,853	05/23/03
10/444,853	Continuation-in-part	10/652,791	08/29/03
10/652,791	Continuation of	10/422,704	04/24/03
10/422,704	Continuation of	10/417,012	04/16/03
10/693,059	Continuation-in-part	PCT/US03/05346	02/20/03
10/693,059	Continuation-in-part	PCT/US03/05028	02/20/03
PCT/US03/05028	An application claiming the benefit under 35 USC 119(e)	60/358,580	02/20/03
PCT/US03/05028	An application claiming the benefit under 35 USC 119(e)	60/363,124	03/11/02
PCT/US03/05028	An application claiming the benefit under 35 USC 119(e)	60/386,782	06/06/02
PCT/US03/05028	An application claiming the benefit under 35 USC 119(e)	60/406,784	08/29/02
PCT/US03/05028	An application claiming	60/408,378	09/05/02

	the benefit under 35 USC 119(e)		
PCT/US03/05028	An application claiming the benefit under 35 USC 119(e)	60/409,293	09/09/02
PCT/US03/05028	An application claiming the benefit under 35 USC 119(e)	60/440,129	01/15/03
PCT/US03/05346	An application claiming the benefit under 35 USC 119(e)	60/358,580	02/20/03
PCT/US03/05346	An application claiming the benefit under 35 USC 119(e)	60/363,124	03/11/02
PCT/US03/05346	An application claiming the benefit under 35 USC 119(e)	60/386,782	06/06/02
PCT/US03/05346	An application claiming the benefit under 35 USC 119(e)	60/406,784	08/29/02
PCT/US03/05346	An application claiming the benefit under 35 USC 119(e)	60/408,378	09/05/02
PCT/US03/05346	An application claiming the benefit under 35 USC 119(e)	60/409,293	09/09/02
PCT/US03/05346	An application claiming the benefit under 35 USC 119(e)	60/440,129	01/15/03
10/693,095	Continuation-in-part	10/427,160	04/30/03
10/693,095	Continuation-in-part	PCT/US02/15876	5/17/02